

**KIPS**  
ENTRY TESTS  
**SERIES**

**PRACTICE BOOK**  
**KIPSSATs**  
*Self Assessment Tests*

# PHYSICS

**MDCAT**

- ▶ 1693 Practice MCQs
- ▶ Questions from Past Papers
- ▶ Answer Keys with Explanatory Notes
- ▶ Topic-wise Practice Exercises
- ▶ Unit-wise Self Assessment Tests
- ▶ Pre-Assessment Test (Diagnostic Test)
- ▶ Post-Assessment Test  
(Sample Paper as per Original Format)



A Kitab Dost Publication

# CONTENT

TOPIC NO.	PRACTICES EXERCISE	PAGE NO.
A	Pre- Prep Assessment .....	1
1	Force and Motion.....	1
2	Work and Energy.....	19
3	Rotational and Circular motion.....	37
4	Waves.....	51
5	Thermodynamics.....	68
6	Electrostatics.....	78
7	Current Electricity.....	96
8	Electromagnetism.....	109
9	Electromagnetic Induction.....	120
10	Electronics.....	130
11	Dawn of Modern Physics.....	137
12	Atomic Spectra.....	147
13	Nuclear Physics.....	153

UNIT NO.	KIPS SATs UNIT WISE	PAGE NO.
1	Force and Motion	169
2	Work and Energy	176
3	Rotational and Circular motion	182
4	Waves	189
5	Thermodynamics	196
6	Electrostatics	203
7	Current Electricity	210
8	Electromagnetism and Electromagnetic Induction	217
9	Electronics & Dawn of Modern Physics	225
10	Atomic Spectra & Nuclear Physics	231
B	Post- Prep Assessment	238

- This Pre Prep Assessment is designed to help you to pinpoint the weak areas in your background. Sit aside 1 hour to take this test. Check your answers with those at the end of the test. Then evaluate yourself.

## PRE-PREP ASSESSMENT

- Q.1 The diagram shows the energy levels for an electron in a certain atom. Which transition shown represents the emission of a photon with most energy?



- A. III  
B. IV  
C. I  
D. II
- Q.2 If the kinetic energy of a free electron doubles. Its DE Broglie wavelength changes by the factor
- A.  $\frac{1}{2}$   
B. 2  
C.  $\frac{1}{\sqrt{2}}$   
D.  $\sqrt{2}$
- Q.3 A nuclear transformation is denoted by  $X (n, \alpha) {}^7_3Li$ . Which of the following is the nucleus of element X?
- A.  ${}^{12}_6C$   
B.  ${}^{10}_5B$   
C.  ${}^9_5B$   
D.  ${}^{11}_4Be$
- Q.4 The bob of a simple pendulum is a spherical hollow ball filled with water. A plugged hole near the bottom of the oscillation bob gets suddenly unplugged. During observation, till water is coming out, the time period of oscillation would
- A. First increase and then decrease to the original value  
B. First decreased then increase to the original value  
C. Remain unchanged  
D. Increase towards a saturation value
- Q.5 An observer moves towards a stationary source of sound, with a velocity one fifth of the velocity of sound. What is the percentage increase in the apparent frequency?
- A. Zero  
B. 0.5%  
C. 5%  
D. 20%
- Q.6 The resistance of hot tungsten filament is about 10 times the cold resistance. What will be the resistance of 100 W and 200 V lamp when not in use?
- A. 40  $\Omega$   
B. 20  $\Omega$   
C. 400  $\Omega$   
D. 200  $\Omega$
- Q.7 A magnetic needle is kept in a non-uniform magnetic field. It experiences
- A. A torque but not a force  
B. Neither a force nor a torque  
C. A force and a torque  
D. A force but not a torque
- Q.8 A machine gun fires a bullet of mass 40 g with a velocity  $1200 \text{ ms}^{-1}$ . The man holding it can exert a maximum force of 144 N on the gun. How many bullets can he fire per second at the most?
- A. One  
B. Four  
C. Two  
D. Three

Q.9 A force  $\vec{F} = (5\hat{i} + 3\hat{j} + 2\hat{k})\text{N}$  is applied over a particle which displaces it from its origin to the point  $\vec{R} = (2\hat{i} - \hat{j})\text{m}$ . The work done on the particle in joules is

- A. 7  
B. +7  
C. +10  
D. +13

Q.10 A body of mass  $m$ , accelerates uniformly from rest  $v_1$  in time  $t_1$ . The instantaneous power delivered to the body as a function of time  $t$  is


- A.  $\frac{mv_1 t}{t_1}$   
B.  $\frac{mv_1^2 t}{t_1^2}$   
C.  $\frac{mv_1 t^2}{t_1}$   
D.  $\frac{mv_1^2 t}{t_1}$

Q.11 If a particle covers half the circle of radius  $R$  with constant speed, then



- A. momentum change is  $mvr$   
B. change in K.E. is  $1/2 mv^2$   
C. change in K.E. is  $mv^2$   
D. change in K.E. is zero

Q.12 The angular displacement covered by a body in the following graph is



- A. 40 rev  
B. 20 rev  
C. 30 rad  
D. 40 rad

Q.13 Restoring force in SHM is

- A. conservative  
B. non-conservative  
C. frictional  
D. centripetal

Q.14 An organ pipe, open at both ends and another organ pipe, closed at one end, will resonate with each other, if their lengths are in ratio of

- A. 1:1  
B. 1:4  
C. 2:1  
D. 1:2

Q.15 When source is moving towards observer with velocity  $u_s$ , then the modified frequency ' $f_c$ ' will be

- A.  $f_c = f \left[ \frac{v - u_s}{v} \right]$   
B.  $f_c = f \left[ \frac{v}{v + u_s} \right]$   
C.  $f_c = f \left[ \frac{v}{v - u_s} \right]$   
D.  $f_c = f \left[ \frac{v + u_s}{v} \right]$


Q.16 A rope of length 5 m is stretched to a tension of 80 N. If its mass is 1 kg, at what speed would a 10 Hz transverse wave travel down the string?

- A. 2 m/s  
B. 5 m/s  
C. 20 m/s  
D. 50 m/s

Q.17 The ratio of adiabatic bulk modulus and isothermal bulk modulus of a gas is  $(\gamma = C_p / C_v)$ .

- A. 1  
B.  $\gamma$   
C.  $\frac{\gamma}{(\gamma-1)}$   
D.  $\frac{(\gamma-1)}{\gamma}$

Q.18 In the figure curves AB and CD represent the relation between pressure  $P$  and volume  $V$  of an ideal gas. One of the curves represents an isothermal expansion and the other represents an adiabatic expansion. Which curve represents an adiabatic expansion?



- A. Curve AB  
B. Curve CD  
C. Both "a" and "b"  
D. None of these

Q.19 The force between two charges situated in air is  $F$ . The force between the same charges if the distance between them is reduced to half and they are situated in a medium having dielectric constant 4 is:

- A.  $F/4$   
B.  $16F$   
C.  $4F$   
D.  $F$

Q.20 If the force between the electron in the first Bohr orbit and the nucleus (proton) in hydrogen atom is  $F$ , then the force between them when the electron is in the second orbit is:

- A.  $4F$   
B.  $F/9$   
C.  $F/4$   
D.  $F/16$

Q.21 The given figure shows the electric lines of force emerging from a charged body. If the electric fields at A and B are  $E_A$  and  $E_B$  respectively and if the displacement between A and B is  $r$ , then:



- A.  $E_A > E_B$   
B.  $E_A = \frac{E_B}{r}$   
C.  $E_A < E_B$   
D.  $E_A = \frac{E_B}{r^2}$

Q.22 A capacitor has a capacitance of  $4\text{F}$  in presence of air. Find the capacitance in  $\mu\text{F}$  when a medium of dielectric constant 8 is placed between the plates

- A. 32  
B. 16  
C. 2  
D. None

Q.23 Two wires made of the same material and of the same length are connected in parallel to the same voltage supply. Wire P has a diameter of 2 mm. Wire Q has a diameter of 1 mm.

What is the ratio  $\frac{\text{current in P}}{\text{current in Q}}$ ?

- A.  $\frac{1}{4}$  B. 4  
C. 2 D.  $\frac{1}{2}$

Q.24 An electric iron is marked 20 volts 500W. The units consumed by it in using it for 24 hours will be \_\_\_\_\_

- A. 12 B. 24  
C. 5 D. 1100

Q.25 The masses of three wires of copper are in the ratio of 1 : 3 : 5 and their lengths are in the ratio 5 : 3 : 1. The ratio of their electrical resistance is:

- A. 1 : 3 : 5 B. 1 : 15 : 125  
C. 5 : 3 : 1 D. 125 : 15 : 1

Q.26 There are three bulbs of 60W, 100W and 200W which bulb has thickest filament.

- A. 100W B. 200W  
C. 60W D. All

Q.27 When a charged particle moving with velocity  $\vec{v}$  is subjected to a magnetic field of induction  $\vec{B}$ , the force on it is non-zero. This implies that:

- A. angle between  $\vec{v}$  and  $\vec{B}$  is either zero or  $180^\circ$   
B. angle between  $\vec{v}$  and  $\vec{B}$  can have any value other than  $90^\circ$   
C. angle between  $\vec{v}$  and  $\vec{B}$  is necessarily  $90^\circ$   
D. angle between  $\vec{v}$  and  $\vec{B}$  can have any value other than zero and  $180^\circ$

Q.28 The magnetic field in a certain region is given by  $40\hat{i} - 18\hat{k}$ . How much flux passes through a  $5.0 \text{ cm}^2$  area loop in this region if loop lies flat in YZ plane?

- A.  $90 \times 10^{-4} \text{ Wb}$  B.  $2 \times 10^{-2} \text{ Wb}$   
C.  $2 \times 10^2 \text{ Wb}$  D.  $9 \times 10^{-4} \text{ Wb}$

Q.29 A small piece of wire is passed through the gap between the poles of a magnet in 0.1 sec. An e.m.f. of  $4 \times 10^{-8} \text{ V}$  is induced in the wire, the magnetic flux between the poles is:

- A. 10 Wb B.  $4 \times 10^{-9} \text{ Wb}$   
C. 0.1 Wb D.  $4 \times 10^{-2} \text{ Wb}$

Q.30 What is the charge induced in coil of 100 turns of resistance  $100\Omega$ , if magnetic flux changes from  $2 \text{ T m}^2$  to  $-2 \text{ T m}^2$ ?

- A. 4 C B. 2.8 C  
C. 2 C D. 0.4 C

Q.31 A coil having an area  $A_0$  is placed in a magnetic field which changes from  $B_0$  to  $4B_0$  in time interval  $t$ . The e.m.f. induced in the coil will be:

- A.  $3A_0B_0/t$  B.  $3B_0/A_0t$   
C.  $4A_0B_0/t$  D.  $4B_0/A_0t$

Q.32 The coils of a stepdown transformer have 500 and 5000 turns. In the primary coil an AC of 4 A at 2200 volt is sent. The value of the current and potential difference in the secondary will be.

- A. 20 A, 22V B. 0.4 A, 22000 A  
C. 40 A, 220V D. 40 A, 22000V

Q.33 If a full wave rectifier circuit is operating from 50 Hz mains, the fundamental frequency in the ripple will be

- A. 50 Hz B. 100 Hz  
C. 70.7 Hz D. 25 Hz

Q.34 In full wave rectification, the output D.C. voltage across the load is obtained for \_\_\_\_\_

- A. The positive half cycle of input A.C. B. The negative half cycle of input A.C.  
C. The complete cycle of input A.C. D. All of the above.

Q.35 A radio station emits 10 kW power of 90.8 MHz. Find the number of photon emitted per second

- A.  $1.6 \times 10^{28}$  B.  $1.6 \times 10^{29}$   
C.  $1.6 \times 10^{30}$  D.  $1.6 \times 10^{32}$

Q.36 If  $n$  number of photon are striking on a metal surface, then total momentum exerted is \_\_\_\_\_

- A.  $nh/\lambda$  B.  $2nh\lambda$   
C. zero D.  $n \times t$

Q.37 Four lowest energy levels of H atom are shown in the figure. The number of possible emission lines would be



- A. 3 B. 4  
C. 5 D. 6

Q.38 Plutonium decays with a half-life of 24000 years. If plutonium is stored for 72000 years, the fraction of it that remains is:

- A. 1/8 B. 1/4  
C. 1/3 D. 1/2

Q.39 The radioactivity of a certain radioactive element drops to 1/64 of its initial value in 30 seconds. Its half-life is

- A. 4 seconds B. 3 seconds  
C. 5 seconds D. 2 seconds

Q.40  $\alpha$  Particle is bombarded on  ${}_{7}\text{N}^{14}$  as a result  ${}_{8}\text{O}^{17}$  is formed. The particle emitted is

- A. Neutron B. Proton  
C. Electron D. Positron


Q.41 An aeroplane flying 490 m above ground level at 100 m/s, releases a block. How far on ground will it strike

- A. 1 km B. 2 km  
C. 0.1 km D. 0.01 km

Q.42 A particle of mass having velocity ' $v$ ' makes head on elastic collision with another particle of the same mass and initially at rest. The velocity of the first particle after the collision is


- A.  $v$  B.  $-v$   
C.  $\frac{v}{2}$  D. 0

- Q.43 A particle of mass  $m$  at rest is acted upon by a force  $F$  for a time  $t$ . Its K.E. after an interval  $t$  is
- A.  $\frac{F^2 t^2}{m}$       B.  $\frac{F^2 t^2}{3m}$   
 C.  $\frac{F^2 t^2}{2m}$       D.  $\frac{Ft}{2m}$
- Q.44 The angle described in 2sec by an object rotating at a rate of 600 rpm is
- A.  $20\pi$  rad      B.  $40\pi$  rad  
 C.  $5\pi$  rad      D. Zero
- Q.45 An empty vessel is partially filled with water. Then the frequency of vibration of air column in the vessel
- A. Remain same      B. Decrease  
 C. Increase      D. First increase then decrease
- Q.46 There are two strings of equal length and diameter but the densities are in the ratio 1:2 they are stretched by a tension  $T$ . The ratio of their fundamental frequencies will be:
- A.  $\sqrt{2} : 1$       B. 1:4  
 C. 1:2      D. 2:1
- Q.47 A polyatomic gas ( $\gamma=4/3$ ) is compressed to 1/8 of its volume adiabatically. If its initial pressure is  $P_0$ , its new pressure will be:
- A.  $16P_0$       B.  $6P_0$   
 C.  $8P_0$       D.  $2P_0$
- Q.48 A gas is compressed from a volume of  $2m^3$  to a volume of  $1m^3$  at a constant pressure of  $100N/m^2$ . Then it is heated at constant volume by supplying 150 J of energy. As a result, the internal energy of the gas
- A. Decreases by 250 J      B. Increases by 50 J  
 C. Increases by 250 J      D. Decreases by 50 J
- Q.49 The variation of electric potential due to a point charge with distance is represented by the graph (where  $V$  along y-axis and  $r$  along x-axis)



- Q.50 The charge of an electron is  $1.6 \times 10^{-19}$  C. How many electrons strike the screen of a cathode ray tube each second when the beam current is 16 mA
- A.  $10^{19}$       B.  $10^{17}$   
 C.  $10^{-19}$       D.  $10^{-17}$
- Q.51 An electron enters a region where the electric field  $E$  is perpendicular to the magnetic field  $B$ . It will suffer no deflection if \_\_\_\_\_
- A.  $E = Bev$       B.  $B = cE/v$   
 C.  $E = Bv$       D.  $E = Bev/2$

- Q.52 In pushing the magnet towards the solenoid, the meter shows deflection towards



- A. Left  
 B. Right  
 C. No deflection  
 D. Vibration
- Q.53 A radioactive substance is at  $t=0$ , the number of atoms is  $8 \times 10^4$ . Its half-life period is 3 years. The number of atoms  $1 \times 10^4$  will remain after interval
- A. 19 years      B. 24 years  
 C. 9 years      D. 6 years
- Q.54 In the nuclear decay
- $${}^A_Z X \longrightarrow {}^A_{Z-1} Y \longrightarrow {}^A_{Z-2} B^* \longrightarrow {}^A_{Z-2} B$$
- The particle emitted in the sequence are
- A.  $\alpha, \beta, \gamma$       B.  $\beta, \alpha, \gamma$   
 C.  $\gamma, \beta, \alpha$       D.  $\beta, \gamma, \alpha$
- Q.55 The unit of radioactivity 'curie' is equal to \_\_\_\_\_
- A.  $3.74 \times 10^9$  disintegration per sec      B.  $3.70 \times 10^{10}$  disintegration per sec  
 C.  $3.55 \times 10^{10}$  disintegration per sec      D.  $3.60 \times 10^{10}$  disintegration per sec
- Q.56 Various types of cancer are treated by \_\_\_\_\_
- A. Cobalt-60      B. Strontium-90  
 C. Carbon-14      D. Nickel-63


# ANSWER KEY

1	A	11	D	21	A	31	A	41	A	51	C
2	C	12	D	22	A	32	C	42	D	52	B
3	B	13	A	23	B	33	B	43	C	53	C
4	A	14	C	24	A	34	C	44	B	54	B
5	D	15	C	25	D	35	B	45	C	55	B
6	A	16	C	26	B	36	A	46	A	56	A
7	C	17	B	27	D	37	D	47	A		
8	D	18	B	28	B	38	A	48	C		
9	B	19	D	29	B	39	C	49	D		
10	B	20	D	30	A	40	B	50	B		

## EXPLANATORY NOTES

$$Q.1 \quad \Delta E \propto \left( \frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

$$Q.2 \quad \lambda = \frac{h}{\sqrt{2km}}$$



Q.4 First CM goes down and then comes to its initial position.

$$Q.5 \quad t = \frac{v+v/5}{v} \quad f = \frac{6f}{5}$$

% increase in frequency = 20%

$$Q.6 \quad R_{\text{eq}} = \frac{1^2}{P} = \frac{200 \times 200}{100} = 400 \Omega$$

Cold resistance =  $R_{\text{eq}} / 10 = 400 / 10 = 40 \Omega$

Q.7 In non-uniform magnetic field, dipole experiences both force and torque.

Q.8 Change in momentum for each bullet fired is

$$= \frac{40}{1000} \times 1200 = 48 \text{ N}$$

If a bullet fired exerts a force of 48 N on man's hand so man can exert maximum force of 144 N

$$\text{Number of bullets that can be fired} = \frac{144}{48} = 3$$

Q.9 Work done  $W = \vec{F} \cdot \vec{s}$

$$\vec{s} = 2\hat{i} - \hat{j}$$

$$W = 5\hat{i} \cdot 3\hat{j} + 2\hat{k} \cdot (2\hat{i} - \hat{j}) = 10 - 3 = 7 \text{ J}$$

$$Q.10 \quad P = \vec{F} \cdot \vec{v} = mav = m \begin{pmatrix} v_x \\ v_y \\ v_z \end{pmatrix} \cdot \begin{pmatrix} v_x \\ v_y \\ v_z \end{pmatrix} = \frac{mv^2}{t}$$

Q.11 As momentum is vector quantity so change in momentum  $\Delta P = mv - (-mv) = 2mv$ . But kinetic energy remains always constant so change in kinetic energy is zero.

$$Q.12 \quad \theta = \omega \times t = 10 \times 4 = 40 \text{ rad}$$

Q.13 Restoring force is conservative force.

$$Q.14 \quad t_1 = t_2$$

$$\frac{v}{2t_1} = \frac{v}{4t_2}$$

$$t_1 : t_2 = 2 : 1$$

$$Q.15 \quad f_1 = \left( \frac{v}{v-u} \right) f$$

$$Q.16 \quad v = \sqrt{\frac{F}{m/l}} \Rightarrow v = \sqrt{\frac{80 \times 5}{1}} = 20 \text{ ms}^{-1}$$

Q.17  $E = P$  For isothermal process

$E = \gamma P$  For adiabatic process

$$\text{So, } \frac{\gamma P}{P} = \gamma$$

Q.18 CD curve is more steeper, it represents rapid process.

$$Q.19 \quad F_{\text{net}} = \frac{F_{\text{cm}}}{e_1}$$

$$Q.20 \quad F = \frac{kq^2}{r^2}$$

For second orbit

$$r_2 = 4r_1$$

$$F' = \frac{F}{16}$$

$$Q.21 \quad E = \frac{kq}{r}$$

$$r_A < r_B$$

$$E_A > E_B$$

$$Q.22 \quad \frac{1}{e_1} = \frac{C}{C_m} \Rightarrow C_m = e_1 C_1 = 4 \times 8 = 32$$

$$Q.23 \quad R = \frac{\rho L}{A} = \frac{\rho L}{\pi \left( \frac{D}{2} \right)^2} \Rightarrow R = \frac{1}{D^2} \Rightarrow \frac{R_x}{R_y} = \frac{D_y^2}{D_x^2}$$

$$V = IR \Rightarrow \frac{I_x}{I_y} = \frac{R_y}{R_x} = \frac{D_x^2}{D_y^2} = \frac{(2)^2}{(1)^2} = 4$$

$$Q.24 \quad \text{Number of units} = \frac{P(W) \times t(h)}{1000} = \frac{500 \times 24}{1000} = 12$$

$$Q.25 \quad R = \rho \frac{L}{A}$$

$$R = \frac{\rho L^2}{V} \left( \because A = \frac{V}{L} \right)$$

$$R = \frac{\rho d L^2}{m} \left( \because V = \frac{m}{d} \right)$$

$$R = \frac{L^2}{m}$$

$$R_1 : R_2 : R_3 = \frac{5^2}{1} : \frac{3^2}{3} : \frac{1^2}{5} = 125 : 15 : 1$$

Q.26 For parallel combination

$$P = \frac{V^2}{R} \Rightarrow P \propto \frac{1}{R} \quad (V = \text{constant})$$

Q.27  $F = qvB \sin \theta$  other than  $0^\circ$  and  $180^\circ$  it gives non zero values.

Q.28  $B = 40\hat{i} + 0\hat{j} - 18\hat{k}$

$$A = (5\hat{i} + 0\hat{j} + 0\hat{k}) \times 10^{-4} \text{ m}^2$$

$$\begin{aligned} \phi_b &= \vec{B} \cdot \vec{A} \\ &= 40 \times 5 \times 10^{-4} \\ &= 2 \times 10^{-3} \text{ Wb} \end{aligned}$$

Q.29  $\epsilon = \frac{N\Delta\phi}{\Delta t}$   
 $\Delta\phi = \epsilon \times \Delta t \quad (\because N = 1)$

Q.30  $\epsilon = \frac{N\Delta\phi}{\Delta t}$   
 $IR = \frac{N\Delta\phi}{\Delta t}$   
 $\frac{Q}{\Delta t} R = \frac{N\Delta\phi}{\Delta t} \Rightarrow Q = \frac{N\Delta\phi}{R}$

Q.31  $\epsilon = \frac{N\Delta\phi}{\Delta t}$   
 $\epsilon = \frac{3A_s B_s}{\Delta t}$

Q.32  $\frac{E_s}{E_p} = \frac{N_s}{N_p} \Rightarrow E_s = \frac{N_s}{N_p} E_p = \frac{500}{5000} \times 2200 = 220 \text{ V}$

As  $E_s i_s = E_p i_p \Rightarrow i_s = \frac{E_p}{E_s} i_p$

Or  $i_s = \frac{N_p}{N_s} i_p = \frac{5000}{500} \times 4 = 40 \text{ A}$

Q.33  $f' = 2f_{(\text{fundamental})}$

Q.34 In full wave rectification, complete cycle of A.C is rectified.

Q.35  $P = \frac{E}{t} = \frac{nhf}{t} \Rightarrow \frac{n}{t} = \frac{P}{hf}$

$$\frac{n}{t} = \frac{10 \times 10^1}{(6.63 \times 10^{-34})(90.8 \times 10^6)}$$

$$n = 1.6 \times 10^{29} \text{ photons per second}$$

Q.36 As  $p = \frac{h}{\lambda}$  for 1 photon

$$\therefore p = \frac{nh}{\lambda} \text{ for } n \text{ photon}$$

Q.37 No. of possible spectral lines =  $\frac{n(n-1)}{2} = \frac{4(4-1)}{2} = 2(3) = 6$

Q.38 Undecayed atoms =  $\frac{1}{2^n}$   
 $\therefore n = \text{number of half lives}$   
 Undecayed atoms =  $\frac{1}{2^3} = \frac{1}{8}$

Q.39  $N = N_0 \left(\frac{1}{2}\right)^{\frac{t}{T}} \Rightarrow \frac{N_0}{64} = N_0 \left(\frac{1}{2}\right)^{\frac{t}{T}} \Rightarrow T = \frac{30}{6} = 5 \text{ sec}$

Q.40  ${}^7\text{N}^{14} + {}^2\alpha^4 \rightarrow {}^8\text{O}^{17} + {}^1\text{H}^1 \Rightarrow$  Proton will be emitted.

Q.41  $S = u \times \sqrt{\frac{2h}{g}} = 100 \times \sqrt{\frac{2 \times 490}{9.8}} = 1000 \text{ m} = 1 \text{ km}$

Q.42 If masses are same their velocities will alter, so velocity of 1<sup>st</sup> ball after collision = velocity of 2<sup>nd</sup> ball before collision = 0

Q.43 Kinetic energy  $E = \frac{P^2}{2m} = \frac{(Ft)^2}{2m} = \frac{F^2 t^2}{2m}$  [As  $P = Ft$ ]

Q.44  $\theta = \alpha t = \frac{600 \times 2\pi}{60} \times 2 = 40\pi \text{ rad}$

Q.45  $f_n = n \left(\frac{v}{4l}\right) \Rightarrow \left(f \propto \frac{1}{l}\right)$ , length of column decrease so frequency will be increase.

Q.46  $f = \frac{1}{2l} \sqrt{\frac{T}{m}} = \frac{1}{2l} \sqrt{\frac{T}{\rho V}} \Rightarrow f \propto \frac{1}{\sqrt{\rho}} \quad \because \rho = \frac{m}{V}$

$$\frac{f_1}{f_2} = \sqrt{\frac{\rho_2}{\rho_1}} = \sqrt{\frac{2}{1}}$$

Q.47  $\frac{P_2}{P_1} = \left(\frac{V_1}{V_2}\right)^4 \Rightarrow P_2 = P_1 \left(\frac{V_1}{V_2}\right)^4 = P_1 (8)^4 = 16P_1$

Q.48 As we know,  $\Delta Q = \Delta U + \Delta W$  (1<sup>st</sup> law of thermodynamics)

$$\Rightarrow \Delta Q = \Delta U + P\Delta V$$

$$150 = \Delta U + 100(1-2) = \Delta U - 100 \therefore \Delta U = 150 + 100 = 250 \text{ J}$$

Q.49  $V = \frac{kq}{r} \Rightarrow V \propto \frac{1}{r}$



1  
TOPIC

## FORCE AND MOTION

## PRACTICE EXERCISE

$$Q.50 \quad i = \frac{ne}{t} \Rightarrow 16 \times 10^{-3} = \frac{n \times 1.6 \times 10^{-19}}{1} \Rightarrow n = 10^{17}$$

$$Q.51 \quad qvB = Eq$$

$$E = Bv$$

Q.52 When magnet is pushed toward solenoid, change in flux with respect to time will increase and induced current will also increase, so it shows deflection of meter towards right.

Q.53 By formula

$$N = N_0 \left(\frac{1}{2}\right)^{t/T} \text{ or } 10^4 = 8 \times 10^4 \left(\frac{1}{2}\right)^{t/3} \text{ or } \left(\frac{1}{2}\right) = \left(\frac{1}{2}\right)^{t/3} \text{ or } \left(\frac{1}{2}\right)^3 = \left(\frac{1}{2}\right)^{t/3} \Rightarrow 3 = \frac{t}{3}$$

Hence  $t = 9$  years

Q.54 Charge number increases by 1, so it is  $\beta$ -emission

Charge number decreases by 2, so it is  $\alpha$ -emission

No change in charge number so it is  $\gamma$ -emission


Q.55 Curie =  $3.7 \times 10^{10}$  disintegrating/sec

Q.56 Cobalt 60 is used for treatment of cancer.

- Q.1 A cyclist moves along a circular path of radius 70m. If he completes one round in 11s, calculate the total length of a path.
- A. 40m  
B. 440m  
C. 0m  
D. 11m
- Q.2 A body is projected horizontally from the top of a cliff with a velocity of 9.8m/s. What time elapses before horizontal and vertical velocities become equal? Take  $g = 9.8\text{m/s}^2$
- A. 9.8s  
B. 0s  
C. 10s  
D. 1s
- Q.3 Two bullets are fired simultaneously, horizontally and with different speeds from the same place. Which bullet will hit the ground first?
- A. The faster one  
B. Depends on their mass  
C. The slower one  
D. Both will reach simultaneously
- Q.4 A body is moving along a straight path. What will happen to the body in the absence of an external field?
- A. It will stop  
B. It will move with the same speed in a different path  
C. It will move with the same speed along the same straight path  
D. It will move with a reduced speed along the same path
- Q.5 A person is standing in a bus. When the bus starts moving forward suddenly
- A. The person moves forward  
B. The person remains stationary  
C. The person is unaffected  
D. The person moves backward
- Q.6 A ship of mass  $3 \times 10^7\text{kg}$  initially at rest is pulled by a force of  $5 \times 10^4\text{N}$  through a distance of 3m. Assuming that the resistance due to water is negligible, the speed of the ship is?
- A. 1.5m/s  
B. 60m/s  
C. 0.1m/s  
D. 5m/s
- Q.7 Which of the following is an example of inelastic collision?
- A. Collision between two vehicles  
B. Collision between glass balls  
C. A bullet fired into a wooden block  
D. Collision between two railway compartments
- Q.8 Mud thrown on a wall and sticking to it is an example for \_\_\_\_\_
- A. Inelastic collision  
B. Elastic collision  
C. Super elastic collision  
D. Perfectly inelastic collision
- Q.9 A Body moves 6 m north. 8 m east and 10m vertically upwards, what is its resultant displacement from initial position
- A.  $10\sqrt{2}m$   
B.  $\frac{10}{\sqrt{2}}m$   
C. 10 m  
D.  $10 \times 2m$

Topic-1

Q.10 A particle starts from center O towards A then moves along AB and stop at B. if  $R=100\text{m}$  then displacement of the particle is



- A. 100 m
- B.  $100\sqrt{2}$  m
- C.  $\frac{100}{\sqrt{2}}$  m
- D. None

Q.11 The instantaneous acceleration is the limit of average acceleration as  $\Delta t \rightarrow 0$  is given by

- A.  $\bar{a}_{ins} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \bar{d}}{\Delta t}$
- B.  $\bar{a}_{ms} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \bar{v}}{\Delta t}$
- C.  $\bar{a}_{ins} = \lim_{\Delta t \rightarrow 0} \frac{\Delta v}{\Delta t}$
- D.  $\bar{a}_{ms} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \bar{a}}{\Delta t}$

Q.12 If an object is moving with constant velocity of  $20\text{ms}^{-1}$  towards north then its acceleration will be

- A.  $5\text{ m s}^{-2}$
- B.  $10\text{ m s}^{-2}$
- C.  $9\text{ m s}^{-2}$
- D.  $0\text{ m s}^{-2}$

Q.13 The retardation is defined as

- A. Increase in velocity per unit time
- B. Decrease in velocity per unit time
- C. Decrease in speed per unit time
- D. Increase in speed per unit time

Q.14 Consider the acceleration, velocity and displacement of a tennis ball as it falls to the ground and bounces back. Directions of which of these changes in the process

- A. Velocity only
- B. Acceleration, velocity and displacement
- C. Displacement and velocity
- D. Displacement and acceleration


Q.15 A particle goes from  $x = -2\text{m}$ ,  $y = 3\text{m}$ ,  $z = 1\text{m}$  to  $x = 3\text{m}$ ,  $y = -1\text{m}$ ,  $z = 4\text{m}$ . Its displacement is:

- A.  $(1\text{m})\hat{i} + (2\text{m})\hat{j} + (5\text{m})\hat{k}$
- B.  $(5\text{m})\hat{i} - (4\text{m})\hat{j} + (3\text{m})\hat{k}$
- C.  $-(5\text{m})\hat{i} + (4\text{m})\hat{j} - (3\text{m})\hat{k}$
- D.  $-(1\text{m})\hat{i} - (2\text{m})\hat{j} - (5\text{m})\hat{k}$

Q.16 Length of the path of a particle is equal to the magnitude of the displacement of that particle. Shape of the path possible

- A. Circle
- B. Arc of a circle
- C. Parabola
- D. Straight line

Q.17 The displacement versus time graph for a body moving in a straight line is shown in figure. Which of the following regions represents the motion when no force is acting on the body?



- A. ab
- B. cd
- C. bc
- D. de

Topic-1

Q.18 The shortest distance between two points is called

- A. Acceleration
- B. Velocity
- C. Speed
- D. Displacement

Q.19 Two cars are moving in opposite directions with speed  $v$ . What is the magnitude of their relative velocity?

- A. 0
- B.  $v$
- C.  $v/2$
- D.  $2v$

Q.20 A man in a car moving with velocity of  $36\text{ km/hr}$ . His speed with respect to the car is

- A.  $10\text{ m/s}$
- B. Zero
- C.  $36\text{ m/s}$
- D. Infinite


Q.21 A body can have constant velocity when it follows a.

- A. Elliptical path
- B. Parabolic path
- C. Circular path
- D. Rectilinear path

Q.22 A man leaves his house for a cycle ride. He comes back to his house after half an hour after covering a distance of one km. What is his average velocity for the ride?


- A.  $2\text{ km per hour}$
- B. 0
- C.  $1/2\text{ km per hour}$
- D.  $1/2\text{ km per second}$

Q.23 A particle moves along the sides AB, BC and CD of a square of side  $25\text{m}$  with a velocity of  $15\text{ms}^{-1}$ . Its average velocity is




- A.  $15\text{ms}^{-1}$
- B.  $7.5\text{ms}^{-1}$
- C.  $10\text{ms}^{-1}$
- D.  $5\text{ms}^{-1}$

Q.24 The Figure shows the velocity time graph of a one-dimensional motion. Which of the following characteristic of the particle is represented by the shaded area?




- A. Distance covered
- B. Speed
- C. Momentum
- D. Acceleration

Q.25 Which graph represents the motion of a car that is travelling along a straight road with a uniformly increasing speed?




Topic-1

Q.26 A particle is moving in a straight line with uniform acceleration. Which graph represents the motion of the particle?




Q.27 In the following graph, distance travelled by the body in metres is




- A. 200
- C. 250

- B. 300
- D. 400

Q.28 Displacement time graph of a ball thrown vertically upward is shown in figure



Which of the following represents v-t graph?



Q.29 If the slope of a velocity time graph gradually decreases, then the body is said to be moving with

- A. Positive acceleration
- B. Negative acceleration
- C. Uniform velocity
- D. None of these

Q.30 If the force acting on a body is doubled, then acceleration becomes

- A. Half
- B. Doubled
- C. One fourth
- D. Constant

Q.31 When force of 1 N is applied on a body of mass 100 g then the acceleration would be

- A.  $5 \text{ m s}^{-2}$
- B.  $0.5 \text{ m s}^{-2}$
- C.  $10 \text{ m s}^{-2}$
- D.  $0.1 \text{ m s}^{-2}$

Q.32 A mass of 10 kg moves with an acceleration of  $10 \text{ m s}^{-2}$ , the force on it is

- A. 5 N
- B. 100 N
- C. 50 N
- D. 25 N

Q.33 Which law of motion defines force?


- A. 1<sup>st</sup> law
- B. 2<sup>nd</sup> law
- C. 3<sup>rd</sup> law
- D. All of these

Topic-1

Q.34 A Force of 12 N gives an object an acceleration of  $4 \text{ m s}^{-2}$ . The force required to give it an acceleration of  $10 \text{ m s}^{-2}$  is

- A. 15 N
- B. 25 N
- C. 20 N
- D. 30 N

Q.35 A balloon is acted upon by three forces, weight, upthrust and sideways force due to the wind, as shown in the diagram




What is the vertical component of the resultant force on the balloon?

- A. 500 N
- B. 1000 N
- C. 10000 N
- D. 10500 N

Q.36 When a force of 4 N acts on a mass of 2 kg for a time of 2 sec, what is the rate of change of momentum?

- A.  $2 \text{ kg m s}^{-2}$
- B.  $4 \text{ kg m s}^{-2}$
- C.  $8 \text{ kg m s}^{-2}$
- D.  $16 \text{ kg m s}^{-2}$

Q.37 Two equal masses travel towards each other on a frictionless air track at speeds of  $60 \text{ cm s}^{-1}$  and  $30 \text{ cm s}^{-1}$ . They stick together on impact.



What is the speed of the masses after impact?

- A.  $15 \text{ cm s}^{-1}$
- B.  $30 \text{ cm s}^{-1}$
- C.  $20 \text{ cm s}^{-1}$
- D.  $45 \text{ cm s}^{-1}$

Q.38 Which is a statement of the principle of conservation of momentum?

- A. A force is equal to the rate of change of momentum of the body upon which it acts.
- B. The momentum of a body is the product of the mass of the body and its velocity
- C. In a perfectly elastic collision, the relative momentum of the bodies before impact is equal to their relative momentum after impact
- D. The total momentum of a system of interacting bodies remains constants, providing no external force acts

Q.39 The average force necessary to stop a hammer with 25 N s momentum in 0.05 s expressed in 'N' is

- A. 500
- B. 125
- C. 50
- D. 25


Q.40 In which of the following cases forces may not be required to keep the

- A. Particle going in a circle
- B. The momentum of the particle constant
- C. Particle going along a straight-line
- D. Acceleration of the particle constant

Q.41 A force of 6 N acts on a mass of 1kg which acquire velocity of  $30 \text{ m s}^{-1}$ . The time for which the force acts is

- A. 26 s
- B. 6 s
- C. 5 s
- D. 2 s

- Q.42 A force of 100 Dynes acts on mass of 5g for 10 sec. The velocity produced is  
 A. 2 cm/sec B. 200 cm/sec  
 C. 20 cm/sec D. 2000 cm/sec
- Q.43 The graph shows how the force acting on a body varies with time. Assuming that the body is moving in a straight line, by how much does its momentum change?



- A.  $40 \text{ kg ms}^{-1}$  B.  $36 \text{ kg ms}^{-1}$   
 C.  $20 \text{ kg ms}^{-1}$  D.  $16 \text{ kg ms}^{-1}$
- Q.44 A gun after firing recoils due to  
 A. Conservation of energy B. Conservation of momentum  
 C. Backward thrust of gases produced D. Newton's first law of motion
- Q.45 A player takes 0.1 s in catching a ball of mass 150 g moving with velocity of 20 m/s. The force imparted by the ball on the hands of the player is:  
 A. 0.3 N B. 3 N  
 C. 300 N D. 30 N
- Q.46 A ball of mass 2 kg travelling at  $8 \text{ ms}^{-1}$  strikes a ball of mass 4 kg travelling at  $2 \text{ ms}^{-1}$ . Both balls are moving along the same straight line as shown



After collision, both balls move at the same velocity  $v$ . What is the magnitude of the velocity  $v$ ?

- A.  $4 \text{ ms}^{-1}$  B.  $6 \text{ ms}^{-1}$   
 C.  $5 \text{ ms}^{-1}$  D.  $8 \text{ ms}^{-1}$
- Q.47 Two similar spheres, each of mass  $m$  and travelling with speed  $v$ , are moving towards each other.



The spheres have a head on elastic collision. Which statement is correct?

- A. The spheres stick together on impact B. The total kinetic energy before impact is zero  
 C. The total kinetic energy after impact is  $mv^2$  D. The total momentum before impact is  $2mv$
- Q.48 Two railway trucks of masses  $m$  and  $3m$  move towards each other in opposite directions with speeds  $2v$  and  $v$  respectively. These trucks collide and stick together. What is the speed of the trucks after the collision?
- A.  $\frac{v}{4}$  B.  $v$   
 C.  $\frac{v}{2}$  D.  $\frac{5v}{4}$
- Q.49 The collision between the two bodies is elastic if bodies are?  
 A. Solid and soft B. Solid and hard  
 C. Hard and elastic D. Soft and elastic

- Q.50 When a very heavy ball 'B<sub>1</sub>' collide with a stationary target 'B<sub>2</sub>' of negligible mass, after collision the final velocity of ball 'B<sub>2</sub>' will  
 A. Become zero B. Become half  
 C. Become doubled as compared to B<sub>1</sub> D. Same as the B<sub>1</sub>
- Q.51 A shell explodes into four unequal parts. Which one of the following is conserved?  
 A. Potential energy B. Kinetic energy  
 C. Momentum D. Both potential and kinetic energy
- Q.52 A handball is tossed vertically upward with a velocity of 19.6 meters per second. Approximately how high will it rise?  
 A. 15 m B. 20 m  
 C. 25 m D. 30 m
- Q.53 Which shows the correct relation between time-of-flight  $T$  and maximum height  $H$ ?  
 A.  $H = \frac{gT^2}{8}$  B.  $H = \frac{8g}{T^2}$   
 C.  $H = \frac{8T^2}{g}$  D.  $H = \frac{8}{gT^2}$
- Q.54 At maximum height on the trajectory which of projectile becomes zero  
 A. Acceleration B. Vertical velocity  
 C. Velocity D. Horizontal velocity
- Q.55 Time taken by a projectile to reach maximum height is  $t =$   
 A.  $\frac{v_i \sin \theta}{2g}$  B.  $\frac{v_i \sin \theta}{g}$   
 C.  $\frac{v_i \sin 2\theta}{g}$  D.  $\frac{2v_i \sin \theta}{g}$
- Q.56 Two projectiles are projected at angle of  $20^\circ$  and  $70^\circ$  with same velocity which one have longer range  
 A. Which is fired at  $20^\circ$  B. Both have same range  
 C. Which is fired at  $70^\circ$  D. None of these
- Q.57 The path followed by a projectile is known as its  
 A. Range B. Trajectory  
 C. Cycle D. Height
- Q.58 A projectile is launched at point O and follows the path OPQRS, as shown. Air resistance may be neglected.



Which statement is true for the projectile when it is at the highest point Q of its path?


- A. The horizontal component of the projectile's acceleration is zero  
 B. The kinetic energy of the projectile is zero  
 C. The horizontal component of the projectile's velocity is zero  
 D. The momentum of the projectile is zero

- Q.59 A projectile is fired horizontally with an initial speed of 20 m/s. Its horizontal speed 3s later is  
 A. 20 m/s  
 B. 6.67 m/s  
 C. 60 m/s  
 D. 29.4 m/s
- Q.60 For which of the following angles range is maximum?  
 A. 45°  
 B. 30°  
 C. 60°  
 D. None
- Q.61 A bullet is fired horizontally from a rifle at a distant target. Ignoring the effect of air resistance, what is the horizontal and vertical acceleration of the bullet?  
 Horizontal  
 A. 9.8 ms<sup>-2</sup>  
 B. 0  
 C. 9.8 ms<sup>-2</sup>  
 D. 0  
 Vertical  
 A. 9.8 ms<sup>-2</sup>  
 B. 9.8 ms<sup>-2</sup>  
 C. 0 ms<sup>-2</sup>  
 D. 0

**PAST PAPER MCQs**

- Q.62 A ball is just allowed to fall from the window of a moving train, it will hit the ground following.  
 A. Circular path  
 B. Hyperbolic  
 C. Straight line path  
 D. Parabolic path
- Q.63 A particle mover along the straight line. The distance x describes in time, t is given by the equation  $x = t^3$  the acceleration at t=1  
 A. A = 1  
 B. A = 2  
 C. A = 3  
 D. A = 4
- Q.64 The vertical velocity of ball thrown upward \_\_\_\_\_ with time.  
 A. Decreases linearly  
 B. Doubles  
 C. Remains constant  
 D. Decreases parabolically
- Q.65 An \_\_\_\_\_ missile is called a ballistic missile.  
 A. Un-powered and guided  
 B. Powered and guided  
 C. Un-guided and powered  
 D. Un-powered and un-guided
- Q.66 A body is moving with an initial velocity of 2 kms<sup>-1</sup>. After a time of 50 secs its velocity becomes 1.5 kms<sup>-1</sup>. Its acceleration will be:  
 A. 30 m s<sup>-2</sup>  
 B. 20 m s<sup>-2</sup>  
 C. 40 m s<sup>-2</sup>  
 D. 10 m s<sup>-2</sup>
- Q.67 In elastic collision, when a massive body collides with light body at conditions  $m_1 \gg m_2$  and  $v_2 = 0$  ms<sup>-1</sup>, then the change in velocity will be written as:  
 A.  $v_1' = -v_1$ ;  $v_2' = v_1$   
 B.  $v_1' = v_1$ ;  $v_2' = 2v_1$   
 C.  $v_1' = v_1$ ;  $v_2' = 0$   
 D.  $v_1' = -v_1$ ;  $v_2' = 0$
- Q.68 If a force of 12N is applied on a body and its momentum is changed form 60 kgms<sup>-1</sup> to 36 kg ms<sup>-1</sup>, then find the time during, which this force acts:  
 A. 1 second  
 B. 12 seconds  
 C. 2 seconds  
 D. 24 seconds
- Q.69 Time of projectile's flight is  
 A.  $\frac{v_i^2 \sin^2 \theta}{g}$   
 B.  $\frac{v_i^2 \sin \theta}{g}$   
 C.  $\frac{2v_i \sin \theta}{g}$   
 D.  $v_i^2 \sin^2 \theta$

- Q.70 If the velocity of the body changes by equal amount in equal intervals of time, the body is said to have:  
 A. Variable acceleration  
 B. Uniform velocity  
 C. Uniform acceleration  
 D. Negative acceleration
- Q.71 For finding the height of projectile, the equation used is:  
 A.  $2S = a(v_f^2 - v_i^2)$   
 B.  $S = 2a(v_f^2 - v_i^2)$   
 C.  $2aS = v_f^2 - v_i^2$   
 D.  $a = 2S(v_f^2 - v_i^2)$
- Q.72 One ball is thrown vertically upward with a velocity of 9.8 m/s. If it takes 10 seconds to reach the highest point, then the acceleration of the ball is  
 A. 9.8ms<sup>-2</sup>  
 B. 980 ms<sup>-2</sup>  
 C. 98ms<sup>-2</sup>  
 D. -9.8ms<sup>-2</sup>
- Q.73 A ball is dropped from the roof of a very tall building. What is its velocity after falling for 5.0s?  
 A. 1.96 ms<sup>-1</sup>  
 B. 9.80 ms<sup>-1</sup>  
 C. 49.0 ms<sup>-1</sup>  
 D. 98.0 ms<sup>-1</sup>
- Q.74 A projectile is launched at 45° to the horizontal with initial K, Energy, E. Assuming air resistance to be negligible, what will be the kinetic energy of the projectile when it reaches its highest point?  
 A. 0.71E  
 B. 0.50E  
 C. 0.87E  
 D. E
- Q.75 Weight rather than mass be used in calculating  
 A. Moment of inertia of a body  
 B. The stress in wire due to load hanging from it  
 C. The binding energy of the nucleus  
 D. The gravitational force between the two bodies
- Q.76 The ratio of displacement along diameter and total distance along circle:  
 A. 1 :  $\pi$   
 B. 2 :  $\pi$   
 C.  $\pi$  : 1  
 D.  $\pi$  : 2
- Q.77 Arshad is driving down 7<sup>th</sup> street. He drives 150 meter in 18 seconds. Assume he does not speed up or slow down. What is his speed?  
 A. 0.38m/s  
 B. 8.33m/s  
 C. 126m/s  
 D. 58.33m/s
- Q.78 The distance travelled by a moving car with velocity 15 m/s in 2 seconds, decelerates at 2m/s<sup>2</sup> is equal to  
 A. 30m  
 B. 16m  
 C. 34m  
 D. 26m
- Q.79 The value of ratio of displacement to distance is:  
 A. Always one  
 B. More than one  
 C. Always less than one  
 D. Equal or less than one
- Q.80 Which of the following v-t graph represents the constant acceleration: (MDCAT 2017)



- A. B. C. D. All of these

(ETEA 2017)

- Q.81 The numerical ratio of displacement to distance is:  
 A. Always less than one  
 B. Always equal to one  
 C. Always more than one  
 D. Equal to or less than one
- Q.82 A bullet of mass  $m$  moving with a velocity  $v$  is fired into large wooden block of mass  $M$  of the bullet remains embedded in the wooden block, the velocity of the system will be: (ETEA 2017)
- A.  $\frac{M}{M+m}v$   
 B.  $\frac{m}{M+m}v$   
 C.  $\frac{M}{M-m}v$   
 D.  $\frac{m}{M-m}v$
- Q.83 If slope of velocity time graph is not constant at different points, then body is moving with  
 A. Uniform velocity  
 B. Average acceleration  
 C. Increasing acceleration  
 D. Constant acceleration
- Q.84 A cyclist is traveling at  $15\text{ms}^{-1}$  she applies brakes so that she doesn't collide with the wall in front of her distance of  $18\text{m}$ . Calculate the magnitude of deceleration. (MDCAT 2018)
- A.  $6.3\text{ms}^{-2}$   
 B.  $13\text{ms}^{-2}$   
 C.  $5.3\text{ms}^{-2}$   
 D.  $12.5\text{ms}^{-2}$
- Q.85 Newton first law of motion is also known is  
 A. Law of inertia  
 B. Law of universal gravity  
 C. Law of electromagnetism  
 D. Law of conservation
- Q.86 If two objects of equal masses ' $m$ ' are moving towards each other with the same speed ' $v$ ' then what will be the total final momentum after elastic head-on collision? (MDCAT 2019)
- A.  $-mv \text{ kg/s}$   
 B.  $2mv \text{ kg/s}$   
 C.  $mv \text{ kg m/s}$   
 D.  $0 \text{ kg m/s}$
- Q.87 For projectile motion in the absence of air resistance:  
 A. vertical speed is constant  
 B. horizontal acceleration is zero  
 C. horizontal force is constant  
 D. vertical acceleration is zero
- Q.88 The range of the projectile depends upon the velocity of the projection and the angle of the projection i.e  $45^\circ$ . For a fixed velocity, when the angle of projection is larger than  $45^\circ$ . Which of the following is correct? (MDCAT 2019)
- A. both the height and the range attained by the projectile will be less  
 B. The height attained by the projectile will be less but the range is more  
 C. both the height and the range attained by the projectile will be more  
 D. The height attained by the projectile will be more but the range is less
- Q.89 The slope of distance - time graph will always be: (NMDCAT 2021)
- A. Negative  
 B. Positive  
 C. Zero  
 D. Maximum
- Q.90 At what angle of projection of a projectile the range becomes half of its maximum value? (NMDCAT 2021)
- A.  $15^\circ$   
 B.  $20^\circ$   
 C.  $30^\circ$   
 D.  $40^\circ$

- Q.91 If we drop an object, its initial velocity is zero. How far will it fall in time ' $t$ '? (NMDCAT 2020)
- A.  $9.8t^2$   
 B.  $4.9t^2$   
 C.  $0.49t^2$   
 D.  $98t^2$
- Q.92 The newton - second is unit of: (NMDCAT 2020)
- A. Work  
 B. Power  
 C. Impulse  
 D. Momentum
- Q.93 Which one of the following is the angle of projection of a projectile if its range is equal to its height? (NUMS 2020)
- A.  $48^\circ$   
 B.  $60^\circ$   
 C.  $90^\circ$   
 D.  $76^\circ$
- Q.94 The product of force and time is equal to: (NUMS 2020)
- A. Angular momentum  
 B. Force  
 C. Change in momentum  
 D. Velocity
- Q.95 At what angle of projection of a projectile the range becomes half of its maximum value. (NMDCAT 2021)
- A.  $15^\circ$   
 B.  $20^\circ$   
 C.  $20^\circ$   
 D.  $40^\circ$
- Q.96 Vertical velocity vs time graph for a projectile motion (NMDCAT 2021)
- A. Varies linearly  
 B. Follow a parabolic path  
 C. Is constant  
 D. Is linear
- Q.97 The magnitude of displacement is (NMDCAT 2021)
- A. Size of object A  
 B. Straight line distance between initial and final position  
 C. Size of object B  
 D. Any distance between initial and final position
- Q.98 If the displacement =  $15\text{m}$  and time =  $10\text{seconds}$  the average velocity is (NMDCAT 2021)
- A.  $12.5\text{m/s}$   
 B.  $1.5 \text{ m/s}$   
 C.  $2.5\text{m/s}$   
 D.  $3\text{m/s}$
- Q.99 Decrease in velocity per unit time is called (NMDCAT 2021)
- A. Acceleration  
 B. Deceleration  
 C. Positive acceleration  
 D. Uniform acceleration
- Q.100 The slope of velocity-time graph represents (NMDCAT 2021)
- A. Displacement  
 B. Acceleration  
 C. Force  
 D. Momentum
- Q.101 An object is accelerating (NMDCAT 2021)
- A. Only when its speed changes  
 B. Only when its direction changes  
 C. When its speed or direction changes  
 D. If its velocity is large
- Q.102 What is the slope of a straight-line graph of position vs. time? (NMDCAT 2021)
- A. Velocity  
 B. Displacement  
 C. Distance  
 D. Acceleration
- Q.103 The coefficient of restitution for a perfectly elastic collision is (NMDCAT 2021)
- A. 1  
 B. 0  
 C.  $\infty$   
 D. -1

- Q.104 What is the accelerating force on the Diwali rocket if it ejects 0.05 kg of gases per second at a velocity of 400 m/s?  
 A. 20 N  
 B. 20 dynes  
 C. 1000 N  
 D. 22 dynes  
 (NMDCAT 2021)
- Q.105 If mass is 2 kg and velocity is 35 m/s, then momentum is  
 A. 70 kg m/s  
 B. 90 kg m/s  
 C. 17.5 kg m/s  
 D. 105 kg m/s  
 (NMDCAT 2021)
- Q.106 If Velocity time graph shape is parallel to Y-axis the acceleration  
 A. Maximum  
 B. 0  
 C. Infinity  
 D. Minimum  
 (NMDCAT 2021)

# ANSWER KEY

## TOPIC-WISE MCQs & PAST PAPER MCQs

1	B	11	C	21	D	31	C	41	C	51	C	61	B	71	C	81	D	91	B	101	C
2	D	12	D	22	B	32	B	42	B	52	B	62	D	72	D	82	B	92	C	102	B
3	D	13	B	23	D	33	A	43	C	53	A	63	A	73	C	83	C	93	D	103	A
4	C	14	C	24	A	34	D	44	B	54	B	64	A	74	B	84	A	94	C	104	A
5	D	15	B	25	A	35	B	45	D	55	B	65	D	75	D	85	A	95	A	105	A
6	C	16	D	26	B	36	B	46	A	56	B	66	D	76	A	86	D	96	A	106	C
7	A	17	B	27	A	37	A	47	C	57	B	67	B	77	B	87	B	97	B		
8	d	18	D	28	C	38	D	48	A	58	A	68	C	78	D	88	D	98	B		
9	A	19	D	29	A	39	A	49	B	59	A	69	C	79	D	89	B	99	B		
10	A	20	B	30	B	40	B	50	C	60	A	70	C	80	A	90	A	100	B		

# EXPLANATORY NOTES

- Radius of the circular path,  $r = 70\text{m}$   
 Time takes to complete one round,  $t = 11\text{s}$   
 Total length of the path,  $s = 2\pi r = 2 \times 22/7 \times 70 = 440\text{m}$ .
- Horizontal velocity at any instant,  $v_x = u = 9.8\text{m/s}$   
 Vertical velocity at any instant,  $v_y = 0 + gt = 9.8t$   
 $9.8 = 9.8t$   
 $t = 1\text{s}$ .
- Both the bullets will reach simultaneously because  
 $T = \sqrt{2h/g}$  Time  $t$  does not depend on horizontal speed.
- Galileo stated that on a horizontal plane an object should move with a constant velocity in a straight-line path. Therefore, a body moving along a straight path will continue to move in the same direction with the same speed even in the absence of external forces.
- A person standing in a bus moves backward when the bus moves forward. When the bus moves forward, the lower part of his body begins to move along with the bus while the upper part of his body continues to remain at rest due to inertia. That is why a person falls backward when the bus starts.
- $a = F/m = (5 \times 10^4)/(3 \times 10^7)$   
 Speed attained by the ship,  $v = \sqrt{2as} = 0.1\text{m/s}$ .
- If there is a loss of kinetic energy during a collision, it is called an inelastic collision. Collision between two vehicles is an example for inelastic collision.
- If two bodies stick together after the collision and move as a single body with a common velocity, then the collision is said to be perfectly inelastic collision. A mud thrown on a wall sticks to the wall, hence it is an example for perfectly inelastic collision.
- $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k} \quad \therefore r = \sqrt{x^2 + y^2 + z^2}$   
 $r = \sqrt{6^2 + 8^2 + 10^2} = 10\sqrt{2}\text{ m}$
- Displacement is the shortest distance between initial and final positions of the body.  
 $\vec{d} = OB = 100\text{m}$
- $\vec{a}_m = \lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{v}}{\Delta t}$
- Since velocity is constant so acceleration is zero.
- Definition of retardation.
- Only direction of displacement and velocity gets changed, acceleration is always directed vertically downward.
- $\vec{A} = -2\vec{i} + 3\vec{j} + \vec{k}$  ,  $\vec{B} = 3\vec{i} - \vec{j} + 4\vec{k}$   
 Displacement  $= \vec{d} = \vec{B} - \vec{A}$   
 $\vec{d} = 3\vec{i} - \vec{j} + 4\vec{k} + 2\vec{i} - 3\vec{j} - \vec{k}$   
 $\vec{d} = 5\vec{i} - 4\vec{j} + 3\vec{k}$

Topic-1

16. When length of the path of a particle is equal to the magnitude of the displacement of that particle, shape of the path possible is Straight line.
17. In part *cd* displacement-time graph shows constant slope i.e. velocity is constant. It means no acceleration or no force is acting on the body.
18. The shortest distance between two points is called displacement
19. For bodies moving in opposite direction, relative velocity is given by  $v_r = v_1 + v_2 \Rightarrow v_r = v + v = 2v$
20. In this case, man is at rest with respect to car.
21. At straight path, body can have constant velocity because direction of motion will remain same.
22. Displacement covered is zero so average velocity will be zero in this case.
23.  $s = 75 \text{ m}$ ,  $v = 15 \text{ ms}^{-1} \Rightarrow t = \frac{s}{v} = \frac{75}{15} = 5 \text{ s}$ , Now  $\bar{v}_{\text{avg}} = \frac{\bar{d}}{t} = \frac{25}{5} = 5 \text{ ms}^{-1}$
24. Area under v-t graph represents distance covered by the body.
25. As velocity of an object is increasing uniformly so it's acceleration is constant.
26. When change in velocity is uniform then uniform acceleration will be produced.
27. Distance = Area covered between graph and displacement axis =  $\frac{1}{2}(30+10)10 = 200 \text{ meter}$ .
28. In this case, initial velocity will be maximum, velocity will be zero at highest position, then velocity will increase when body comes back.
29. If the slope of velocity time graph gradually decreases, then the body is said to be moving with positive acceleration. i.e (graph is in 1<sup>st</sup> quadrant)
30.  $F = ma \Rightarrow a \propto F$
31.  $a = \frac{F}{m} = \frac{1}{0.1} = 10 \text{ m s}^{-2}$
32.  $F = ma = 10 \times 10 = 100 \text{ N}$
33. Newton's 1<sup>st</sup> law of motion defines force.
34.  $\frac{F'}{F} \propto \frac{a'}{a} \Rightarrow F' \propto \frac{10}{4} \times 12 = 30 \text{ N}$
35. Since forces are antiparallel  
 $F = F_{\text{upthrust}} - F_{\text{weight}}$   
 so  $= 10000 - 9000 \Rightarrow F = 1000 \text{ N}$
36. Rate of change of momentum is equal to applied force. So,  $F = \frac{\Delta p}{\Delta t}$
37.  $m_1 v_1 + m_2 v_2 = (m_1 + m_2) v$   
 $(m)(60) + m(-30) = (m + m) v$   
 $30m = 2mv \Rightarrow v = 15 \text{ cm s}^{-1}$
38. Statement of law of conservation of momentum.

Topic-1

39.  $\Delta p = F \times t \Rightarrow F = \frac{\Delta p}{t} = \frac{25}{0.05} = 500 \text{ N}$
40. If momentum remains constant then force will be zero because  $F = \frac{dP}{dt}$
41.  $F = \frac{\Delta p}{t} \Rightarrow t = \frac{mv}{F} = \frac{1 \times 30}{6} = 5 \text{ s}$
42.  $F = \frac{mv}{t} \Rightarrow v = \frac{F \times t}{m} = \frac{100 \times 10}{5} \Rightarrow v = 200 \text{ cms}^{-1}$
43. Area under "F - t" graph represents change in momentum.  
 $\Delta P = \text{area of triangle} + \text{area of trapezium}$   
 $\Delta P = 2 \times 2 + \frac{1}{2}(6+2) \times (4) \Rightarrow \Delta P = 20 \text{ kgms}^{-1}$
44. A gun after firing recoils due to conservation of momentum.
45.  $F = \frac{mv}{t} = \frac{0.15 \times 20}{0.1} = 30 \text{ N}$
46. For inelastic collision, momentum is still conserved.  
 Momentum before collision = momentum after collision  
 $m_1 u_1 + m_2 u_2 = (m_1 + m_2) v$   
 $(2)(8) + (4)(2) = (2+4)v \Rightarrow v = 4 \text{ ms}^{-1}$
47. Kinetic energy is conserved for elastic collision.  
 $\frac{1}{2} m v^2 + \frac{1}{2} m v^2 = m v^2$
48. This is a perfectly inelastic collision.



$$m(2v) + 3m(-v) = (m + 3m) v_1$$

$$-mv = 4mv_1$$

$$v_1 = -\frac{1}{4} v$$

49. Kinetic energy will remain conserved if energy loss due to compression is zero. It is possible in case of solid and hard object.
50. In this case  
 $m_1 \gg m_2$  and  $v_2 = 0$   
 so  $v_2' = 2v_1$
51. Momentum will be conserved.
52.  $h = \frac{v^2}{2g} = \frac{(19.6)^2}{2 \times 9.8} = 19.6 \text{ m}$



Topic-1

53.  $H = \frac{v_i^2 \sin^2 \theta}{2g} \Rightarrow H = \frac{4g}{4g} \frac{v_i^2 \sin^2 \theta}{2g}$
54. At maximum height projectile have minimum velocity as  $v_y = 0$
55.  $T = \frac{v_i \sin \theta}{g}$  is the time taken to reach maximum height?
56. For complementary angles ranges are same.  
If  $\theta_1 + \theta_2 = 90^\circ$ , then  $R_1 = R_2$
57. Path followed by projectile is known as its trajectory.
58. The horizontal component of the projectile's acceleration is zero
59. Horizontal component of projectiles' velocity remains same.
60. For an angle closer to  $45^\circ$ , range will be maximum; so in this case at  $\theta = 43^\circ$ , rang will be maximum as compared to range at other angles.
61. For the projectile's, horizontal acceleration will be zero and vertical acceleration =  $9.8 \text{ ms}^{-2}$
62. It has an initial horizontal velocity, which is same as that of train. It acquires a vertical component under the force of gravity.
63.  $s = x = t^3$ ,  $v = s/t = t^3/t = t^2$   
 $A = v/t = t^2/t = t$ ,  $A = t = 1$
64. The vertical velocity of ball thrown upward Decreases linearly with time.
65. An unpowered and unguided missile is called ballistic missile
66.  $v_f = v_i + a_i$   
 $a = \frac{v_f - v_i}{t} = \frac{1.5 \times 10^3 - 2 \times 10^3}{50} = \frac{-0.5 \times 10^3}{50} = -10 \text{ m/s}^2$
67. When a massive body collide with a light body then  
 $v'_1 = v_1 \Rightarrow v'_2 = 2v_1$
68.  $F = \frac{\Delta P}{\Delta t} \Rightarrow \Delta t = \frac{P_f - P_i}{F}$   
 $\Delta t = \frac{36 - 60}{12} = \frac{-24}{12} = 2 \text{ sec}$   
Time is always taken as positive.
69.  $T = \frac{2v_i \sin \theta}{g}$
70. Definition of uniform acceleration.
71. For finding the height of projectile, the equation used is  
 $2as = v_f^2 - v_i^2$

Topic-1

72.  $a = \frac{v_f - v_i}{t} = \frac{0 - 9.8}{10} = \frac{-9.8}{10} = -0.98 \text{ ms}^{-2}$
73.  $v_f = gt$
74. Initial K.E = E  
K.E at highest point =  $\frac{1}{2}mv^2 \cos^2 \theta = (E) \cos^2 45^\circ = E \times (0.7)^2 = 0.49E = 0.50E$
75.  $w = mg$
76.  $|\vec{d}| = 2r$  (along diameter)  
 $S = 2\pi r$  (distance along circle)  
 $\frac{|\vec{d}|}{S} = \frac{2r}{2\pi r} = 1 : \pi$
77.  $|\vec{V}_{\text{min}}| = \frac{|\vec{d}|}{t} = \frac{150}{18} = 8.33 \text{ ms}^{-1}$
78.  $S = \frac{1}{2}at^2 + v_i t = \frac{1}{2}(-2)(2)^2 + (15 \times 2)$   
 $S = -4 + 30 = 26 \text{ m}$
79. Value of ratio of displacement to distance is equal to 1 for straight path and less than 1 for others path.
80. Slope =  $a = \frac{\Delta v}{\Delta t} = \text{constant}$
81.  $\frac{|\vec{s}|}{s} \leq 1$  if initial and final point is same on a straight line then equal to 1 generally distance is greater than 1.
82.  $(m + M)V = mv + Mu \Rightarrow (m + M)V = mv + M(0) \Rightarrow V = \frac{m}{m + M}v$
83. The acceleration either increase or decrease.
84.  $2as = v_f^2 - v_i^2 \quad \because v_f = 0$   
 $a = \frac{-v_i^2}{2s} = \frac{-15^2}{2 \times 18} = 6.25 \text{ ms}^{-2}$
85. Newton first law of motion is also known is law of inertia.
86.  $m_1 v_1 + m_2 v_2 = (m_1 + m_2) v'$   
 $m(2v) + 3m(v) = 4mv' \Rightarrow v' = \frac{5v}{4}$
87. By using law of conservation of linear momentum.  
Total final momentum = Total initial momentum  
Total final momentum =  $mv - mv = 0$

Topic-1


88.  $F_y = ma_y = 0$   
 $a_y = 0 \Rightarrow v_y = \text{constant}$
89. Distance of moving body can't be zero and negative
90.  $R = R_{\max} \sin 2\theta$   
 If  $R = \frac{R_{\max}}{2}$   
 $\frac{R_{\max}}{2} = R_{\max} \sin 2\theta \Rightarrow \frac{1}{2} = \sin 2\theta \Rightarrow \theta = 15^\circ$
91.  $v = Vt + \frac{1}{2}gt^2$   
 $V = 0$   
 $s = \frac{1}{2}gt^2 = \frac{1}{2} \times 9.8t^2 = 4.9t^2$
92.  $J = F \times t = Ns$
93.  $4H = R \tan \theta$
94.  $F \times \Delta t = \Delta P$
95.  $R = R_{\max} \sin 2\theta$   
 $R_{\max} 2 = R_{\max} \sin 2\theta$   
 $\sin 2\theta = \frac{1}{2}$   
 $2\theta = \sin^{-1} (1/2)$   
 $\theta = 15^\circ$
96. Vertical velocity vs time graph for a projectile motion varies linearly
97. It is straight line distance between initial and final position
98.  $v_{av} = \frac{15}{10} = 1.5 \text{ ms}^{-1}$
99. Decrease in velocity per unit time is called deceleration
100. The slope of velocity-time graph represents acceleration
101. An object is accelerating when its speed or direction changes.
102. The slope of a straight-line graph of position vs. time is displacement
103. The coefficient of restitution for a perfectly elastic collision is 1.
104.  $F = \frac{m}{t} v = \frac{0.05}{1} \times 400 = 20$
105.  $P = mv = 2 \times 35 = 70 \text{ kg ms}^{-1}$
106. Slope of vt graph =  $a = \frac{\Delta v}{\Delta t} = \frac{\Delta v}{0} = \infty$

2  
TOPIC

WORK AND ENERGY

PRACTICE EXERCISE


- Q.1 A truck and a car are moving with equal velocity. On applying brakes, both will stop after a certain distance, then?  
 A. Truck will cover less distance before stopping  
 B. Car will cover less distance before stopping  
 C. Both will cover equal distance  
 D. None of the mentioned
- Q.2 What sort of energy does flying bird possess?  
 A. Potential energy  
 B. Kinetic energy  
 C. Elastic energy  
 D. Both potential and kinetic energy
- Q.3 When the velocity of an aero plane is doubled, the momentum \_\_\_\_\_  
 A. Remains unchanged  
 B. Is conserved  
 C. Becomes zero  
 D. Increases uniformly
- Q.4 A machine gun fires 60 bullets per minute, with a velocity of 700m/s. If each bullet has a mass of 50g, find the power developed by the gun.  
 A. 1225W  
 B. 12250W  
 C. 122.5W  
 D. 122W
- Q.5 A bullet fired from a gun can pierce a target due to its \_\_\_\_\_  
 A. Mechanical energy  
 B. Heat energy  
 C. Kinetic energy  
 D. Acceleration
- Q.6 Kinetic energy with any reference must be \_\_\_\_\_  
 A. Zero  
 B. Positive  
 C. Negative  
 D. Either negative or positive
- Q.7 At what angle work done will be maximum?  
 A.  $0^\circ$   
 B.  $45^\circ$   
 C.  $90^\circ$   
 D.  $30^\circ$
- Q.8 Which one of the following is a greater work?  
 A. +100 J  
 B. -1000 J  
 C. -100 J  
 D. +200 J
- Q.9 The figure shows the force distance curve of a body moving along a straight line. The work done by the force:



- A. 10 J  
 B. 30 J  
 C. 20 J  
 D. 40 J
- Q.10 Work done will be zero if angle between Force and displacement is:  
 A.  $0^\circ$   
 B.  $270^\circ$   
 C.  $60^\circ$   
 D.  $360^\circ$

Topic-2

- Q.11 A force  $2\hat{i} + \hat{j}$  has moved its point of application from (2,3) to (6,5). What is work done?  
 A. -10 B. -18  
 C. +10 D. +18
- Q.12 At what angle the work done will be half of its maximum value  
 A.  $0^\circ$  B.  $30^\circ$   
 C.  $45^\circ$  D.  $60^\circ$
- Q.13 A man pushes a wall with 50 (N) and it displaces it zero (m), his work is  
 A. Negative B. No work  
 C. Positive D. May all possible
- Q.14 If a mass of 5 Kg is lifted upto 5m height, what will be the work done against the gravitational field  
 A. 245 J B. 25 J  
 C. 49 J D. 98 J
- Q.15 A person walks 2 m with an acceleration of  $5 \text{ m s}^{-2}$ , holding an object of mass 2 kg. The net work done on the object is  
 A. 20 J B. 10 J  
 C. 5 J D. 0 J
- Q.16 A force of  $3\hat{i} + 2\hat{j} + 4\hat{k}$  N gives displacement of  $10\hat{j}$  m. The work done is  
 A. 20 J B. 26 J  
 C. 32 J D. Zero
- Q.17 A body travels displacement of 10 m by force of 5 N If work done is 25 J then angle between  $\vec{F}$  and  $\vec{d}$  is  
 A.  $0^\circ$  B.  $45^\circ$   
 C.  $30^\circ$  D.  $60^\circ$
- Q.18 A person holds a bucket of weight 60 N. He walks 7 m along the horizontal path and then climbs up a vertical distance of 5 m. The work done by the gravity is:  
 A. 300 N-m B. 720 N-m  
 C. 420 N-m D. None of these
- Q.19 A force F acting on an object varies with distance x as shown in fig. The work done by the force in moving the object from  $x = 0$  to  $x = 6$  m is




- A. 18 J B. 9 J  
 C. 13.5 J D. 4.5 J
- Q.20 If force and displacement of particle in direction of force are doubled. Work would be  
 A. Double B. 1/4 times  
 C. Half D. 4 times
- Q.21 A person is holding a bucket by applying a force of 10N. He moves a horizontal distance of 5m and then climbs up a vertical distance of 10m. Find the total work done by him?  
 A. 50J B. 100J  
 C. 150J D. 200J

Topic-2

- Q.22 A gardener pushes a lawn roller through a distance of 20m. If he applies a force of 20kg weight in a direction inclined at  $60^\circ$  to the ground, find the work done by him. ( $g=9.8\text{m/s}^2$ )  
 A. 400J B. 250J  
 C. 1960J D. 2514J
- Q.23 If velocity is double, then.  
 A. Momentum increase 4 times and K.E increases 2 times  
 B. Momentum increases 2 times and K.E increase constant  
 C. Momentum and K.E remain same  
 D. Momentum increases 2 times and K.E increases 4 times
- Q.24 What will be the ratio of kinetic energies of alpha particle and proton if their linear momentum will be same?  
 A. 18 : 1 B. 4 : 1  
 C. 1 : 4 D.  $10^4 : 1$
- Q.25 The Bodies of one kg and four kg have same kinetic energy. The ratio in their momenta will be  
 A. 1 : 2 B. 1 : 4  
 C. 1 : 16 D. 1 : 1
- Q.26 The velocity and momentum of a moving body are  $10,000 \text{ cm s}^{-1}$  and  $10,000 \text{ g cm s}^{-1}$  respectively. The K.E will be  
 A.  $5 \times 10^7 \text{ J}$  B.  $5 \times 10^8 \text{ J}$   
 C.  $5 \times 10^{-2} \text{ J}$  D.  $5 \times 10^9 \text{ J}$
- Q.27 If momentum of a moving object is doubled then its kinetic energy will be  
 A. Doubled B. Four times  
 C. Halved D. Same
- Q.28 The momentum and kinetic energy of a ball is numerically equal. The numerical value of velocity is  
 A.  $1 \text{ m s}^{-1}$  B.  $3 \text{ m s}^{-1}$   
 C.  $2 \text{ m s}^{-1}$  D.  $4 \text{ m s}^{-1}$
- Q.29 Kinetic energy of a body moving with speed of  $10 \text{ m s}^{-1}$  is 30 J. If its speed becomes 30 m/s its K.E will be  
 A. 10 J B. 90 J  
 C. 180 J D. 270 J
- Q.30 Car X is traveling at half the speed of car Y. Car X has twice mass of car Y. Which statement is correct?  
 A. Car X has half the kinetic energy of car Y  
 B. Car X has twice the kinetic energy of car Y  
 C. Car X has one quarter of the kinetic energy of car Y  
 D. The two cars have the same kinetic energy
- Q.31 A ball of mass 2 kg and another of mass 4 kg are dropped together from a 60 feet tall building. After a fall of 30 feet each towards earth, their respective kinetic energies will be in the ratio of:  
 A.  $\sqrt{2} : 1$  B. 1 : 2  
 C. 1 : 4 D.  $1 : \sqrt{2}$

- Q.32 A bomb of mass 30 kg at rest explodes into two pieces of masses 18 kg and 12 kg. The velocity of 18 kg mass is  $6 \text{ m s}^{-1}$ . The K.E of other mass is  
 A. 324 J B. 256 J  
 C. 486 J D. 524 J
- Q.33 The kinetic energy acquired by a body of mass  $m$  is travelling some distance  $s$ , starting from rest under the actions of a constant force, is directly proportional to  
 A.  $m^0$  B.  $m$   
 C.  $m^2$  D.  $m^{1/2}$
- Q.34 When force and displacement are in the same direction, the kinetic energy of the body  
 A. Increases B. Remains constant  
 C. Decreases D. Becomes zero
- Q.35 A truck and a car are moving with equal velocity. On applying brakes, both will stop after a certain distance, then?  
 A. Truck will cover less distance before stopping  
 B. Both will cover equal distance  
 C. Car will cover less distance before stopping  
 D. None of the mentioned
- Q.36 Potential energy per unit volume is given by  
 A.  $mgh$  B.  $gh$   
 C.  $\frac{mgh}{\rho}$  D.  $\rho gh$
- Q.37 A body is falling from a height  $h$ . After it has fallen a height  $h/2$ , it will possess  
 A. Only potential energy B. Half potential and half kinetic energy  
 C. Only kinetic energy D. More kinetic and less potential energy
- Q.38 Energy stored in the spring of watch is  
 A. Electrical energy B. potential energy  
 C. Kinetic energy D. Elastic potential energy
- Q.39 Initially four identical uniform block, each of mass  $m$  and thickness  $h$ , are spread on a table, work require to stack them on one another is



- A.  $mgh$  B.  $4mgh$   
 C.  $3mgh$  D.  $6mgh$
- Q.40 A stone is thrown up from the surface of earth when it reaches at maximum height. Its total energy is equal to  
 A.  $mgh$  B.  $\frac{1}{2}mv^2$   
 C. Zero D.  $2mgh$
- Q.41 Energy consumed by 60-watt bulb in 2 minutes is equal to  
 A. 7.2 kilo joules B. 120 joules  
 C. 720 joules D. 72000 joules
- Q.42 The consumption of energy by 60-watt bulb in 2 seconds is:  
 A. 20J B. 30J  
 C. 120 J D. 0.02 J

- Q.43 100 joules work has been done by an agency in 10 seconds. What is power of agency?  
 A. 1000-watt B. 10 watt  
 C. 100 D. 0.10 watt
- Q.44 A 500N force is applied on an object and it moves with velocity  $10 \text{ ms}^{-1}$ . If value of power is 2500 watt. Then what will be the angle between force and displacement  
 A.  $0^\circ$  B.  $90^\circ$   
 C.  $60^\circ$  D.  $23^\circ$
- Q.45 The time taken by an engine of power 10 kW to lift a mass of 200 kg to the height of 40 m is  
 A. 2 s B. 8 s  
 C. 4 s D. 16 s
- Q.46 To travel at a constant speed, a car engine provides 24 kW of useful power. The driving force on the car is 600 N. At what speed does it travel?  
 A.  $25 \text{ ms}^{-1}$  B.  $2.5 \text{ ms}^{-1}$   
 C.  $4.0 \text{ ms}^{-1}$  D.  $40 \text{ ms}^{-1}$
- Q.47 A force of 1000 N is needed to lift the hook of a crane at a steady velocity. The crane is then used to lift a load of mass 1000 kg at a velocity of  $0.50 \text{ m s}^{-1}$ . How much of the power developed by the motor of the crane is used in lifting the hook and the load? (Take  $g$  as  $10 \text{ m s}^{-2}$ ).  
 A. 5.0 kW B. 5.5 kW  
 C. 20 kW D. 22 kW
- Q.48 The power output of a lamp is 6W. How much energy does the lamp give out in 2 minutes?  
 A. 3 J B. 720 J  
 C. 120 J D. 430 J
- Q.49 A man  $M_1$  of mass 80 kg runs up a staircase in 15 s. Another man  $M_2$  also of mass 80 kg runs up the same staircase in 20 s. The ratio of the power developed by them will be:  
 A. 1 B. 16/9  
 C. 4/3 D. none of these
- Q.50 An engine pumps up 100 kg of water through a height of 10 m in 5s. Given that the efficiency of the engine is 60%, what is the power of the engine? (Take  $g = 10 \text{ m s}^{-2}$ )  
 A. 33 kW B. 0.33 kW  
 C. 3.3 kW D. 0.033 kW
- Q.51 An engine pumps out 40 kg of water in one second. The water comes out vertically upwards with a velocity of  $3 \text{ m s}^{-1}$ . What is the power of engine in kilowatt?  
 A. 1.2 kW B. 120 kW  
 C. 12 kW D. 1200 Kw
- Q.52 An elevator's motor produces 3000 W power. The speed with which it can lift a 1000 kg load is:  
 A.  $30.6 \text{ m s}^{-1}$  B.  $0.306 \text{ m s}^{-1}$   
 C.  $3.06 \text{ m s}^{-1}$  D.  $300.6 \text{ m s}^{-1}$
- Q.53 The power needed to lift a mass of 5000g to height of 1 m in 2 second is  
 A. 2.45-watt B. 24.5 watt  
 C. 245-watt D. 2.45 k watt
- Q.54 An engine pulls a car of mass 1500 kg on a level road at a constant speed of  $5 \text{ ms}^{-1}$ . If the frictional force is 500 N, what power does the engine generate?  
 A. 5.0 kW B. 10 kW  
 C. 2.5 kW D. 12.5 kW

Topic-2


- Q.55 Work done on a body equals change in its \_\_\_\_\_ energy.  
 A. Total  
 B. Kinetic  
 C. Potential  
 D. All of these
- Q.56 If a certain force acts on an object and changes its kinetic energy from 65 J to 130 J, then work done by the force will be:  
 A. 92.5 J  
 B. 65J  
 C. 97.5 J  
 D. 130 J
- Q.57 The amount of work required to stop a moving object is equal to:  
 A. The velocity of the object  
 B. The kinetic energy of the object  
 C. The mass of the object times its velocity  
 D. The mass of the object times its acceleration
- Q.58 The work energy principle is valid for  
 A. Change in K.E  
 B. Change in E.P.E  
 C. Change in P.E  
 D. All type of energies
- Q.59 Mathematical form of work energy principle is  
 A.  $Fd = \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2$   
 B.  $Fd = \frac{1}{2}mv_f - \frac{1}{2}mv_i$   
 C.  $Fd = \frac{1}{2}mv_f^2 + \frac{1}{2}mv_i^2$   
 D.  $Fd = \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2$

PAST PAPER MCQs


- Q.60 A force  $2i + j$  has moved its point of application from (2,3) to (6,5). What is work done? (MCAT 2008)  
 A. -10  
 B. -18  
 C. +10  
 D. +18
- Q.61 100 joules work has been done by an agency in 10 seconds. What is power of agency? (MCAT 2008)  
 A. 1000-watt  
 B. 10 watt  
 C. 100  
 D. 0.10 watt
- Q.62 Work done on a body equals change in its \_\_\_\_\_ energy. (MCAT 2008)  
 A. Total  
 B. Kinetic.  
 C. Potential  
 D. All of these
- Q.63 If a certain force acts on an object and changes its kinetic energy from 65 J to 130 J, then work done by the force will be: (MCAT 2009)  
 A. 92.5 J  
 B. 65J  
 C. 97.5 J  
 D. 130 J
- Q.64 The consumption of energy by 60-watt bulb in 2 seconds is: (MCAT 2010)  
 A. 20J  
 B. 30J  
 C. 120 J  
 D. 0.02 J
- Q.65 Which one of the following is a non-conservative force? (MCAT 2010)  
 A. Electric force  
 B. Gravitational force  
 C. Elastic spring force  
 D. Frictional force
- Q.66 If velocity is double, then. (MCAT 2010)  
 A. Momentum increase 4 times and K.E increases 2 times  
 B. Momentum increases 2 times and K.E increase constant  
 C. Momentum and K.E remain same  
 D. Momentum increases 2 times and K.E increases 4 times

Topic-2

- Q.67 The heat energy dissipated by 40 watts also in one hour is (ETE A 2010)  
 A. 1440 J  
 B. 14400 J  
 C. 144000 J  
 D. 1440.000 J
- Q.68 The gravitational potential energy per unit mass is called: (ETE A 2010)  
 A. Gravitational potential  
 B. Absolute P.E  
 C. P.E  
 D. Potential hill
- Q.69 A 6.0 kg block is released from rest 80m above the ground. When it has fallen 60m its kinetic energy is approximately: (ETE A 2015)  
 A. 4800 J  
 B. 3500 J  
 C. 1200 J  
 D. 120 J
- Q.70 Potential energy per unit volume is given by (MDCAT 2016)  
 A. mgh  
 B. gh  
 C.  $\frac{mgh}{\rho}$   
 D.  $\rho gh$
- Q.71 Total work done in figure (MDCAT 2017)



- A. 24 Nm  
 B. 8 Nm  
 C. 16 Nm  
 D. Zero Nm
- Q.72 Work done will be zero if angle between Force and displacement is: (MDCAT 2017)  
 A.  $0^\circ$   
 B.  $270^\circ$   
 C.  $60^\circ$   
 D.  $360^\circ$
- Q.73 If mass 'm' is dropped from height 'h' vertically, f is the force of friction during downward motion and 'v' is the velocity at bottom, following equation will be hold: (MDCAT 2017)  
 A.  $\frac{1}{2}mv^2 = mgh + fh$   
 B.  $fh = mgh + \frac{1}{2}mv^2$   
 C.  $mgh = \frac{1}{2}mv^2 - fh$   
 D.  $mgh = \frac{1}{2}mv^2 + fh$
- Q.74 At what angle work done will be maximum? (MDCAT 2017)  
 A.  $0^\circ$   
 B.  $45^\circ$   
 C.  $90^\circ$   
 D.  $30^\circ$
- Q.75 Which one of the following is a greater work? (MDCAT 2017)  
 A. +100 J  
 B. -1000 J  
 C. -100 J  
 D. +200 J
- Q.76 The figure shows the force distance curve of a body moving along a straight line. The work done by the force: (MDCAT 2017)



- A. 10 J  
 B. 30 J  
 C. 20 J  
 D. 40 J

## Topic-2

- Q.77 A man of mass 60 kg climbs up a 20m long staircase to the top of a building 10m high. What is the work done by him: Take  $g = 10\text{ms}^{-2}$  (ETEA 2017)  
 A. 12 KJ  
 B. 6 KJ  
 C. 3 KJ  
 D. None he above
- Q.78 When a force retards the motion of a body the work done is: (ETEA 2017)  
 A. Zero  
 B. Negative  
 C. Positive  
 D. Positive or negative depending upon the magnitude of force and displacement
- Q.79 An engine pumps out 40 kg of water in one second. The water comes out vertically upwards with a velocity of  $3\text{ms}^{-1}$ , the power of engine in kilowatt is: (ETEA 2017)  
 A. 1.2 kW  
 B. 12 kW  
 C. 120 kW  
 D. 1200 kW
- Q.80 Two boys weighing in the ratio 4:5 goes up stair taking time in the ratio 5:4. The ratio of their power is: (ETEA 2017)  
 A. 1  
 B. 16/25  
 C. 25/16  
 D. 4/5
- Q.81 Energy consumed by 60-watt bulb in 2 minutes is equal to (MDCAT 2018)  
 A. 7.2 kilo joules  
 B. 120 joules  
 C. 720 joules  
 D. 72000 joules
- Q.82 A stone of mass 2.0 kg is dropped from a rest position 5.0m above the ground. What is its velocity at a height of 3.0m above the ground? (MDCAT 2018)  
 A. 12.5m/s  
 B. 9.3m/s  
 C. 6.3m/s  
 D. 16.0m/s
- Q.83 The rate at which work is being done is called: (MDCAT 2018)  
 A. Power  
 B. Density  
 C. Energy  
 D. Force
- Q.84 A man has a mass of 80 kg. He ties himself to one end of rope which passes over a single fixed pulley. He pulls on the other end of the rope to lift himself up at an average speed of  $50\text{cms}^{-1}$ . What is the average useful power at which he is working? (ETEA 2018)  
 A. 40W  
 B. 0.39kW  
 C. 40kW  
 D. 39kW
- Q.85 If the momentum of a body decreases by 20% the percentage decreases in K.E will be: (ETEA 2018)  
 A. 44%  
 B. 36%  
 C. 28%  
 D. 20%
- Q.86 A man carries a 1 kg body 10m horizontally on a level ground. The work done by the man is: (ETEA 2018)  
 A. 10 J  
 B. 1 J  
 C. 0 J  
 D. 5 J
- Q.87 An automobile is moving forwards with uniform velocity due to the force exerted by its engine. If that force is double with the velocity remaining constant what happens to the total power? (MDCAT 2018)  
 A. It does not change  
 B. It is halved  
 C. It is squared  
 D. It is doubled

## Topic-2

## Work and Energy

- Q.88 Which of the following statement shows that no work is done? (MDCAT 2019)  
 A. Pushing a car to start it moving  
 B. Lifting the weights  
 C. Writing an essay on a page  
 D. The moon orbiting the earths
- Q.89 A 1.75 m heighted weight-lifter raises weights with a mass of 50 kg to a height of .5m above his head. How much work is being done by him? ( $g=10\text{ms}^{-2}$ ) (NMDCAT 2020)  
 A. 2125J  
 B. 2500 J  
 C. 50J  
 D. 1225 J
- Q.90 When the speed of your car is halved, by what factor does its kinetic energy decreases? (NMDCAT 2020)  
 A.  $\frac{1}{2}$   
 B.  $\frac{1}{4}$   
 C.  $\frac{1}{8}$   
 D.  $\frac{1}{6}$
- Q.91 Which of the following force is non-conservative force? (NMDCAT 2020)  
 A. Fractional force  
 B. Gravitation force  
 C. Electric force  
 D. Elastic spring force
- Q.92 The area under force – displacement graph gives us: (NUMS 2020)  
 A. Displacement  
 B. Power  
 C. Work  
 D. Acceleration
- Q.93 Kilowatt-hour is unit of? (NUMS 2020)  
 A. Electric energy  
 B. Power  
 C. Momentum  
 D. Torque
- Q.94 A field in which the work is done in a moving a body along a closed path is zero is called (NMDCAT 2021)  
 A. Electric field  
 B. Conservative field  
 C. Electromagnetic field  
 D. Gravitational field
- Q.95 When a force is parallel to the direction of motion of the body, then work done on the body is (NMDCAT 2021)  
 A. Zero  
 B. Minimum  
 C. Infinity  
 D. Maximum
- Q.96 If a body a mass of 2 kg is raised vertically through 2m, then the work done will be (NMDCAT 2021)  
 A. 38.2 J  
 B. 392.1 J  
 C. 39.2 J  
 D. 3.92 J
- Q.97 The relation between horsepower and watt is (NMDCAT 2021)  
 A. 1 hp = 546 watts  
 B. 1 hp = 746 watts  
 C. 1 hp = 1000 watts  
 D. 1 hp = 946 watts
- Q.98 The area under the force displacement graph represents (NMDCAT 2021)  
 A. Area  
 B. Work done  
 C. Power  
 D. None of these
- Q.99 A machine does 2500 J of work in 1 min. What is the power developed by the machine? (NMDCAT 2021)  
 A. 21 W  
 B. 42 W  
 C. 150 W  
 D. 2500 W

Topic-2

- Q.100 A moving car possesses  
 A. Sound energy  
 C. Heat energy
- Q.101 kWh is the unit of  
 A. Force  
 C. Time
- Q.102 2kg mass is uplifted by the machine through the height of 200m for 10 sec calculate power deliver to the load. ( $g = 10\text{m/s}^2$ )  
 A. 40W  
 C. 440W
- Q.103 Light body A and heavy body B have equal K. E of translation. Then  
 A. A has larger momentum than B  
 C. A and B have same momentum
- Q.104 If momentum is increased by 13% then K.E increase by  
 A. 37%  
 C. 3.47%
- B. Mechanical energy  
 D. Chemical energy
- B. Power  
 D. Energy
- B. 400W  
 D. 200W
- B. B has larger momentum than A  
 D. None
- B. 2.27%  
 D. None

(NMDCAT 2021)

(NMDCAT 2021)

(NMDCAT 2021)

(NMDCAT 2021)

(NMDCAT 2021)

# ANSWER KEY

## TOPIC-WISE MCQs & PAST PAPER MCQs

1	B	14	A	27	B	40	A	53	B	66	D	79	A	92	C
2	D	15	A	28	C	41	A	54	C	67	C	80	B	93	A
3	B	16	A	29	D	42	C	55	A	68	A	81	A	94	B
4	B	17	D	30	A	43	B	56	B	69	B	82	C	95	D
5	C	18	A	31	B	44	C	57	B	70	D	83	A	96	C
6	B	19	C	32	C	45	B	58	D	71	D	84	B	97	B
7	A	20	D	33	A	46	D	59	C	72	B	85	B	98	B
8	B	21	B	34	A	47	B	60	C	73	D	86	C	99	B
9	A	22	C	35	C	48	B	61	B	74	A	87	D	100	B
10	B	23	D	36	D	49	C	62	D	75	B	88	D	101	D
11	C	24	C	37	B	50	C	63	B	76	A	89	D	102	B
12	D	25	A	38	D	51	A	64	C	77	B	90	B	103	B
13	B	26	D	39	D	52	B	65	D	78	B	91	A	104	D

Topic-2

# EXPLANATORY NOTES

- Being lighter than a truck, the car has less kinetic energy. On applying brakes with the same force, the car will cover less distance before coming to rest.
- A flying bird possesses both kinetic and potential energies because it is at a certain height above the ground and moving with a certain velocity.
- If velocity of the aeroplane is doubled, its momentum is also doubled. However, combined momentum of air and aeroplane is conserved. As the momentum of aeroplane increases, the momentum of air also increases by an equal amount in the opposite direction.
- Mass of the bullets =  $60 \times 50 = 3000\text{g} = 3\text{kg}$   
 $v = 700\text{m/s}$   $t = 1\text{min} = 60\text{s}$   
 Power =  $W/t = (\text{Kinetic energy})/t = 12250\text{W}$ .
- The kinetic energy of a body by virtue of its motion is called kinetic energy. A moving object can do work. Therefore, a bullet fired from a gun can pierce a target due to its kinetic energy.
- $K.E = 1/2 mv^2$   
 It is always positive.
- $W = Fd \cos \theta$   
 $0 = 0^\circ$   
 $w = Fd \cos(0) \Rightarrow w = Fd$
- 1000 J is a greater work in given options.
- $W = (10)(1) - (10)(1) + (10)(1) = 10\text{ J}$
- $W = Fd \cos \theta = Fd \cos 270^\circ = 0$
- $\hat{d} = (6-2)\hat{i} + (5-3)\hat{j}$   
 $\hat{d} = 4\hat{i} + 2\hat{j}$   
 $W = \hat{F} \cdot \hat{d}$   
 $= (2\hat{i} + \hat{j}) \cdot (4\hat{i} + 2\hat{j}) = 8(\hat{i} \cdot \hat{i}) + 2(\hat{j} \cdot \hat{j})$   $\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = 1$   
 $= 8 + 2 = 10$
- $W = \frac{W_{\max}}{2}$   
 $W = W_{\max} \cos \theta \Rightarrow \frac{W_{\max}}{2} = W_{\max} \cos \theta \Rightarrow 0 = \cos^{-1} \left( \frac{1}{2} \right) = 60^\circ$
- $W = Fd \cos \theta$   
 since  $d = 0 \Rightarrow W = 0$
- $0 = 0^\circ \Rightarrow W = mgh \Rightarrow W = 245\text{J}$
- $W = Fd = (ma)d = 2 \times 5 \times 2 = 20\text{ J}$

Topic-2

16.  $W = \vec{F} \cdot \vec{d} = (3\hat{i} + 2\hat{j} + 4\hat{k}) \cdot (10\hat{j}) \Rightarrow W = 20\text{J}$
17.  $W = Fd \cos \theta$   
 $0 = \cos^{-1} \left( \frac{W}{Fd} \right) = \cos^{-1} \left( \frac{25}{10 \times 5} \right) = \cos^{-1} \left( \frac{1}{2} \right) \Rightarrow \theta = 60^\circ$
18.  $W_f = W_{\text{horizontal}} + W_{\text{vertical}}$   
 $W_f = 0 + 60 \times 5$   
 $W_f = 300 \text{ Nm}$
19.  $W = \text{Area under } F-x \text{ graph}$   
 $W = \frac{1}{2} (\text{Sum of parallel sides}) (\text{Perpendicular distance between parallel sides})$   
 $= \frac{1}{2} (6+3)(3) = 13.5 \text{ J}$
20. Work = (Force) (Displacement). If force and displacement both are doubled then work would be four times
21.  $F = 10\text{N}$ ,  $s = 5\text{m}$ ,  $\theta = 90^\circ$   
 Work done,  $W_1 = F s \cos \theta = 10 \times 5 \times \cos 90^\circ = 0$   
 For vertical motion, the angle between force and displacement is  $0^\circ$ .  
 Here,  $F = 10\text{N}$ ,  $s = 10\text{m}$ ,  $\theta = 0^\circ$   
 Work done,  $W_2 = 10 \times 10 \times \cos 0 = 100\text{J}$   
 Total work done =  $W_1 + W_2 = 100\text{J}$ .
22.  $F = w = mg = 20 \times 9.8\text{N}$   
 $s = 20\text{m}$ ,  $\theta = 60^\circ$   
 $W = F s \cos \theta = 20 \times 9.8 \times 20 \times \cos 60^\circ$   
 $W = 1960\text{J}$
23. If velocity is doubled then  $v' = 2v$   

$$K.E = \frac{1}{2} m v^2$$

$$P = mv$$

$$P' = m(2v) = 2mv$$

$$P' = 2P$$

$$K.E' = \frac{1}{2} m (2v)^2$$

$$= 4 \left[ \frac{1}{2} m v^2 \right]$$

$$K.E' = 4K.E$$
24. Mass of  $\alpha$ -particle =  $6.644 \times 10^{-27} \text{ kg}$   
 mass of proton =  $1.672 \times 10^{-27} \text{ kg}$   
 $P_\alpha = P_p$   
 $K.E = \frac{P^2}{2m}$ ,  $K.E \propto \frac{1}{m}$   
 $\frac{K.E_\alpha}{K.E_p} = \frac{m_p}{m_\alpha} = \frac{1.672 \times 10^{-27}}{4(1.672 \times 10^{-27})} = \frac{1}{4}$

Topic-2

25.  $K.E_1 = K.E_2$   
 $P_1^2 \propto m$   
 $\frac{P_1^2}{m_1} = \frac{P_2^2}{m_2} \Rightarrow \frac{P_1}{P_2} = \sqrt{\frac{m_1}{m_2}} \Rightarrow \frac{P_1}{P_2} = \sqrt{\frac{1}{4}} = \frac{1}{2}$
26.  $K.E = \frac{1}{2} P v \Rightarrow K.E = \frac{1}{2} \left( \frac{10,000 \times 10^2}{1000} \right) (10,000 \times 10^2)$   
 $= \frac{1}{2} \left( \frac{10000}{1000} \right) = 5\text{J} \Rightarrow K.E = 5 \times 10^8 \text{ J}$
27.  $K.E = \frac{P^2}{2m}$ , if  $P' = 2P$   
 $K.E = \frac{(2P)^2}{2m} = \frac{4P^2}{2m} = 4K.E \Rightarrow K.E' = 4K.E$
28.  $P = K.E$   
 $\frac{1}{2} m v = \frac{1}{2} m' v'^2$   
 $2v = v'^2 \Rightarrow v = 2\text{ms}^{-1}$
29.  $\frac{K.E_1}{K.E_2} = \frac{v_1^2}{v_2^2}$   
 $K.E_2 = K.E_1 \frac{v_2^2}{v_1^2} \Rightarrow K.E_2 = 30 \times \frac{900}{100} = 270\text{J}$
30.  $v_x = \frac{v_y}{2}$ ,  $m_x = 2m_y$   
 $K.E_x = \frac{1}{2} m_x v_x^2 = \frac{1}{2} (2m_y) \left( \frac{v_y}{2} \right)^2 = \frac{1}{2} (2m_y) \left( \frac{v_y^2}{4} \right) = \left( \frac{1}{2} m_y v_y^2 \right) \frac{1}{2} = \frac{K.E_y}{2}$
31. As both balls falling from same height so speed is same  $v = \sqrt{2g(h_1 - h_2)}$   
 $\frac{K.E_1}{K.E_2} = \frac{\frac{1}{2} m_1 v^2}{\frac{1}{2} m_2 v^2} = \frac{2}{4} = \frac{1}{2}$
32.  $m_1 v_1 = m_2 v_2 \therefore v_2 = \frac{18 \times 6}{12} = 9\text{ms}^{-1} \Rightarrow K.E = \frac{1}{2} \times 12(9)^2 = 486\text{J}$
33.  $K.E = \frac{1}{2} m v^2 \Rightarrow K.E \propto v^2 \therefore$  does not depend upon mass for a single body (As mass is constant)
34. When force and displacement are in the same direction, the kinetic energy of the body increases. The increase in kinetic energy is equal to the work done on the body.



Topic-2

35. Being lighter than a truck, the car has less kinetic energy. On applying brakes with the same force, the car will cover less distance before coming to rest.
36.  $\frac{P.E.}{V} = \frac{mgh}{V} = \rho gh$
37.  $P.E = mgh$   
 $P.E' = \frac{mgh}{2}$  ( $\because h' = \frac{h}{2}$ )  $\Rightarrow P.E' = \frac{P.E}{2}$   
 So at  $h/2$ , body has half P.E and half K.E.
38. Spring has elastic potential energy.
39.  $W = mg(0h) + mg(1h) + mg(2h) + mg(3h)$   
 $W = 0 + mgh + 2mgh + 3mgh \Rightarrow W = 6mgh$
40. At maximum height:  
 $T.E = K.E + P.E$   
 $= 0 + mgh = mgh$
41.  $W = P \times t = 60 \times 120 = 7200J = 7.2KJ$
42.  $P = \frac{W}{t} \Rightarrow W = P \times t = 60 \times 2 = 120J$
43.  $P = \frac{W}{t} = \frac{100}{10} = 10W$
44.  $P = \vec{F} \cdot \vec{v}$   
 $P = Fv \cos\theta$   
 $0 = \cos^{-1}\left(\frac{P}{Fv}\right) = \cos^{-1}\left(\frac{2500}{500 \times 10}\right) \Rightarrow \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ$
45.  $P = \frac{W}{t} = \frac{mgh}{t}$   
 $t = \frac{200 \times 9.8 \times 40}{10 \times 10^3} = 7.84s \Rightarrow t = 8s$
46.  $P = \vec{F} \cdot \vec{v}$   
 $P = Fv \cos\theta \Rightarrow \theta = 0^\circ$   
 $P = Fv \Rightarrow v = \frac{P}{F} = \frac{24000}{600} = 40 \text{ ms}^{-1}$
47.  $F = F_{\text{max}} + F_{\text{min}}$   
 $F = 1000 + 1000 \times 10 = 11000N \Rightarrow P = Fv$   
 $P = 11000 \times 0.50 = 5500W = 5.5kW$
48.  $t = 2 \times 60 = 120s$   
 $P = \frac{E}{t} \Rightarrow E = Pt = 6 \times 120 = 720J$

Topic-2

49.  $P = \frac{mgh}{t}$   
 As,  $M_1 = M_2 = m$   
 $h = \text{same}$   
 $\frac{P_1}{P_2} = \frac{t_2}{t_1} = \frac{20}{15} = \frac{4}{3}$
50.  $(P) = \frac{mgh}{t}$   
 $P = \frac{100 \times 10 \times 10}{5 \times 0.6} \Rightarrow P = 3333.3W \Rightarrow P = 3.3kW$
51.  $P = \frac{mgh}{t}$   
 $P = mgv = 40 \times 10 \times 3 = 1200W = 1.2 \times 10^3W = 1.2KW$
52.  $P = Fv$   
 $P = mgv \Rightarrow v = \frac{P}{mg} = \frac{3000}{1000 \times 10} = 0.306 \text{ms}^{-1}$
53.  $P = \frac{mgh}{t} = \frac{5000 \times 10 \times 1}{1000 \times 2} = \frac{50}{2} = 24.5W$
54.  $P = Fv = 500 \times 5 = 2500 = 2.5kW$
55. According to Work-energy principle  
 W.D on a body = change in K.E.  
 W.D on a body = change in P.E (W.D on spring)  
 W.D on a body = change in T.E (W.D on mass-spring system)
56. According to work-energy principle  
 $W.D = \Delta K.E = 130 - 65 = 65J$
57. Work energy principle(statement)
58. Work energy principle is valid for all kinds of mechanical energies
59. Amount of work = change in K.E
60.  $\vec{d} = (6-2)\hat{i} + (5-3)\hat{j}$   
 $\vec{d} = 4\hat{i} + 2\hat{j}$   
 $W = \vec{F} \cdot \vec{d}$   
 $= (2\hat{i} + \hat{j}) \cdot (4\hat{i} + 2\hat{j})$   
 $= 8(\hat{i}\hat{i}) + 2(\hat{j}\hat{j})$  ( $\hat{i}\hat{i} = \hat{j}\hat{j} = 1$ )  
 $= 8 + 2 = 10$

Topic-2

61.  $P = \frac{W}{t} = \frac{100}{10} = 10W$
62. According to Work-energy principle  
 W.D on a body = change in K.E  
 W.D on a body = change in P.E (W.D on spring)  
 W.D on a body = change in T.E (W.D on mass-spring system)
63. According to work-energy principle  
 W.D =  $\Delta K.E = 130 - 65 = 65J$
64.  $P = \frac{W}{t}$   
 $W = P \times t = 60 \times 2 = 120J$
65. Frictional force is a non-conservative force
66. If velocity is doubled then  $v' = 2v$   
 $K.E = \frac{1}{2}mv^2$   
 $P = mv$   
 $P' = m(2v) = 2mv$   
 $P' = 2P$   
 $K.E' = \frac{1}{2}m(2v)^2$   
 $= 4\left[\frac{1}{2}mv^2\right]$   
 $K.E' = 4K.E$   
 $W = P \times t = 40 \times 3600 = 144000 J$
67.  $W = P \times t$   $P = 40\text{watt}$   $t = 60 \times 60 = 3600\text{sec}$
68. Gravitational potential =  $U/m$
69.  $v = \sqrt{2gh}$ ,  $K.E = \frac{1}{2}mv^2$
70.  $\frac{P.E}{V} = \frac{mgh}{V} = \rho gh$
71. Work done in closed path is zero.
72.  $W = Fd \cos \theta = Fd \cos 270^\circ = 0$
73.  $mgh = \frac{1}{2}mv^2 + fh$
74.  $W = Fd \cos \theta$   
 $\theta = 0^\circ$   
 $w = Fd \cos(0)$   
 $w = Fd$
75. 1000 J is a greater work in given options.
76.  $W = (10)(1) - (10)(1) + (10)(1) = 10 J$
77.  $W.D = mgh = 60 \times 10 \times 10 \times 6000J = 6 \times 10^3 J = 6KJ$
78. Force and displacement are opposite in direction.
79.  $\vec{r} = Fx = mgv = 40 \times 10 \times 3 = 1200 = 1.2kW$

Topic-2

80.  $\frac{P_1}{P_2} = \frac{W_1/t_1}{W_2/t_2} = \frac{W_1 \times t_2}{W_2 \times t_1} = \frac{m_1 g h_1 \times t_2}{m_2 g h_2 \times t_1}$   
 Here  $h_2 = h_1 = h$   
 $\frac{P_1}{P_2} = \frac{m_1 \times t_2}{m_2 \times t_1} = \frac{4 \times 4}{5 \times 5} = \frac{16}{25}$
81.  $W = P \times t = 60 \times 120 = 7200J = 7.2KJ$
82.  $v_i = 0$   
 $v_f = \sqrt{2g(h_2 - h_1)}$   
 $v_f = \sqrt{2 \times 9.8(2)} \Rightarrow v_f = 6.26 \text{ ms}^{-1}$
83.  $P = \frac{W}{t}$
84.  $P = W/t = F \cdot v = mg \cdot v$  (As  $m = 80kg, g = 9.8ms^{-2}, v = 50m/sec = 0.5m/sec$ )  
 $P = mg \cdot v = 80 \times 9.8 \times 0.5 = 392W = 0.39kW$
85.  $K.E = \frac{P^2}{2m} \Rightarrow \%K.E = 2(\%P) - \frac{P^2}{100} \Rightarrow 2(20\%) - \left(\frac{400}{100}\right) \Rightarrow 40\% - 4\% = 36\%$
86.  $w = \vec{F} \cdot \vec{d} = mgh \cos 0$
87.  $P = \vec{F} \cdot \vec{v}$   
 $P \propto F \Rightarrow P' \propto F'$   
 if  $F' = 2F$   
 $P' = 2P$
88. Work done in a closed path is zero.
89.  $W = mgh = 50 \times 10 \times (1.75 + 0.5) = 1125J$
90.  $K.E = \frac{1}{2}mv^2 \Rightarrow v' = \frac{v}{2}$   
 $K.E' = \frac{1}{2}m\left(\frac{v}{2}\right)^2 = \frac{K.E}{4}$
91. Frictional force is path dependent so, it is non-conservative force.
92. The area under force – displacement graph gives us work.
93. Kilowatt-hour is unit of electric energy.
94. A field in which the work is done in a moving a body along a closed path is zero is called conservative field
95.  $\theta = 0^\circ$   
 $W = Fd \cos 0 = Fd$
96.  $W = P.E = mgh = 2 \times 9.8 \times 2 = 39.2J$

## Topic-2

1 hp = 746 watt

The area under the force displacement graph represents work done

$$99. \quad P = \frac{mgh}{t} = \frac{2500}{60} = 42W$$

100. A moving car possesses mechanical energy

101. kWh is the unit of energy

$$102. \quad P = \frac{mgh}{t}$$

$$103. \quad P = \sqrt{2m K.E}$$

$$104. \quad K.E = \frac{P^2}{2m}$$

New momentum = 1.13 P

$$K.E' = \frac{(1.13P)^2}{2m} = 1.27 K.E$$

$$\% \text{ change in } K.E = \frac{K.E' - K.E}{K.E} \times 100$$


$$= \frac{1.27 K.E - K.E}{K.E} \times 100 = 27.69\%$$

# 3 ROTATIONAL & CIRCULAR MOTION

## TOPIC

## PRACTICE EXERCISE

- Q.1 A girl attaches a rock to a string, which she then swings counter-clockwise in a horizontal circle. The string breaks at point P in the figure, which shows a bird's-eye view (as seen from above). Which path (A-E) will the rock follow?



- A. Path A  
B. Path B  
C. Path C  
D. Path D
- Q.2 The net acceleration of a particle in circular motion is \_\_\_\_\_  
A. Towards the centre  
B. Always along the radius  
C. Irregular  
D. Circular in motion
- Q.3 A stone tied to the end of string 80cm long is whirled in horizontal circle with a constant speed. If the stone makes 14 revolutions in 25 sec, what is the magnitude of acceleration of the stone?  
A. 680 cm/s<sup>2</sup>  
B. 860 cm/s<sup>2</sup>  
C. 720 cm/s<sup>2</sup>  
D. 990 cm/s<sup>2</sup>
- Q.4 Motion of the tip of second hand of the clock is an example for \_\_\_\_\_  
A. Uniform circular motion  
B. Projectile motion  
C. Motion in a plane with uniform velocity  
D. Motion in a plane with constant acceleration
- Q.5 Calculate the angular speed of flywheel making 420 revolutions per minute.  
A. 42300 rad/sec  
B. 1200 rad/sec  
C. 10/4200 rad/sec  
D. 44 rad/sec
- Q.6 Find the magnitude of the centripetal acceleration of a particle on the tip of a fan blade of length 0.30 metre, rotation at 1200 rev/minute  
A. 40 m/s<sup>2</sup>  
B. 4737.6 m/s<sup>2</sup>  
C. 245 m/s<sup>2</sup>  
D. 20 m/s<sup>2</sup>
- Q.7 The racing cars of masses  $m_1$  and  $m_2$  are moving in circles of radii  $r_1$  and  $r_2$  respectively. Their speeds are such that each makes a complete circle in the same length of time. The ratio of the angular speed of the first car to that of the second car is  
A.  $m_1 : m_2$   
B. 1 : 1  
C.  $r_1 r_2$   
D.  $m_1 r_1 : m_2 r_2$
- Q.8 The ratio of angular speeds of minute hand and hour hand of a watch is  
A. 6 : 1  
B. 12 : 1  
C. 1 : 12  
D. 1 : 6
- Q.9 The angular velocity of a particle rotating in a circular orbit 100 times per minute is  
A. 1.66 rad/s  
B. 10.47 rad/s  
C. 10.47 deg/s  
D. 60 deg/s

Topic-3


- Q.10 Angular speed of a particle increases from  $2\text{rads}^{-1}$  to  $4\text{rads}^{-1}$  across any two diametrically opposite positions. Its angular acceleration will be?  
 A.  $6\text{rads}^{-2}$   
 B.  $\frac{\pi}{6}\text{rads}^{-2}$   
 C.  $\frac{\pi}{3}\text{rads}^{-2}$   
 D.  $\pi\text{rads}^{-2}$
- Q.11 Radian is a unit of angular displacement which can also be measured in degrees. How many radians are equal to one degree?  
 A.  $\frac{180}{\pi}$   
 B.  $\frac{2\pi}{180}$   
 C.  $\frac{\pi}{180}$   
 D.  $\frac{\pi}{57.3}$
- Q.12 If a rotating body is moving counter clockwise, direction of angular velocity will be  
 A. Along linear velocity  
 B. Towards the center  
 C. Along the axis of rotation  
 D. Away from center
- Q.13 The ratio of angular frequency and linear frequency is  
 A.  $2\pi$   
 B.  $\pi$   
 C.  $\frac{1}{2\pi}$   
 D.  $\frac{\pi}{2}$
- Q.14 A flywheel gains a speed of 540 rpm in 6 second. Its angular acceleration is  
 A.  $3\pi\text{rad s}^{-2}$   
 B.  $6\pi\text{rad s}^{-2}$   
 C.  $9\pi\text{rad s}^{-2}$   
 D.  $12\pi\text{rad s}^{-2}$
- Q.15 The angular speed of a fly wheel making 120 revolutions/minute is  
 A.  $2\pi\text{rad/s}$   
 B.  $4\pi\text{rad/s}$   
 C.  $4\pi^2\text{rad/s}$   
 D.  $\pi\text{rad/s}$
- Q.16 For positive angular displacement the rotation would be  
 A. Clockwise  
 B. Anti-clockwise  
 C. Parallel  
 D. Perpendicular
- Q.17 Ten seconds after an electric fan is turned on, the fan rotates at 300rev/min. Its average angular acceleration is  
 A.  $30\text{rad/s}^2$   
 B.  $30\text{rev/s}^2$   
 C.  $3.14\text{rad/s}^2$   
 D.  $500\text{rev/s}^2$
- Q.18 The angular speed in radian/hour for daily rotation of the earth is  
 A.  $2\pi$   
 B.  $4\pi$   
 C.  $\frac{\pi}{6}$   
 D.  $\frac{\pi}{12}$
- Q.19 The shaft of a motor rotates at a constant angular speed of 360rev/min. Angle turned through in 1 sec in radian is  
 A.  $\pi$   
 B.  $6\pi$   
 C.  $3\pi$   
 D.  $12\pi$
- Q.20 The angular velocity of the minute hand of a clock is  
 A.  $\frac{2\pi}{60}\text{rads}^{-1}$   
 B.  $\frac{\pi}{24}\text{rads}^{-1}$   
 C.  $\frac{2\pi}{3600}\text{rads}^{-1}$   
 D.  $\frac{\pi}{3600}\text{rads}^{-1}$

Topic-3

- Q.21 If a wheel of radius  $r$  turns through an angle of  $30^\circ$ , then the distance through which any point on its rim moves is  
 A.  $\frac{\pi}{3}r$   
 B.  $\frac{\pi}{6}r$   
 C.  $\frac{\pi}{30}r$   
 D.  $\frac{\pi}{180}r$
- Q.22 A body moving along the circumference of a circle, completes two revolutions. If the radius of the circular path is  $r$ , total angular displacement covered is  
 A.  $\pi r$   
 B.  $2\pi r$   
 C. zero  
 D.  $4\pi$
- Q.23 When a body moves in a circle, the angle between its velocity  $\vec{v}$  and angular velocity  $\vec{\omega}$  is always  
 A.  $0^\circ$   
 B.  $180^\circ$   
 C.  $360^\circ$   
 D.  $90^\circ$
- Q.24 An object is moving along a circular path of radius 4m. What will be its angular displacement if it moves 14m on this circular path?  
 A. 5.5 radians  
 B. 5.0 radians  
 C. 3.5 radians  
 D. 4.5 radians
- Q.25 Which of the following gives the relationship between linear velocity and angular velocity?  
 A.  $v = r\omega$   
 B.  $v = s\omega$   
 C.  $v = r\theta$   
 D.  $v = s\theta$
- Q.26 A body moves in a circle with increasing angular velocity. At time  $t = 6\text{sec}$ , the angular velocity is  $27\text{rad/s}$ . What is the radius of circle made by the body where linear velocity is  $81\text{cm/s}$ ?  
 A. 6cm  
 B. 9cm  
 C. 3cm  
 D. 7cm
- Q.27 A wheel of radius 1 m covers an angular displacement of  $180^\circ$ . Its linear displacement is  
 A. 3.14 m  
 B. 6.28 m  
 C.  $\pi$  rad  
 D. 0.157 m
- Q.28 Linear velocity or tangential velocity of any particle moving in a circular path of radius 2 m with angular velocity  $8\text{rads}^{-1}$  will be:  
 A.  $16\text{ms}^{-1}$   
 B.  $10\text{ms}^{-1}$   
 C.  $4\text{ms}^{-1}$   
 D.  $6\text{ms}^{-1}$
- Q.29 The linear and angular velocities of a particle moving about the centre of a circle of radius  $r$ , are related by  
 A.  $\vec{v} = \vec{\omega} \times \vec{r}$   
 B.  $\vec{v} = \vec{r} \times \vec{\omega}$   
 C.  $\vec{v} \times \vec{\omega} = \vec{r}$   
 D.  $\vec{\omega} \times \vec{v} = \vec{r}$
- Q.30 If a car moves with a uniform speed of  $2\text{ms}^{-1}$  in a circle of radius 0.4 m. Its angular speed is  
 A.  $4\text{rad s}^{-1}$   
 B.  $1.6\text{rad s}^{-1}$   
 C.  $5\text{rad s}^{-1}$   
 D.  $2.8\text{ms}^{-1}$

Topic-3

Q.31 A disc of radius  $R=20$  cm is rotating about its axis with an angular velocity  $\omega = 20 \text{ rad s}^{-1}$  on a horizontal smooth surface. The linear speed of point C on the disc is



- A.  $1 \text{ m s}^{-1}$   
 B.  $4 \text{ m s}^{-1}$   
 C.  $2 \text{ m s}^{-1}$   
 D.  $4\pi \text{ m s}^{-1}$

Q.32 The length of the second hand of a watch is 1 cm. The velocity vector of the tip of the second hand in cm per second is

- A.  $2\pi$   
 B.  $\frac{2\pi}{60}$   
 C.  $\frac{2\pi}{12 \times 60}$   
 D.  $\frac{2\pi}{24 \times 60}$

Q.33 If the position vector of a particle is  $\vec{r} = (3\hat{i} + 4\hat{j})$  meter and its angular velocity is  $\vec{\omega} = (\hat{j} + 2\hat{k})$  rad/sec then its linear velocity is (in m/s).

- A.  $-(8\hat{i} - 6\hat{j} + 3\hat{k})$   
 B.  $(3\hat{i} - 6\hat{j} + 8\hat{k})$   
 C.  $-(3\hat{i} - 6\hat{j} + 6\hat{k})$   
 D.  $(6\hat{i} - 8\hat{j} + 3\hat{k})$

Q.34 For a particle in uniform circular motion the relation  $\vec{a} = r \alpha$  of accelerations hold. The acceleration 'a'

- A. Is centripetal acceleration  
 B. Is tangential acceleration  
 C. Is radial acceleration  
 D. Both A and B

Q.35 A point on the rim of a wheel 4 m in diameter has a velocity of  $1600 \text{ cm s}^{-1}$ . The angular velocity of the wheel is

- A.  $2 \text{ rad s}^{-1}$   
 B.  $4 \text{ rad s}^{-1}$   
 C.  $6 \text{ rad s}^{-1}$   
 D.  $8 \text{ rad s}^{-1}$

Q.36 The acceleration of a motor car is  $8 \text{ m/s}^2$ . If the diameter of its wheel be 2m. It's angular acceleration will be

- A.  $8 \text{ rad/s}^2$   
 B.  $10 \text{ m/s}^2$   
 C.  $16 \text{ rad/s}^2$   
 D.  $10 \text{ rad/s}^2$

Q.37 When a wheel 1m in diameter makes 30 rev/min, the linear speed of point on it's rim in  $\text{m s}^{-1}$  is

- A.  $2\pi$   
 B.  $\frac{\pi}{2}$   
 C.  $3\pi$   
 D.  $4\pi$

Q.38 A body is moving in a circular path with constant speed. The magnitude of tangential and centripetal acceleration are:

	Tangential	Centripetal
A.	$v^2$	0
B.	0	$v^2/r$
C.	0	0
D.	$v^2/r$	$v^2/r$

Topic-3

Q.39 Work done due to centripetal force for circular motion will be:

- A. Reduced  
 B. Half  
 C. Maximum  
 D. Zero

Q.40 A 500 kg car takes a round turn of radius 50 m with a velocity of 36 km/hr. The centripetal force is

- A. 250 N  
 B. 1000 N  
 C. 750 N  
 D. 1200 N

Q.41 A cycle wheel of radius 0.4 m completes one revolution in one second then the acceleration of a point on the cycle wheel will be

- A.  $0.8 \text{ m/s}^2$   
 B.  $1.6\pi^2 \text{ m/s}^2$   
 C.  $0.4 \text{ m/s}^2$   
 D.  $0.4\pi^2 \text{ m/s}^2$

Q.42 A stone of mass m tied to a string of length l is rotated in a circle with the other end of the string as the centre. The speed of the stone is v. If the string breaks, the stone will move

- A. Toward the centre of circle  
 B. Away from centre of circle  
 C. Along the tangent  
 D. All of these

Q.43 The mud flies off the tyre of a fast moving car in the direction

- A. Parallel to the moving tyre  
 B. Anti-parallel to the moving tyre  
 C. Tangent to the moving tyre  
 D. None of these

Q.44 The force required to bend the normally straight path of a particle into a circular path is called \_\_\_\_\_ force.

- A. Traveling  
 B. Centrifugal  
 C. Bending  
 D. Centripetal

Q.45 Which of the following is the correct vector form of centripetal force?

- A.  $m\omega\vec{r}$   
 B.  $m\omega^2\vec{r}$   
 C.  $-\frac{m\omega^2}{r}\hat{r}$   
 D.  $-m\omega^2\vec{r}$

Q.46 A body rotates with uniform speed in a circle of radius r and takes time T to complete one revolution. What are the magnitudes of the angular velocity  $\omega$ , the linear velocity v and the acceleration a?

Angular velocity, $\omega$	Linear velocity, v	Acceleration, $\alpha$
A. $\frac{1}{T}$	$\frac{4\pi r}{T}$	$\frac{2\pi r}{T^2}$
B. $\frac{2\pi}{T}$	$\frac{2\pi r}{T}$	$\frac{4\pi^2 r}{T^2}$
C. $\frac{2\pi}{T}$	$\frac{2\pi r}{T}$	$\frac{2\pi r}{T^2}$
D. $\frac{2\pi}{T}$	$\frac{4\pi r}{T}$	$\frac{4\pi^2 r}{T^2}$

Q.47 A particle revolves round a circular path with a constant speed. The acceleration of the particle is

- A. Along the circumference of the circle  
 B. Along the radius  
 C. Along the tangent  
 D. Zero

Topic-3

- Q.48 A car is moving with high velocity when it has a turn. A force acts on it outwardly because of  
 A. Centripetal force  
 B. Gravitational force  
 C. Centrifugal force  
 D. All the above
- Q.49 A cyclist turns around a curve at 15 miles/hour. If he turns at double the speed, the tendency to overturn is  
 A. Quadrupled  
 B. Halved  
 C. Unchanged  
 D. Doubled
- Q.50 A body of mass 5 kg is moving in a circle of radius 1m with an angular velocity of 2 radian/sec. The centripetal force is  
 A. 10 N  
 B. 20 N  
 C. 30 N  
 D. 40 N
- Q.51 The direction of centripetal force is  
 A. Towards the center  
 B. Along the tangential velocity  
 C. Away from center  
 D. Along the axis of rotation
- Q.52 A car of mass 1000kg traveling at 40 ms<sup>-1</sup> rounds a curve of radius 100m. what is the F<sub>c</sub>  
 A. 100 N  
 B. 1.6 × 10<sup>6</sup>N  
 C. 1.6 × 10<sup>4</sup>N  
 D. 8 × 10<sup>4</sup>N
- Q.53 If the radius of the circular path of a moving body is half without changing speed of rotation then the F<sub>c</sub> becomes  
 A. Half  
 B. One third  
 C. Doubled  
 D. One forth
- Q.54 A stone of mass 250 g is tied to the end of a string of length 1.0 m. It is whirled in a horizontal circle with a frequency of 30 rev./min. What is the tension in the string?  
 A.  $\frac{\pi^2}{4}$  N  
 B.  $\pi^2$  N  
 C.  $\frac{\pi^2}{2}$  N  
 D.  $2\pi^2$  N
- Q.55 A particle of mass m is executing uniform circular motion on a path of radius r. If p is the magnitude of its linear momentum. The radial force acting on the particle is  
 A.  $\frac{mp^2}{r}$   
 B.  $\frac{rm}{p}$   
 C.  $\frac{p^2}{rm}$   
 D.  $\frac{pmr}{r}$
- Q.56 A stone ties to the end of a string 1m long is whirled in a horizontal circle with a constant speed. If the stone makes 22 revolution in 44 seconds, what is the magnitude and direction of acceleration of the stone  
 A.  $\pi^2/4$  ms<sup>-2</sup> and direction along the radius towards the centre  
 B.  $\pi^2$  ms<sup>-2</sup> and direction along the radius towards the centre  
 C.  $\pi^2$  ms<sup>-2</sup> and direction along the radius away from the centre  
 D.  $\pi^2$  ms<sup>-2</sup> and direction along the tangent to the circle

PAST PAPER MCQs

- Q.57 The force required to bend the normally straight path of a particle into a circular path is called \_\_\_\_\_ force. (MCAT 2008)  
 A. Traveling  
 B. Centrifugal  
 C. Bending  
 D. Centripetal

Topic-3

- Q.58 Linear velocity or tangential velocity of any particle moving in a circular path of radius 2 m with angular velocity 8 rads<sup>-1</sup> will be: (MCAT 2009)  
 A. 16 ms<sup>-1</sup>  
 B. 10 ms<sup>-1</sup>  
 C. 4 ms<sup>-1</sup>  
 D. 6 ms<sup>-1</sup>
- Q.59 A wheel of radius 1 m covers an angular displacement of 180. Its linear displacement is (MCAT 2010)  
 A. 3.14 m  
 B. 6.28 m  
 C.  $\pi$  rad  
 D. 0.157 m
- Q.60 When a body moves in a circle the angle between its linear velocity and angular velocity is always: (ETEA 2010)  
 A. 0°  
 B. 180°  
 C. 360°  
 D. 90°
- Q.61 Radian is a unit of angular displacement which can also be measured in degrees. How many radlans are equal to one degree? (MCAT 2011)  
 A.  $\frac{180}{\pi}$   
 B.  $\frac{\pi}{180}$   
 C.  $\frac{2\pi}{180}$   
 D.  $\frac{\pi}{57.3}$
- Q.62 The centripetal acceleration of a car travelling at constant speed around a frictionless circular track: (ETEA 2013)  
 A. Is zero  
 B. Has constant magnitude but varying direction  
 C. Has constant direction but varying magnitude  
 D. Has varying magnitude and direction
- Q.63 A wheel starts from rest and has an angular acceleration of 4.0 rad/s<sup>2</sup>. When it has made 10 rev its angular velocity is: (ETEA 2016)  
 A. 16 rad/s  
 B. 22 rad/s  
 C. 32 rad/s  
 D. 250 rad/s
- Q.64 A child riding on a large merry-go-round, travels a distance of 3000m in a circle of diameter 40m. The total angle through which she revolves is: (ETEA 2016)  
 A. 50 rad  
 B. 75 rad  
 C. 150 rad  
 D. 314 rad
- Q.65 The angular velocity for daily rotation of the earth is: (ETEA 2016)  
 A.  $\frac{\pi}{3}$  radian hr<sup>-1</sup>  
 B.  $\frac{\pi}{6}$  radian hr<sup>-1</sup>  
 C.  $\frac{\pi}{12}$  radian hr<sup>-1</sup>  
 D.  $12\pi$  radian hr<sup>-1</sup>
- Q.66 A body moves in a circle with increasing angular velocity. At time t= 6sec, the angular velocity is 27rad/s. What is the radius of circle made by the body where linear velocity is 81cm/s? (MDCAT 2017)  
 A. 6cm  
 B. 9cm  
 C. 3cm  
 D. 7cm
- Q.67 Angular speed of minutes hand of mechanical watch is: (MDCAT 2017)  
 A.  $\pi / 30$  rad min<sup>-1</sup>  
 B.  $\pi / 2$  rad min<sup>-1</sup>  
 C.  $\pi$  rad mi:  
 D. None of these

Topic-3

- Q.68 A body is moving in a circular path with constant speed. The magnitude of tangential and centripetal acceleration are:  

	Tangential	Centripetal
A.	$rv^2$	0
B.	0	$v^2/r$
C.	0	0
D.	$v^2/r$	$v^2/r$

 (MDCAT 2017)
- Q.69 A centripetal force  $F$  acts on a body moving with angular speed  $\omega$ . If the angular speed is tripled, then the magnitude of centripetal force becomes:  
 A.  $8F$   
 B.  $9F$   
 C.  $3F$   
 D.  $4F$   
 (ETEAT 2017)
- Q.70 A fly wheel rotates at a constant speed of 3000 rpm (rev/min). The angle described by the shaft in radian in one second is:  
 A.  $2\pi$   
 B.  $30\pi$   
 C.  $100\pi$   
 D.  $300\pi$   
 (ETEAT 2017)
- Q.71 A wheel starts rotating from rest with angular acceleration of  $2 \text{ rads}^{-2}$  till its angular speed becomes  $6 \text{ rad/s}$ . The angular displacement of the wheel will be equal to  
 A.  $4 \text{ rad}$   
 B.  $12 \text{ rad}$   
 C.  $9 \text{ rad}$   
 D.  $7 \text{ rad}$   
 (MDCAT 2018)
- Q.72 Which of the following gives the relationship between linear velocity and angular velocity?  
 A.  $v = r\omega$   
 B.  $v = s\omega$   
 C.  $v = r\theta$   
 D.  $v = s\theta$   
 (MDCAT 2019)
- Q.73 An object is moving along a circular path of radius  $4\text{m}$ . What will be its angular displacement if it moves  $14\text{m}$  on this circular path?  
 A.  $5.5$  radians  
 B.  $5.0$  radians  
 C.  $3.5$  radians  
 D.  $4.5$  radians  
 (MDCAT 2019)
- Q.74 If during circular motion, tangential velocity of a body becomes double, then centripetal force becomes:  
 A. Double  
 B. One half  
 C. Four times  
 D. One fourth  
 (NMDCAT 2020)
- Q.75 Under what condition an object will have zero displacement but non zero distance?  
 A. Linear motion  
 B. Circular motion  
 C. Random motion  
 D. Oscillation  
 (NMDCAT 2020)
- Q.76 1 radian is equal to  
 A.  $57.1^\circ$   
 B.  $57.2^\circ$   
 C.  $57.3^\circ$   
 D.  $57.4^\circ$   
 (NUMS 2020)
- Q.77 A  $10\text{N}$  force moves a body around a circular path of radius  $50 \text{ cm}$ . What is work done in completing one revolution?  
 A.  $5\text{J}$   
 B.  $0$   
 C.  $31.42\text{J}$   
 D.  $500\text{J}$   
 (NMDCAT 2021)
- Q.78 Number of revolutions in  $720$  degrees.  
 A. 4 revolutions  
 B. 3 revolutions  
 C. 2 revolutions  
 D. 9 revolutions  
 (NMDCAT 2021)

Topic-3

- Q.79 Car moves in circular path due to  
 A. Centripetal force  
 B. Frictional force  
 C. Gravitational force  
 D. None  
 (NMDCAT 2021)
- Q.80 Radian is used to measure  
 A. Small angles  
 B. Large angles  
 C. Both  
 D. None  
 (NMDCAT 2021)
- Q.81 If earth covers  $360$  degrees in  $24$  hours, then how many degrees are in two hours  
 A.  $15^\circ$   
 B.  $30^\circ$   
 C.  $45^\circ$   
 D.  $60^\circ$   
 (NMDCAT 2021)
- Q.82 Earth completes one revolution in  
 A.  $24$  hours  
 B.  $6$  months  
 C.  $12$  months  
 D. None  
 (NMDCAT 2021)

ANSWER KEY

TOPIC-WISE MCQs & PAST PAPER MCQs

1	B	11	C	21	B	31	C	41	B	51	A	61	B	71	C	81	B
2	A	12	C	22	D	32	B	42	C	52	C	62	B	72	A	82	A
3	D	13	A	23	D	33	A	43	C	53	A	63	A	73	C		
4	A	14	A	24	C	34	B	44	D	54	A	64	C	74	C		
5	D	15	B	25	A	35	D	45	D	55	C	65	C	75	B		
6	B	16	B	26	C	36	A	46	B	56	B	66	C	76	C		
7	B	17	C	27	A	37	B	47	B	57	D	67	A	77	B		
8	B	18	D	28	A	38	B	48	C	58	A	68	D	78	C		
9	B	19	D	29	A	39	D	49	A	59	A	69	B	79	B		
10	C	20	C	30	C	40	B	50	B	60	D	70	C	80	B		

## EXPLANATORY NOTES

- Along the tangent.
- The net acceleration of a particle in a circular motion is towards the centre only if its speed is constant.
- $a = r\omega^2 = r(\theta/t)^2 = 80 \times (14 \times 3.14 \times 2)^2 / 25^2 = 990 \text{ cm/s}^2$
- If a particle moves along a circular path with a constant speed, then its motion is said to be a uniform circular motion. Hence, the motion of the tip of the second hand of a clock is an example for uniform circular motion.
- $\omega = 420 \text{ revolution/minute} = 420 \times 2\pi/60 \text{ rad/sec} = 44 \text{ rad/sec}$

$$a = r\omega^2$$

$$a = (0.30) \left( \frac{1200 \times 2\pi}{60} \right)^2 = 4737.6 \text{ ms}^{-2}$$

- Both cars complete one rotation after same time interval so have same angular velocity.

$$\text{Hence } \frac{\omega_1}{\omega_2} = 1:1$$

$$8. \frac{\omega_{\text{min hand}}}{\omega_{\text{hour hand}}} = \frac{\frac{1 \text{ rot}}{\text{hour}}}{\frac{1 \text{ rot}}{12 \text{ hours}}} = 12:1$$

$$9. \omega = \frac{100 \text{ rot}}{1 \text{ minute}} = \frac{100(2\pi) \text{ rad}}{60 \text{ s}} = 10.47 \text{ rads}^{-1}$$

- For semicircle angle covered must be  $\pi$  rad

$$2\alpha\theta = \omega_f^2 - \omega_i^2 \Rightarrow \alpha = \frac{4^2 - 2^2}{2(\pi)} = \frac{12}{2\pi} = \frac{6}{\pi} \text{ rads}^{-2}$$

$$11. 2\pi \text{ rad} = 360^\circ$$

$$1^\circ = \frac{2\pi}{360} \text{ rad} \Rightarrow 1^\circ = \frac{\pi}{180} \text{ rad}$$

- Angular velocity is a axial vector so it is always along the axis of rotation



$$13. \frac{\omega}{f} = \frac{2\pi f}{f} = 2\pi$$

$$14. \alpha = \frac{\Delta\omega}{\Delta t} = \frac{540 \times 2\pi}{60 \times 6} = \frac{540 \times 2\pi}{360} = 3\pi \text{ rad/s}^2$$

- $\omega = 120 \text{ rev/min} = 120 \times \frac{2\pi}{60} \text{ rad/sec} = 4\pi \text{ rad/sec}$
- According to convention in anti-clock wise rotation angular displacement is taken as positive.
- $\omega_1 = 0 \text{ rads}^{-1}$   
 $\omega_f = 300 \text{ rev/min} = 10\pi \text{ rads}^{-1}$   
 $\alpha = \frac{\omega_f - \omega_i}{t} = \frac{10\pi}{10} \Rightarrow \alpha = 3.14 \text{ rads}^{-2}$
- $\omega = \frac{\theta}{t} = \frac{2\pi}{24} = \frac{\pi}{12}$
- $\theta = \omega t = \frac{360 \times 2\pi}{60} \times 1 \Rightarrow \theta = 12\pi \text{ radian}$
- $\theta = \frac{2\pi}{T} = \frac{2\pi}{60 \times 60} \Rightarrow \theta = \frac{2\pi}{3600} \text{ rads}^{-1}$
- $\theta = 30^\circ = \frac{\pi}{6} \text{ rad}$   
 $S = r\theta \Rightarrow S = r \times \frac{\pi}{6}$
- one revolution =  $2\pi$  radiation  
two revolution =  $4\pi$  radiation  
 $S = r\theta$   
 $\theta = \frac{s}{r} = \frac{2\pi r}{r} \times 2$   
 $\theta = 4\pi$
- $\vec{v}$  and  $\vec{\omega}$  always perpendicular to each other
- $S = r\theta \Rightarrow \theta = \frac{S}{r} = \frac{14}{4} = 3.5 \text{ rad}$
- $v = r\omega$
- $v = r\omega \Rightarrow r = \frac{v}{\omega} = \frac{81}{27} = 3 \text{ cm}$
- $S = r\theta \quad \theta = 180^\circ$   
 $= 1 \times \pi \quad \theta = 180^\circ \times \frac{\pi}{180^\circ} = \pi \text{ rad}$   
 $= \pi m = 3.14 \text{ m}$
- $V = r\omega = 2 \times 8 = 16 \text{ m/s}$
- Relation between linear and angular velocity in vector form is  $\vec{v} = \vec{\omega} \times \vec{r}$
- $\omega = \frac{v}{r} = \frac{2}{0.4} = 5 \text{ rads}^{-1}$



Topic-3

$$v = r\omega \quad \therefore r = \frac{R}{2}$$

$$= 10 \times 10^{-2} \times 20 \Rightarrow v = 2 \text{ ms}^{-1}$$

$$v = r\omega$$

$$\therefore \omega = \frac{2\pi}{60} \text{ rad s}^{-1}$$

$$v = 1 \text{ cm} \times \frac{2\pi}{60} = \frac{2\pi}{60} \text{ cm s}^{-1}$$

$$33. \quad \vec{v} = \vec{\omega} \times \vec{r} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 1 & 2 \\ 3 & 4 & 0 \end{vmatrix} = -8\hat{i} + 6\hat{j} - 3\hat{k} = -(8\hat{i} - 6\hat{j} + 3\hat{k})$$

$$34. \quad a_t = r\alpha$$

Here  $a_t$  is tangential acceleration

$$35. \quad \omega = \frac{v}{r} = \frac{16}{2} = 8 \text{ rad s}^{-1}$$

$$36. \quad a = r\alpha \quad \therefore r = \frac{d}{2} = \frac{2}{2} = 1 \text{ m}$$

$$\alpha = \frac{a}{r} = \frac{8}{1} \Rightarrow \alpha = \frac{\text{rad}}{\text{sec}} = 8 \text{ rad/sec}^2$$

$$37. \quad v = r\omega \Rightarrow v = \frac{1}{2} \times \frac{30 \times 2\pi}{60} \Rightarrow v = \frac{\pi}{2} \text{ m s}^{-1}$$

$$38. \quad \text{For constant speed } \Delta v = 0, \quad a_t = \frac{\Delta v}{\Delta t} = 0 \Rightarrow a_c = \frac{v^2}{r}$$

$$39. \quad W = Fd \cos \theta$$

$$\theta = 90^\circ \Rightarrow W = Fd \cos 90^\circ = 0$$

$$40. \quad v = 36 \frac{\text{km}}{\text{h}} = 10 \frac{\text{m}}{\text{s}} \therefore F = \frac{mv^2}{r} = \frac{500 \times 100}{50} = 1000 \text{ N}$$

$$41. \quad a = r\omega^2 = r \left( \frac{\theta}{t} \right)^2 = 0.4 \times \left( \frac{2\pi}{1} \right)^2 = 0.4 \times 4\pi^2 = 1.6\pi^2 \text{ ms}^{-2}$$

42. When centripetal force vanishes objects moves along straight path i.e along tangent to circle.

43. The mud's flies off the tyre of a fast moving car in the direction tangent to the moving car.

44. Definition of centripetal force.

$$45. \quad \vec{F} = -mr\omega^2 \hat{r} = -mr\omega^2 \left( \frac{\vec{r}}{r} \right) = -m\omega^2 \vec{r}$$

$$46. \quad \text{As we know, } \omega = \frac{2\pi}{T} \quad \text{Also, } v = r\omega = r \left( \frac{2\pi}{T} \right) = \frac{2\pi r}{T}, \quad \text{And, } a = r\omega^2 = r \left( \frac{2\pi}{T} \right)^2 = \frac{4\pi^2 r}{T^2}$$

47. When body revolve with uniform speed then  $a_t$  and  $\alpha$  remain zero. Only centripetal acceleration presents in the body which is directed along the center of the circle.

Topic-3

48. A car is moving with high velocity when it has a turn. A force acts on it outwardly because of Centrifugal force

$$49. \quad F_c = \frac{mv^2}{r}, \quad \text{So, } F_c \propto v^2$$

If  $v$  increases to double then tendency to overturn will become four times

$$50. \quad F_c = mr\omega^2 = 5(1)(2)^2 = 20 \text{ N}$$

51. Centripetal force is towards the centre of circle.

$$52. \quad F_c = \frac{mv^2}{r} = \frac{1000(40)^2}{100} = 1.6 \times 10^4 \text{ N}$$

$$53. \quad F_c = mr\omega^2 \quad \text{here } \omega \text{ is constant}$$

$$\text{So, } F_c \propto r$$

$$54. \quad F_c = mr\omega^2 = \frac{250}{1000} \times 1 \times \frac{4\pi^2}{60 \times 60} \times 30 \times 30 \Rightarrow F_c = \frac{\pi^2}{4} \text{ N}$$

$$55. \quad \text{Radial force} = \frac{mv^2}{r} = \frac{m}{r} \left( \frac{p}{m} \right)^2 = \frac{p^2}{mr} \quad [\text{As } p = mv]$$

$$56. \quad a = \frac{v^2}{r} = \omega^2 r = 4\pi^2 n^2 r = 4\pi^2 \left( \frac{22}{44} \right)^2 \times 1 = \pi^2 \text{ m/s}^2 \text{ and its direction is always along the radius}$$

and towards the centre.

57. Definition of centripetal force.

$$58. \quad v = r\omega$$

$$= 2 \times 8 = 16 \text{ m/s}$$

59.

$$S = r\theta \quad \theta = 180^\circ$$

$$= 1 \times \pi \quad \theta = 180^\circ \times \frac{\pi}{180^\circ} = \pi \text{ rad}$$

$$= \pi \text{ m} = 3.14 \text{ m}$$

$$60. \quad \vec{v} = \vec{\omega} \times \vec{r} \Rightarrow \vec{v} \perp \vec{\omega}$$

61.

$$2\pi \text{ rad} = 360^\circ$$

$$1^\circ = \frac{2\pi}{360} \text{ rad} \Rightarrow 1^\circ = \frac{\pi}{180} \text{ rad}$$

$$62. \quad \vec{a}_c = \frac{v^2}{r} \vec{r}$$

$$63. \quad 2\alpha\theta = \omega_f^2 - \omega_i^2$$

$$64. \quad \theta = \frac{S}{r} = \frac{3000}{D/2} = \frac{3000}{40/2} = \frac{3000}{20} = 150 \text{ rad}$$

Topic-3

65.  $\omega = \frac{\theta}{t} = \frac{2\pi \text{ radian}}{24}$ ;  $\omega(\text{earth}) = \frac{\pi}{12} \text{ rad/hr}$
66.  $v = r\omega$   
 $r = \frac{v}{\omega} = \frac{81}{\frac{\pi}{27}} = 3\text{cm}$
67.  $\omega = \frac{\theta}{t} = \frac{2\pi}{60} = \frac{\pi}{30} \text{ rad/min}$
68. For constant speed  $\Delta v = 0$   
 $a_t = r\alpha \Rightarrow \alpha = 0$   
 $a_r = 0 \Rightarrow a_c = \frac{v^2}{r}$
69.  $F = mr\omega^2$ ,  $F' = mr\omega'^2 = F^2 = mr(3\omega)^2 = F' = 9mr\omega^2 = F' = 9F$
70.  $\theta = \omega t = 3000 \text{ rpm} = 3000 \text{ rev/min}$   $t = 3000 \times \frac{2\pi}{60} = 50 \times 2\pi = 100\pi$
71.  $2\theta\alpha = \omega_1^2 - \omega_2^2 \because \omega_2 = 0$   
 $\theta = \frac{\omega_1^2}{2\alpha} = \frac{6^2}{2(2)} = \frac{36}{4} = 9 \text{ rad}$
72.  $v = r\omega$
73.  $S = r\theta$   
 $\theta = \frac{S}{r} = \frac{14}{4} = 3.5 \text{ rad}$
74.  $F_c = \frac{mv^2}{r} \Rightarrow F_c \propto v^2$
75. During circular motion for complete rotation displacement = zero and distance =  $2\pi r$ .
76.  $360^\circ = 2\pi \text{ rad}$ .  
 $1 \text{ rad} = \frac{360^\circ}{2\pi} \Rightarrow 1 \text{ rad} = 57.3^\circ$
77. Centripetal force does zero work.
78.  $2\pi = 360^\circ = 1 \text{ rev}$   
 $720^\circ = \frac{1}{360} \times 720 = 2 \text{ rev}$
79. Car moves in circular path due to frictional force between tyre and road.
80.  $1 \text{ rad} = 57.3^\circ$   
 So, radian is used to measure large angles.
81.  $24 \text{ hours} = 360^\circ$   
 $2 \text{ hours} = \frac{360}{24} \times 2 = 30^\circ$
82. Earth completes one revolution in 24 hours.

4  
TOPIC


WAVES

PRACTICE EXERCISE

- Q.1 A simple pendulum is attached to the roof of a lift. If the time period of oscillation, when the lift is stationary is T, then the frequency of oscillation when the lift falls freely, will be \_\_\_\_\_  
 A. Zero  
 B. T  
 C. 1/T  
 D.  $\infty$
- Q.2 The length of a simple pendulum executing simple harmonic motion is increased by 21%. The percentage increase in the time period of the pendulum of increased length is?  
 A. 50%  
 B. 21%  
 C. 30%  
 D. 10.5%
- Q.3 If a simple pendulum oscillates with amplitude of 50mm and time period of 2s, then its maximum velocity is?  
 A. 0.10m/s  
 B. 0.16m/s  
 C. 0.24m/s  
 D. 0.32m/s
- Q.4 A spring of force constant k is cut into two pieces, such that one piece is double the length of the other. Then the long piece will have a force constant of \_\_\_\_\_  
 A. (2/3) k  
 B. (3/2) k  
 C. 3k  
 D. 6k
- Q.5 The phase difference between the acceleration of a particle executing simple harmonic motion and the instantaneous velocity is?  
 A.  $\pi$   
 B.  $0.707\pi$   
 C.  $\pi/2$   
 D.  $2\pi$
- Q.6 A siren emitting sound of frequency 800Hz is going away from a static listener with a speed of 30m/s. Frequency of the sound to be heard by the listener is (Take velocity of sound as 330m/s).  
 A. 733.3Hz  
 B. 644.8Hz  
 C. 481.2Hz  
 D. 286.5Hz
- Q.7 Lights can travel in a vacuum but not sound, because \_\_\_\_\_  
 A. Speed of sound is very slow than light  
 B. Light waves are electromagnetic in nature  
 C. Sound waves are electromagnetic in nature  
 D. Light waves are not electromagnetic in nature
- Q.8 When sound travels from air to water, which parameter does not change?  
 A. Wavelength  
 B. Frequency  
 C. Velocity  
 D. Temperature
- Q.9 A whistle giving out 450Hz approaches a stationary observer at a speed of 33m/s. The frequency heard by the observer in Hz is (speed of sound = 330m/s).  
 A. 409  
 B. 429  
 C. 517  
 D. 500
- Q.10 A boat at anchor is rocked by waves, whose crests are 100m apart and velocity is 25m/s. The boat bounces up once in every \_\_\_\_\_  
 A. 2500s  
 B. 75s  
 C. 4s  
 D. 0.25s


Topic- 4

Q.11 The diagram shows a transverse wave at a particular instant. The wave is traveling to the right. The frequency of the wave is 12.5 Hz.



At the instant shown the displacement is zero at point 'P'. What is shortest time to elapse before the displacement is zero at point 'Q'?

- A. 0.01 s  
B. 0.03 s  
C. 0.08 s  
D. 0.10 s
- Q.12 In a transverse wave the distance between a crest and a trough is equal to  
A.  $\frac{\lambda}{2}$   
B.  $\frac{\lambda}{4}$   
C.  $\lambda$   
D.  $2\lambda$
- Q.13 If a wave travelling at a speed of 130 m/s and has a wavelength of 5m. Then find out the frequency of the wave?  
A. 650 Hz  
B.  $3.8 \times 10^2$  Hz  
C. 20 Hz  
D. 26 Hz
- Q.14 A sinusoidal wave is traveling toward the right as shown. Which letter correctly labels the amplitude of the wave?



- A. A  
B. B  
C. D  
D. C
- Q.15 The sound of lightning flash is heard 3 second after the flash is seen. The distance of the lightning is 1020 metre. The speed of sound is:  
A. 340 m/s  
B. 1400 m/s  
C. 332 m/s  
D. none of these
- Q.16 When water waves pass from deep water into shallow water how do the frequency, wave length and speed change

Frequency	Wavelength	Speed
A. Increases	Decreases	No change
B. No change	Increases	Increases
C. No change	Decreases	Decreases
D. None of these		

- Q.17 Which of the following waves can be transmitted through solids, liquids and gases?  
A. Transverse waves  
B. Mechanical waves  
C. Electromagnetic waves  
D. Longitudinal waves

Topic- 4

- Q.18 If two sound waves having a phase difference of  $60^\circ$ , then they will have a path difference of  
A.  $\frac{\lambda}{6}$   
B.  $\lambda$   
C.  $\lambda/3$   
D.  $3\lambda$
- Q.19 All the points above the mean position in a rope  
A. Crest  
B. Trough  
C. Compression  
D. Rarefaction
- Q.20 The distance between consecutive compression and rarefaction in a spring  
A.  $\lambda$   
B.  $2\lambda$   
C.  $\lambda/2$   
D.  $3\lambda$
- Q.21 Velocity of sound in vacuum is:  
A.  $332 \text{ ms}^{-1}$   
B. Zero  
C.  $320 \text{ ms}^{-1}$   
D.  $224 \text{ ms}^{-1}$
- Q.22 Increase in velocity of sound in the air for  $1^\circ\text{C}$  rise in temperature is:  
A.  $1.61 \text{ ms}^{-1}$   
B.  $0.61 \text{ ms}^{-1}$   
C.  $61.0 \text{ ms}^{-1}$   
D.  $2.00 \text{ ms}^{-1}$
- Q.23 The velocity of sound in air would become double than its velocity at  $0^\circ\text{C}$  at temperature:  
A.  $313^\circ\text{C}$   
B.  $819^\circ\text{C}$   
C.  $586^\circ\text{C}$   
D.  $1172^\circ\text{C}$
- Q.24 Laplace found that the alternate compressions and rarefactions produced in sound waves follow:  
A. Isothermal law  
B. Isochoric law  
C. Adiabatic law  
D. All of the above
- Q.25 What is added when two waves superimpose?  
A. Amplitude  
B. Wavelength  
C. Velocities  
D. None of these
- Q.26 Which parameter get affected after superposition?  
A. Amplitude  
B. Wavelength  
C. Frequency  
D. Direction
- Q.27 The frequency of the first harmonic of a string stretched between two points is 100 Hz. The frequency of the third overtone is  
A. 200 Hz  
B. 400 Hz  
C. 300 Hz  
D. 600 Hz
- Q.28 "Stationary waves" are so called because in them  
A. The particles of the medium are not disturbed  
B. There occurs no flow of energy along the wave  
C. The particles of the medium do not execute SHM  
D. The interference effect can't be observed
- Q.29 The frequency of the nth mode of vibration of a string stretched by a tension T and having mass m and length is given by  
A.  $f_n = \frac{n}{2} \sqrt{\frac{T}{m\ell}}$   
B.  $f_n = \frac{n}{2\ell} \sqrt{\frac{T}{m}}$   
C.  $f_n = \frac{n}{2} \sqrt{\frac{T}{m}}$   
D.  $f_n = \frac{n}{2} \sqrt{\frac{T}{m}}$

## Topic- 4

- Q.30 If the string vibrates in 'n' loops, the wavelength is given by  
 A.  $\lambda_n = \frac{2}{nl}$  B.  $\lambda_n = \frac{l}{2n}$   
 C.  $\lambda_n = \frac{2l}{n}$  D. None of these
- Q.31 If the successive overtones of a vibrating string clamped at its ends are 280 Hz and 350 Hz, the frequency of fundamental is:  
 A. 350 Hz B. 140 Hz  
 C. 280 Hz D. 70 Hz
- Q.32 If the number of loops of a stationary wave are increasing, then  
 A.  $\lambda$  increases B.  $\lambda$  remains same  
 C.  $\lambda$  decreases D.  $\lambda$  may increase or decrease
- Q.33 A string of length 2m fixed between two supports vibrates in two loops. The distance between node and antinode is:  
 A. 50 cm B. 100 cm  
 C. 200 cm D. 10 cm
- Q.34 The distance between two particles in a wave motion in the same phase is  
 A.  $\frac{\lambda}{4}$  B.  $\frac{\lambda}{2}$   
 C.  $\frac{3\lambda}{4}$  D.  $\lambda$
- Q.35 The phase between two consecutive antinodes is:  
 A.  $\frac{\pi}{4}$  B.  $\frac{\pi}{2}$   
 C.  $\pi$  D.  $2\pi$
- Q.36 When the string vibrates in three loops then the length 'l' of the string is expressed as  
 A.  $l = \frac{3\lambda}{4}$  B.  $l = \frac{3\lambda}{2}$   
 C.  $l = \frac{\lambda}{2}$  D.  $l = \frac{2\lambda}{3}$
- Q.37 Consider a stretched string under tension and fixed at both ends. If the tension is doubled and the cross-sectional area halved, then the frequency becomes  
 A. Twice B. Half  
 C. Four times D. Eight times
- Q.38 When the antinodes are all at their extreme displacements, the energy stored is  
 A. K.E B. P.E  
 C. Thermal energy D. All of these
- Q.39 When an observer moves towards source with a velocity  $u_o$ , then the modified frequency ' $f_a$ ' becomes  
 A.  $f_a = f(v - u_o)$  B.  $f_a = f\left(\frac{v + u_o}{v}\right)$   
 C.  $f_a = \left(\frac{v + u_o}{2}\right)f$  D.  $f_a = \left(\frac{v - u_o}{v}\right)f$
- Q.40 Doppler's effect is not applicable for:  
 A. Microwaves B. Ultrasonic  
 C. Electromagnetic waves D. Standing waves

## Topic- 4

- Q.41 Doppler shift in frequency does not depend upon  
 A. The actual frequency of the wave  
 B. The velocity of the source  
 C. The distance of the source from the listener  
 D. The velocity of the observer
- Q.42 The source is moving towards a stationary observer then the pitch of the sound will  
 A. Sometimes increases and sometimes decreases  
 B. Decrease  
 C. Remains constant  
 D. Increase
- Q.43 The apparent frequency of the whistle of an engine changes in the ratio 6:5 as engine passes a stationary observer. If the speed of sound is 352 m/s. Then the speed of engine will be  
 A. 22 m/s B. 27 m/s  
 C. 32 m/s D. 36 m/s
- Q.44 A source of sound moves towards a stationary observer with a speed one third that of sound. If the frequency of the sound from the source is 100 Hz, the apparent frequency of the sound heard by the observer is  
 A. 67 Hz B. 100 Hz  
 C. 150 Hz D. 75 Hz
- Q.45 If a Radar system designed in accordance with the Doppler's effect, if an airplane is moving away from a Radar, the wavelength of the reflected wave from the air plane would be:  
 A. Smaller than the transmitting wave  
 B. Same as that of the transmitting wave  
 C. Larger than the transmitting wave  
 D. Either smaller or larger than the transmitting wave
- Q.46 Stars moving towards earth shows  
 A. Blue shift  
 B. Red shift  
 C. No shift  
 D. May be 'A' may be 'B' depending upon speed of stars
- Q.47 Bats navigate and find food by  
 A. Ultrasonic B. Echo location  
 C. Amplitude D. Refraction
- Q.48 Which one of the following explains that all the galaxies are receding from us?  
 A. White shift B. Black holes  
 C. Neutrons stars D. Red shift
- Q.49 A whistle giving out 450Hz approaches a stationary observer at a speed of 33m/s. The frequency heard by the observer in Hz is (speed of sound = 330m/s).  
 A. 409 B. 517  
 C. 429 D. 500
- Q.50 The source of sound generating of a frequency of 3 kHz reaches an observer with a speed of 0.5 times the velocity of sound in air. The frequency heard by the observer is?  
 A. 1 kHz B. 4 kHz  
 C. 3 kHz D. 6 kHz

Q.51 A whistle producing sound waves of frequencies 9500 Hz and it is approaching a stationary person with speed  $v \text{ ms}^{-1}$ . The velocity of sound in air is  $300 \text{ ms}^{-1}$ . If the person can hear frequencies upto a maximum of 10,000 Hz, the maximum value of  $v$  upto which he can hear the whistle is

- A.  $30 \text{ m s}^{-1}$
- B.  $10 \text{ m s}^{-1}$
- C.  $15 \text{ m s}^{-1}$
- D.  $20 \text{ m s}^{-1}$


Q.52 Total distance traveled by vibrating body in one vibration is equal to:

- A. A
- B. 4A
- C. 2A
- D. Zero


Q.53 A particle oscillating in simple harmonic motion is:

- A. Never in equilibrium because there is always a force
- B. In equilibrium at the center of its path because the acceleration is zero there
- C. Never in equilibrium because it is in motion
- D. In equilibrium at the ends of its path because its velocity is zero there

Q.54 Which one is the correct graph between acceleration 'a' and displacement 'x' for SHM?



Q.55 A restoring force  $F$  acts on a particle moving with simple harmonic motion. Which graph shows the variation with displacement  $r$  of force  $F$ ?



Q.56 The distance covered by a body in one complete vibration is 20 cm. What is the amplitude of body?

- A. 10 cm
- B. 15 cm
- C. 5 cm
- D. 7.5 cm

Q.57 A body moves with simple harmonic motion and makes  $n$ -complete oscillations in one second. What is its angular frequency?

- A.  $n \text{ rad s}^{-1}$
- B.  $\frac{1}{n} \text{ rad s}^{-1}$
- C.  $2\pi n \text{ rad s}^{-1}$
- D.  $\frac{2\pi}{n} \text{ rad s}^{-1}$

Q.58 The SI unit of force constant is identical with that of

- A. Force
- B. Surface tension
- C. Pressure
- D. Loudness

Q.59 A mass of 0.5 kg is suspended from a spring. The spring is stretched by 0.0980m. Its spring constant will be

- A.  $1 \text{ Nm}^{-1}$
- B.  $5 \text{ Nm}^{-1}$
- C.  $50 \text{ Nm}^{-1}$
- D.  $0.5 \text{ Nm}^{-1}$

- Q.60 Which of the following is a necessary and sufficient condition for S.H.M.?
- A. Proportionality between acceleration and displacement from equilibrium position
  - B. Constant acceleration
  - C. Proportionality between restoring force and displacement from equilibrium position
  - D. Constant period

PAST PAPER MCQs

- Q.61 A standing wave pattern is formed when the length of string is an integral multiple of \_\_\_\_\_ wavelength. (MCAT 2008)
- A. Triple
  - B. Half
  - C. Full
  - D. Double
- Q.62 Transverse waves cannot be setup in \_\_\_\_\_. (MCAT 2008)
- A. Metals
  - B. Fluids
  - C. Solids
  - D. Gases
- Q.63 Speed of the waves is equal to: (MCAT 2009)
- A.  $f\lambda$
  - B.  $\lambda T$
  - C. Both A and B
  - D.  $\lambda T$
- Q.64 What is it that we use to calculate the speeds of distant stars and galaxies? (MCAT 2009)
- A. Doppler Effect
  - B. Beats
  - C. Interference
  - D. All of the above
- Q.65 Speed of light, radio waves and microwaves in vacuum is: (MCAT 2009)
- A.  $3 \times 10^5 \text{ ms}^{-1}$
  - B.  $3 \times 10^6 \text{ ms}^{-1}$
  - C.  $3 \times 10^3 \text{ ms}^{-1}$
  - D.  $3 \times 10^8 \text{ ms}^{-1}$
- Q.66 An organ pipe closed at one end has a length of 25 cm. Wavelength of the fundamental note is (MCAT 2010)
- A. 25 cm
  - B. 100 cm
  - C. 50 cm
  - D. 75 cm
- Q.67 Two waves of same amplitude are traveling in the same direction and are out of phase, their resultant wave is: (MCAT 2010)
- A. Zero
  - B. Equal to sum of their amplitudes
  - C. Equal to difference of their amplitudes
  - D. Equal to half of their amplitude
- Q.68 The spectrum of a star's light is measured and the wavelength of one of the lines as the sodium's line is found to be 589 nm. The same line has the wavelength of 497 nm when observed in the laboratory. This means the star is (MCAT 2011)
- A. Moving away from the earth
  - B. Stationary
  - C. Moving towards the north
  - D. Revolving around the planet
- Q.69 When the source of sound moves towards the stationary observer, the value of apparent frequency ' $f_o$ ' is: (MCAT 2012)
- A.  $f_o = \left(\frac{v+u}{v}\right) f$
  - B.  $f_o = \left(\frac{v}{v+u_2}\right) f$
  - C.  $f_o = \left(\frac{v}{v-u}\right) f$
  - D.  $f_o = \left(\frac{v-u}{v}\right) f$

## Topic- 4

- Q.70 The displacement 'x' of a particle at time 't' is given by  $x = 10 \sin 4t$  the particle oscillates with period. (E TEA 2014)  
 A.  $\pi/10$ s  
 B.  $\pi/5$ s  
 C.  $\pi/4$ s  
 D.  $\pi/2$ s
- Q.71 In a vibrating cord the point where the particles are stationary is called (E TEA 2014)  
 A. Crest  
 B. Anti-node  
 C. Node  
 D. Trough
- Q.72 An observer moves with velocity 'u<sub>o</sub>' toward a stationary source, then the number waves received in one second is (M CAT 2015)  
 A.  $f' = f \left( \frac{v}{v+u_o} \right)$   
 B.  $f' = f \left( \frac{v+u_o}{v} \right)$   
 C.  $f' = f \left( \frac{v}{v-u_o} \right)$   
 D.  $f' = f \left( \frac{v-u_o}{v} \right)$
- Q.73 The ratio between the velocity of sound in air at 4 atm and that at 3 atm pressure would be: (E TEA 2015)  
 A. 1 : 1  
 B. 4 : 1  
 C. 1 : 4  
 D. 3 : 1
- Q.74 The red shift measurement of Doppler effect of galaxies indicate that the universe is (M CAT 2016)  
 A. Expanding  
 B. Stationary  
 C. Contracting  
 D. Oscillating
- Q.75 In stationary wave (E TEA 2016)  
 A. There is not transfer of energy  
 B. Energy is constant at all points  
 C. Phase is the same for all points  
 D. both (A) & (B)
- Q.76 If a wave travelling at a speed of 130 m/s and has a wavelength of 5m. Then find out the frequency of the wave? (M CAT 2017)  
 A. 650 Hz  
 B.  $3.8 \times 10^2$  Hz  
 C. 20 Hz  
 D. 26 Hz
- Q.77 A metallic wire of 2m length hooked between two points has tension of 10N. If mass per unit length of wire is 0.004 kg/s then fundamental frequency emitted by wire on vibration is: (M CAT 2017)  
 A. 12.5 Hz  
 B. 24 Hz  
 C. 48 Hz  
 D. 6.25 Hz
- Q.78 A source of sound moves towards a stationary observer with speed one third speed of sound. If the frequency of the sound from the source is 100 Hz, the apparent frequency of the sound heard by the observer is: (M CAT 2017)  
 A. 60 Hz  
 B. 200 Hz  
 C. 100 Hz  
 D. 150 Hz
- Q.79 In a stationary wave the distance between consecutive antinodes is 25 cm. If the wave velocity is 300 ms<sup>-1</sup> then the frequency of the wave will be: (E TEA 2017)  
 A. 150 Hz  
 B. 300 Hz  
 C. 600 Hz  
 D. 750 Hz
- Q.80 The speed of sound in air at NTP is 300m/s. If the air pressure become 4 times then the speed of the sound will be (E TEA 2017)  
 A. 150m/s  
 B. 300m/s  
 C. 600m/s  
 D. None

## Topic- 4

- Q.81 Standing waves are produced in 10m long stretched string. If the string vibrates in 5 segments and wave velocity is 20ms<sup>-1</sup>. Its frequency is: (E TEA 2017)  
 A. 2 Hz  
 B. 4 Hz  
 C. 5 Hz  
 D. 10 Hz
- Q.82 A shock wave is produced due to an earthquake which makes the building move in the direction of the shock wave. Which progressive wave would this be? (M CAT 2018)  
 A. Longitudinal wave  
 B. Material wave  
 C. Transverse wave  
 D. Particle wave
- Q.83 Which one of the following varies when an object execute simple harmonic motion? (E TEA 2018)  
 A. Angular frequency  
 B. Total energy  
 C. Force  
 D. Amplitude
- Q.84 Which one of the following is not a characteristic of stationary wave? (E TEA 2018)  
 A. Half wavelength is half the distance between the adjacent nodes  
 B. Amplitude is not the same  
 C. Phase is identical between two adjacent nodes  
 D. Energy of the stationary waves travels outwards
- Q.85 The wavelength of the electromagnetic wave having frequency of 3 kHz will be? (M CAT 2019)  
 A. 80 km  
 B. 100 km  
 C. 140 km  
 D. 120 km
- Q.86 What will be the expression for the observed frequency, if the source is moving towards the observer? (M CAT 2019)  
 A.  $f_o = \left( \frac{v}{v-u_s} \right) f$   
 B.  $f_o = \left( \frac{v}{v+u_s} \right) f$   
 C.  $f_o = \left( \frac{v}{v+u_o} \right) f$   
 D.  $f = \left( \frac{v}{v-u_o} \right) f_o$
- Q.87 The speed of sound in air is 332 m/s. The speed of sound at 22 °C will be: (N M CAT 2020)  
 A. 345.2m/s  
 B. 340 m/s  
 C. 350m/s  
 D. 330 m/s
- Q.88 Astronomers calculate speed of distant stars and galaxies using which of the following phenomena? (N M CAT 2020)  
 A. Beats  
 B. Interference  
 C. Superposition principle  
 D. Doppler effect
- Q.89 In a ripple tank, 40 waves pass through a certain point in 1 second. If the wavelength of the wave is 5cm. then speed of the wave is: (N M CAT 2020)  
 A. 0.5 ms<sup>-1</sup>  
 B. 1 ms<sup>-1</sup>  
 C. 1.5 ms<sup>-1</sup>  
 D. 2 ms<sup>-1</sup>
- Q.90 In ripple tank 40 waves pass through a certain point in one second. If the wavelength of the waves is 5cm, then find the speed of wave. (N UMS 2020)  
 A. 2.7 ms<sup>-1</sup>  
 B. 3 ms<sup>-1</sup>  
 C. 200 ms<sup>-1</sup>  
 D. 2 ms<sup>-1</sup>

Topic- 4

(NUMS 2020)

- Q.91 Trough of a wave acts as:  
 A. Concave lens  
 B. Convex lens  
 C. Convex mirror  
 D. Plane mirror  
 (NMDCAT 2021)
- Q.92 Doppler's effect shows?  
 A. Change in frequency due to relative motion between source and observer  
 B. Change in amplitude due to relative motion between source and observer  
 C. Change in frequency due to motion of source only  
 D. Change in frequency due to motion of observer only  
 (NMDCAT 2021)
- Q.93 Distance Between two Consecutive node is  
 A. Wave length  
 B. Wave length by 2  
 C. Wave length by 4  
 D. twice of Wave length  
 (NMDCAT 2021)
- Q.94 Stationary waves  
 A. Move forward  
 B. Move backwards  
 C. Don't Move  
 D. Move in both direction  
 (NMDCAT 2021)
- Q.95 Crest and trough are found in?  
 A. Transvers wave  
 B. Longitudinal wave  
 C. Progressive wave  
 D. None  
 (NMDCAT 2021)
- Q.96 Ratio of fundamental frequency of Open Organ pipe to closed organ pipe is  
 A. 1 : 2  
 B. 2 : 1  
 C. 1 : 4  
 D. 4 : 1  
 (NMDCAT 2021)

- Q.97 A whistler with velocity  $33 \text{ ms}^{-1}$  approaches towards a stationary observer with frequency 450Hz what is the apparent frequency heard by the observer:  
 A. 500  
 B. 400  
 C. 430  
 D. 450  
 (NMDCAT 2021)
- Q.98 Ripple tank is used to study features of  
 A. Particle  
 B. Wave  
 C. Light  
 D. Solid  
 (NMDCAT 2021)
- Q.99 Waves which deal with fast moving elementary particle  
 A. Transverse Waves  
 B. Longitudinal waves  
 C. Matter waves  
 D. Periodic waves  
 (NMDCAT 2021)
- Q.100 Infrasonic wave has frequency  
 A. Less than 20Hz  
 B. Greater than 20Hz  
 C. 20Hz  
 D. Greater than 20 kHz  
 (NMDCAT 2021)

Topic- 4


# ANSWER KEY

## TOPIC-WISE MCQs & PAST PAPER MCQs

1	A	11	A	21	B	31	D	41	C	51	C	61	B	71	C	81	C	91	A
2	D	12	A	22	B	32	C	42	D	52	B	62	B	72	B	82	A	92	A
3	B	13	D	23	B	33	A	43	C	53	B	63	C	73	A	83	C	93	B
4	B	14	C	24	C	34	D	44	C	54	B	64	A	74	A	84	D	94	C
5	C	15	A	25	A	35	C	45	C	55	D	65	D	75	A	85	B	95	A
6	A	16	C	26	A	36	B	46	A	56	C	66	B	76	D	86	A	96	B
7	B	17	D	27	B	37	A	47	B	57	C	67	A	77	A	87	A	97	A
8	B	18	A	28	B	38	B	48	D	58	B	68	A	78	D	88	D	98	B
9	D	19	A	29	B	39	B	49	D	59	C	69	C	79	C	89	D	99	C
10	C	20	A	30	C	40	D	50	D	60	A	70	D	80	B	90	D	100	A

# EXPLANATORY NOTES

- In a freely falling lift,  
 $g = 0$   
 $f = 1/2\pi \times \sqrt{\frac{g}{l}} = 1/2\pi \times \sqrt{\frac{0}{l}} = 0$
- %age increase in  $T = (\sqrt{1 + \% \text{age increase in } l} - 1) \times 100$   
 %age increase in  $T = \left(\sqrt{1 + \frac{21}{100}} - 1\right) \times 100$   
 %age increase in  $T = \left(\sqrt{\frac{121}{100}} - 1\right) \times 100 = 10\%$
- $v_{\max} = \omega A = 2\pi/T \times A$   
 $v_{\max} = (2 \times 3.14 \times 0.05)/2 = 0.16 \text{ m/s}$
- Force constant,  $k = F/x$   
 The length of the long piece is  $2x/3$   
 So, its force constant is  
 $k' = F/(2x/3) = 3F/2x = 3/2 \times k$
- The phase difference between the instantaneous velocity and acceleration of a particle executing simple harmonic motion is  $\pi/2$ .  
 $f' = v/(v+v_s) \Rightarrow f = 330/(330+30) \times 800 = 733.3 \text{ Hz}$
- Light can travel in a vacuum because light waves are electromagnetic in nature.
- Frequency remains unchanged when sound travels from air to water.
- $f' = v/(v-v_s) \times f$   
 $f' = 330/(330-33) \times 450 = 500 \text{ Hz}$
- $\lambda = 100 \text{ m}$ ,  $v = 25 \text{ m/s}$   
 $T = \lambda/v = 100/25 = 4 \text{ s}$
- $T = \frac{1}{12.5} = 0.08 \text{ s}$   
 for shortest time  $= \frac{0.08}{8} = 0.01 \text{ s}$
- Distance between two consecutive crests  $= \lambda$   
 Distance between consecutive crests and trough  $= \frac{\lambda}{2}$
- $v = f\lambda \Rightarrow f = \frac{v}{\lambda} = \frac{130}{5} = 26 \text{ Hz}$
- Maximum displacement from mean position
- Speed  $= \frac{\text{distance}}{\text{time}}$   
 $= \frac{1020}{3} = 340 \text{ m s}^{-1}$

- Frequency does not depend upon nature of material.
  - Longitudinal waves can be transmitted through all the three types of media
  - $\Delta\phi = 60^\circ = \pi/3$   
 $\Delta x = \lambda/2\pi \times \Delta\phi = \frac{\lambda}{2\pi} \times \frac{\pi}{3} = \frac{\lambda}{6}$
  - Wave characteristics.
  - Wave characteristics.
  - (Because speed of sound depends on medium).
  - $v_t = v_s + 0.61t$   
 So  $1^\circ\text{C}$  rise in temperature, velocity increases to  $0.61 \text{ m/s}$ .
  - $\frac{v_t}{v_s} = \sqrt{\frac{T}{273}} \Rightarrow 2 \frac{v_t}{v_s} = \sqrt{\frac{T}{273}}$   
 $4 = \frac{T}{273}$   
 $T = 1092 \text{ K} = 1092 - 273 = 819^\circ\text{C}$
  - According to Laplace, compression & rarefactions of sound waves follow adiabatic law.
  - When two waves super impose, amplitude is added.
  - After super position, amplitude is added.
  - Third overtone mean fourth harmonic  $f_4 = 4f_1$
  - Energy in a wave moves because of the motion of particle of the medium. The node always remains at rest so energy cannot flow past these point.
  - $f_n = \frac{n}{2\ell} \sqrt{\frac{T}{m}}$
  - $\lambda_n = \frac{2\ell}{n}$
  - $f' = f_2 - f_1$
  - $\lambda_n = \frac{2\ell}{n} \Rightarrow \lambda_n \propto \frac{1}{n}$
  - $\frac{\lambda}{2} = 1 \Rightarrow \lambda = 2 \text{ metre}$
- Distance between nearest node and antinode is
- $$\frac{\lambda}{4} = \frac{2}{4} = 0.5 \text{ m} = 50 \text{ cm}$$
- 
- Phase difference between two in phase points is  $\lambda, 2\lambda, 3\lambda, \dots$
  - Distance between two consecutive antinodes is  $\frac{\lambda}{2} = x$  and phase different  $= \frac{2\pi x}{\lambda} = \frac{2\pi}{\lambda} \cdot \frac{\lambda}{2} = \pi$



## Topic- 4

$$36. \quad \lambda = \frac{2\ell}{n} \Rightarrow n=3 \Rightarrow \lambda = \frac{2\ell}{3} \Rightarrow \ell = \frac{3\lambda}{2}$$

$$37. \quad f = \frac{1}{2l} \sqrt{\frac{T}{m}} \text{ where } m = \frac{\rho V}{l} = \frac{\rho(A)l}{l} = \rho A$$

$$\text{So, } f \propto \sqrt{\frac{T}{A}} \text{ and } f' \propto \sqrt{\frac{2T}{A}} \Rightarrow \frac{f'}{f} = 2 \text{ OR } f' = 2f$$

38. At extreme point energy stored is P.E.

$$39. \quad f_A = \left( \frac{v+u_s}{v} \right) f$$

40. Doppler's effect is applicable on light and sound waves.

41. Doppler shift in frequency does not depend upon distance of the source from listener.

42. As source is moving towards stationary observe.

$$\therefore f' = \frac{v}{v-u_s} f \Rightarrow f' > f \text{ so pitch will increase.}$$

$$43. \quad \frac{f'}{f''} = \frac{v+u}{v-u} = \frac{352+u}{352-u} = \frac{6}{5}$$

$$5(352+u) = 6(352-u)$$

$$1760+5u = 2112-6u$$

$$11u = 352 \Rightarrow u = 32 \text{ ms}^{-1}$$

$$f_A = \left( \frac{v}{v-u_s} \right) f \Rightarrow f_A = \left( \frac{v}{v-\frac{v}{3}} \right) 100 \Rightarrow f_A = \frac{3}{2} \times 100 = 150 \text{ Hz}$$

45. Source is moving away from observer so apparent frequency decreases and wavelength increases.

46. When star is moving towards earth, according to Doppler's shift wavelength decreases, as blue light has smaller wavelength so blue shift appears.

47. Bats use echolocation to navigate and find food in the dark. To echolocate, bats send out sound waves from their mouth or nose. When sound waves hit an object they produce echoes. Bats use this, to avoid flying into objects.

48. When galaxies or stars are receding from us its emitted light wavelength increase. So last colour of spectrum is red.

$$49. \quad f' = v/(v-v_s) \times f$$

$$f' = 330/(330-33) \times 450 = 500 \text{ Hz}$$

$$50. \quad f' = v/(v-v_s) \times f = v/(v-0.5v) \times 3 \text{ kHz} = 6 \text{ kHz}$$

## Topic- 4

$$51. \quad f_A = \left( \frac{v}{v-u_s} \right) f$$

$$10,000 = \left( \frac{300}{300-v} \right) 9500$$

$$v = 15 \text{ ms}^{-1}$$

52. Total distance in one vibration =  $4A$  as  $A+A+A+A=4A$  in complete vibration

53. At center of its path, no net force acts on it

54.  $a \propto -x$ , So graph lies in 2<sup>nd</sup> and 4<sup>th</sup> quadrant.

55.  $F \propto -x$

$$56. \quad S = \frac{20}{4} = 5 \text{ cm}$$

$$57. \quad \omega = 2\pi f \quad \therefore f = n$$

$$\omega = 2\pi n \frac{\text{rad}}{\text{s}}$$

58.  $F = kx \Rightarrow k = \frac{F}{x} = \text{Nm}^{-1}$  is same as that of surface tension.

59. As,  $F = kx$

$$k = \frac{F}{x} = \frac{mg}{x} \quad \therefore F = w = mg$$

$$= \frac{0.5 \times 9.8}{0.0980} = 50 \text{ Nm}^{-1}$$

60. Characteristic of SHM

61. A standing wave pattern is formed when length of string is an integral multiple of half wavelength e.g.,  $\ell_n = n \frac{\lambda_n}{2}$  here  $n=1,2,3,\dots$

62. Transverse waves cannot be set up in fluids, because they required rigidity of material.

$$63. \quad V = \frac{\lambda}{T} = f\lambda$$

64. We use Doppler's effect to calculate speed of distant stars and galaxies.

65. Radio waves and microwaves in vacuum travels with speed of light ( $3 \times 10^8 \text{ ms}^{-1}$ )

$$66. \quad \lambda = \frac{4\ell}{n} = \frac{4 \times 25 \text{ cm}}{1} = 100 \text{ cm}$$

67. When two waves are out of phase superpose of same amplitude their resultant will be zero.

$$68. \quad \lambda' = 589 \text{ nm}$$

$$\lambda = 497 \text{ nm}$$

$\lambda' > \lambda$  so star is moving away from earth.

## Topic- 4

69. When source move towards stationary observer  $f_o = \left(\frac{v}{v-u}\right)f$ .
70.  $2\pi f = \omega$   
 $\frac{2\pi}{T} = \omega \Rightarrow T = \frac{\pi}{\omega} \text{ sec}$
71. Definition.
72. When observer moves toward stationary source  $f' = f\left(\frac{v+u_o}{v}\right)$ .
73.  $\frac{v_{\text{air}}}{v_{\text{vac}}} = 1:1$  speed of sound is independent of pressure.
74.  $\lambda_{\text{red}} = \text{large}$ , at red shift, universe is expanding.
75. Because stationary wave is not a energy carrier.
76.  $v = f\lambda \Rightarrow f = \frac{v}{\lambda} = \frac{130}{5} = 26 \text{ Hz}$
77.  $f = \frac{1}{2l} \sqrt{\frac{F}{m}} = \frac{1}{2(2)} \sqrt{\frac{10}{0.04}} = 12.5 \text{ Hz}$
78.  $f' = \left(\frac{v}{v-u_s}\right) f \Rightarrow \left(\frac{v}{v-\frac{v}{3}}\right) 100 = \left(\frac{v}{\frac{3v-v}{3}}\right) 100 = \frac{3v}{2v} \times 100 = 150 \text{ Hz}$
79.  $\frac{\lambda}{2} = \frac{25}{100} \Rightarrow \lambda = 0.5 \text{ m} \Rightarrow f = \frac{v}{\lambda} = \frac{300}{0.5} = 600 \text{ Hz}$
80. Speed of sound is independent of pressure.
81. As  $n = 5$ ,  $l = 10 \text{ m} \Rightarrow \lambda n = \frac{2l}{n} = \frac{2 \times 10}{5} = 4 \text{ m}$   
 $f = \frac{v}{\lambda} = \frac{20}{4} = 5 \text{ Hz}$
82. Definition of longitudinal waves.
83.  $f \propto -x$
84. Energy of stationary wave travels outwards is not characteristics of stationary waves.
85.  $c = f\lambda \Rightarrow \lambda = \frac{c}{f} = \frac{3 \times 10^8}{3000} = 100 \text{ km}$
86.  $f_o = \left(\frac{v}{v-u_s}\right) f$
87.  $V_t = V_o + 0.6t^\circ\text{C} = 332 + 0.6 \times 22 = 332 + 13.2 = 345.2 \text{ ms}^{-1}$
88. Doppler's effect is applicable for light waves also. Speed of distant stars is measured by Doppler's effect.
89.  $v = f\lambda = (40)(0.05) = 2 \text{ m/s}$

## Topic- 4

90.  $v = f\lambda = 40 \times 5 \times 10^{-2} = 2 \text{ ms}^{-1}$
91. Trough of a wave like a concave lens in geometry.
92. Doppler's effect shows change in frequency due to relative motion between source and observer.
93. Distance Between two Consecutive node is wave length by 2
94. Stationary waves don't Move
95. Crest and trough are found in transvers wave
96.  $\frac{f_o}{f_s} = \frac{v}{2l} = 2:1$   
 $\frac{v}{4l}$
97.  $f' = \left(\frac{v}{v-u_s}\right) f$   
 $\frac{333}{333-33} \times 450 = 500 \text{ Hz}$
98. Ripple tank is used to study features of wave
99. Waves associated with moving particles are matter waves.
100. Infrasonic wave has frequency less than 20Hz

# 5 TOPIC

## THERMODYNAMICS

### PRACTICE EXERCISE

- Q.1 Heat was given to a body, which raises its temperature by  $1^\circ\text{C}$  is \_\_\_\_\_  
 A. Water equivalent  
 B. Temperature gradient  
 C. Thermal capacity  
 D. Specific heat
- Q.2 The thermal capacity of 40g of aluminium (specific heat =  $0.2 \text{ cal}^\circ\text{g}^{-1} \text{ C}^{-1}$ ) is \_\_\_\_\_  
 A.  $40 \text{ cal}^\circ\text{C}^{-1}$   
 B.  $160 \text{ cal}^\circ\text{C}^{-1}$   
 C.  $200 \text{ cal}^\circ\text{C}^{-1}$   
 D.  $8 \text{ cal}^\circ\text{C}^{-1}$
- Q.3 80g of water at  $30^\circ\text{C}$  is poured on a large block of ice which is at  $0^\circ\text{C}$ . The mass of ice that melts is \_\_\_\_\_  
 A. 30g  
 B. 80g  
 C. 150g  
 D. 1600g
- Q.4 Mud houses are cooler in summer and warmer in winter because \_\_\_\_\_  
 A. Mud is a good conductor of heat  
 B. Mud is a superconductor of heat  
 C. Mud is a bad conductor of heat  
 D. Mud can adapt according to temperature
- Q.5 Heat capacity of a substance is infinite. It means \_\_\_\_\_  
 A. Heat is given out  
 B. Heat is taken out  
 C. No change in temperature  
 D. Heat is first given out and then taken out
- Q.6 The change in internal energy, when a gas is cooled from  $927^\circ$  to  $27^\circ$  is?  
 A. 100%  
 B. 300%  
 C. 200%  
 D. 75%
- Q.7 Two cylinders of equal size are filled with equal amount of ideal diatomic gas at room temperature. Both the cylinders are fitted with pistons. In cylinder A the piston is free to move, while in cylinder B the piston is fixed. When same amount of heat is added to cylinder A raises by 20 K. What will be the rise temperature of gas in cylinder B?  
 A. 28K  
 B. 20K  
 C. 15K  
 D. 10K
- Q.8 The relation for the 1<sup>st</sup> law of thermodynamics can be expressed as:  
 A.  $\Delta Q = \Delta W$   
 B.  $\Delta Q = \Delta U + \Delta W$   
 C.  $\Delta Q = \Delta U$   
 D.  $\Delta Q = \frac{\Delta U}{\Delta W}$
- Q.9 Examples of first law of thermodynamics are  
 A. Working of bicycle pump  
 B. Brakes applied by an automobile  
 C. Human metabolism  
 D. All of these
- Q.10 1<sup>st</sup> law of thermodynamics is consequence of conservation of  
 A. Work  
 B. Heat  
 C. Energy  
 D. All of these
- Q.11 Which statement about the first law of thermodynamics is correct?  
 A. The heating of a system equal to the increase of its internal energy plus the work done on the system  
 B. The increase in the internal energy of a system equal the heating of the system minus the work done by the system  
 C. The increase in the internal energy of system equal the heating of the system plus the work done by the system  
 D. The work done on a system equals the increase of its thermal energy plus the heating of the system

## Topic-5

## Thermodynamics

- Q.12 The first law of thermodynamics may be expressed as shown.

$$\Delta U = Q + W$$

Where  $\Delta U$  is the change in internal energy,  $Q$  is the heating of the system,  $W$  is the work done on the system. A fixed mass of ideal gas at high pressure is contained in a balloon. The balloon suddenly bursts, causing the gas to expand and cool.


In this situation, which row describes the values of  $\Delta U$ ,  $Q$  and  $W$ ?

	$\Delta U$	$Q$	$W$
A.	negative	negative	positive
B.	negative	zero	negative
C.	positive	zero	negative
D.	positive	negative	positive

- Q.13 In a thermodynamic system working substance is an ideal gas, its internal energy is in the form of  
 A. Kinetic energy only  
 B. Potential energy  
 C. Kinetic and potential energy  
 D. None of these
- Q.14 In an ideal gas, the molecules possess  
 A. Only K.E  
 B. Only P.E  
 C. K.E and P.E both  
 D. Only gravitational energy
- Q.15 Which one is true for internal energy?  
 A. It is sum of all forms of molecular energies of a system  
 B. It is proportional to transnational K.E of the molecules  
 C. It is a state function of a system  
 D. All are correct
- Q.16 If two system X and Y are in thermal equilibrium. If X is heated at constant volume and Y is heated at constant pressure, and again finally maintained at thermal equilibrium, then heat  $Q$  given to the systems X and Y and internal energy  $U$  stored in the systems X and Y are  
 A.  $Q_x = Q_y$  and  $U_x = U_y$   
 B.  $Q_x = Q_y$  and  $U_x < U_y$   
 C.  $Q_x < Q_y$  and  $U_x < U_y$   
 D.  $Q_x < Q_y$  and  $U_x = U_y$
- Q.17 The internal energy of a body is maximum when its temperature is  
 A. 0 K  
 B. 273 K  
 C.  $-273 \text{ K}$   
 D.  $-273^\circ\text{C}$
- Q.18 An ideal gas is pressed at a constant temperature. Its internal energy  
 A. Decreases  
 B. First increases and then decreases  
 C. Increases  
 D. Remains the same
- Q.19 When 20 J of work was done on a gas, 40J of heat energy was released. If the initial internal energy of the gas was 70J, what is the final internal energy?  
 A. 50J  
 B. 90J  
 C. 60J  
 D. 110J

Topic-5

Q.20 A system goes from A to B via two processes I and II as shown in figure. If  $\Delta U_1$  and  $\Delta U_2$  are the changes in internal energies in the processes I and II respectively, then



- A.  $\Delta U_{II} > \Delta U_I$
- B.  $\Delta U_I = \Delta U_{II}$
- C.  $\Delta U_{II} < \Delta U_I$
- D. Relation between  $\Delta U_I$  and  $\Delta U_{II}$  cannot be determined

Q.21 By rubbing the objects together, their internal energy:

- A. Increases
- B. Remains constant
- C. Decreases
- D. Becomes zero

Q.22 The internal energy of an ideal gas depends upon only:

- A. Pressure
- B. Volume
- C. Temperature
- D. All of these


Q.23 If a system undergoes contraction of volume, then the work done by the system will be

- A. Zero
- B. Negative
- C. Negligible
- D. Positive

Q.24 The work done in the isochoric process is

- A. Constant
- B. Zero
- C. Variable
- D. Depends on situation


Q.25 A gas undergoes the cycle of pressure and volume changes  $W \rightarrow X \rightarrow Y \rightarrow Z \rightarrow W$  shown in the diagram.



What is the net work done by the gas?

- A. -600 J
- B. 0 J
- C. 200 J
- D. -200 J

Q.26 Calculate the heat absorbed by the system in going through the process as shown in figure.




- A. 31.4 J
- B.  $31.4 \times 10^6$  J
- C. 3.14 J
- D. None

Q.27 Work done by air when it expands from 50 litres to 150 litres at a constant pressure of 1 atmosphere is

- A.  $2 \times 10^4$  joules
- B.  $2 \times 10^5 \times 100$  joules
- C.  $2 \times 100$  joules
- D.  $2 \times 10^{-5} \times 100$  joules


Topic-5

Q.28 A system is taken from state A to B through three different paths 1, 2, 3. The work done is maximum in




- A. process 3
- B. process 2
- C. process 1
- D. equal in all processes

Q.29 What's the total work performed on the gas as it's transformed from state a to state c, along the path indicated?



- A. 1,500 J
- B. 4,500 J
- C. 3,000 J
- D. 5,000 J

Q.30 An ideal gas is taken around ABCA as shown in the above P-V diagram. The work done during a cycle is



- A. 2PV
- B. 1/2PV
- C. PV
- D. Zero


Q.31 The equation  $W = P(V_2 - V_1)$  represent work done by a gas in

- A. Free expansion
- B. An isothermal expansion
- C. An adiabatic expansion
- D. An expansion at constant pressure

Q.32 Find the change in internal energy of the system when a system absorbs 2 kilocalorie of heat and at the same time does 500 joules of work

- A. 8200 J
- B. 5600 J
- C. 7900 J
- D. 6400 J

Q.33 In the following indicator diagram, the net amount of work done will be




- A. Positive
- B. Negative
- C. Zero
- D. Infinity


Q.34 A system is described in terms of thermodynamics variables

- A. Pressure (P)
- B. Temperature (T)
- C. Volume (V)
- D. All of these

- Q.35 The concept of temperature is related to  
 A. Zeroth law of thermodynamics  
 B. Second law of thermodynamics  
 C. First law of thermodynamics  
 D. Third law of thermodynamics
- Q.36 Which of the following statement is correct for any thermodynamic system?  
 A. The internal energy changes in all processes  
 B. The change in entropy can never be zero  
 C. Internal energy and entropy are state functions  
 D. The work done in an adiabatic process is always zero
- Q.37 Thermodynamic is the study of relationship between  
 A. Heat & Surrounding  
 B. Heat & other form of energy  
 C. Heat & Liquid  
 D. Heat & chemical energy
- Q.38 In the figure curves AB and CD represent the relation between pressure P and volume V of an ideal gas. One of the curves represents an isothermal expansion and the other represents an adiabatic expansion. Which curve represents an adiabatic expansion?




- A. Curve AB  
 B. Curve CD  
 C. Both "A" and "B"  
 D. None of these
- Q.39 Heat added at constant volume of a gas is used to  
 A. To do external work  
 B. To increase its internal energy  
 C. Either "A" or "C"  
 D. Both "A" and "C"
- Q.40 When heat is given to a gas in an isobaric process then  
 A. The work is done by the gas  
 B. Internal energy of the gas increases  
 C. Both 'A' and 'B'  
 D. None of these
- Q.41 If the volume of a gas is decreased by 10% during isothermal process than its pressure will  
 A. Decrease by 10%  
 B. Decrease by 11.11%  
 C. Increase by 10%  
 D. Increase by 11.11%
- Q.42 During which process the volume of system remains constant  
 A. Isothermal  
 B. Isochoric  
 C. Isobaric  
 D. Adiabatic
- Q.43 In pressure-volume diagram given below, the isochoric, isothermal, and isobaric parts respectively, are



- A. AB, BC, CD  
 B. DC, CB, BA  
 C. CD, DA, AB  
 D. BA, AD, DC
- Q.44 A gas does 10J of external work in adiabatic process while expanding, then the change in internal energy is:  
 A. 10 J  
 B. -10 J  
 C. 20 J  
 D. 0 J

Q.45 A fixed mass of an ideal gas undergoes the changes represented by  $X \rightarrow Y \rightarrow Z \rightarrow X$  as shown below



	XY	YZ	ZX
A.	Isothermal compression	Adiabatic expansion	Pressure reduction at constant volume
B.	Adiabatic compression	Isothermal expansion	Pressure reduction at constant volume
C.	Isothermal compression	Adiabatic expansion	Compression at constant pressure
D.	Adiabatic compression	Isothermal expansion	Compression at constant pressure

- Q.46 During the adiabatic expansion of 2 moles of a gas, the internal energy of the gas is found to decrease by 2 joules, the work done during the process on the gas will be equal to  
 A. 2 J  
 B. 1 J  
 C. -2 J  
 D. -1 J
- Q.47 Starting with the same initial conditions, an ideal gas expands from volume  $V_1$  to  $V_2$  in three different ways. The work done by the gas is  $W_1$ , if the process is purely isothermal.  $W_2$  if purely isobaric and  $W_3$  if purely adiabatic then  
 A.  $W_2 > W_1 > W_3$   
 B.  $W_1 > W_2 > W_3$   
 C.  $W_2 > W_3 > W_1$   
 D.  $W_1 > W_3 > W_2$

PAST PAPER MCQs

- Q.48 What is the factor upon which change in internal energy of an ideal gas depends?  
 (MCAT 2013)  
 A. Change in volume  
 B. Change in temperature  
 C. Changed in volume and temperature  
 D. Path followed to change internal energy
- Q.49 The amount of heat required to raise the temperature of 1kg of substance through 1 K is called;  
 (ETEA 2014)  
 A. Heat capacity  
 B. 1 Joule  
 C. Specific heat  
 D. One calorie
- Q.50 If one mole of an ideal gas is heated at constant pressure, then the first law of thermodynamics can be written as:  
 (MDCAT 2018)  
 A.  $C_p \Delta T = C_v \Delta T + P \Delta V$   
 B.  $C_p \Delta T = C_v \Delta T + V \Delta P$   
 C.  $C_v \Delta T = C_p \Delta T + P \Delta T$   
 D.  $\Delta C_v T = \Delta C_v T + P \Delta V$
- Q.51 If  $C_v = 5/2 R$ ,  $C_p$  will be  
 (MDCAT 2018)  
 A.  $5/5 R$   
 B.  $5/2 R$   
 C.  $2/7 R$   
 D.  $7/2 R$
- Q.52 The amount of heat required to raise the temperature of 10 moles of water from 70K to 80K (molar heat capacity of water 75.24J) is:  
 (ETEA 2018)  
 A. 0.7524J  
 B. 7524J  
 C. 95.24J  
 D. 572.4J

# EXPLANATORY NOTES

Topic-5

- Q.53 The sum of all forms of molecular energies (kinetic and potential) of a substance is termed as? (MDCAT 2019)  
 A. Internal energy  
 B. Heat energy  
 C. Elastic energy  
 D. Absolute energy
- Q.54 In which process the entire of heat supplied to the gas is converted to the internal energy of the gas? (NMDCAT 2020)  
 A. Isochoric process  
 B. Isobaric process  
 C. Isothermal process  
 D. Adiabatic process
- Q.55 The internal energy of a system during an isothermal process:  
 A. Decreases  
 B. Increases  
 C. Become zero  
 D. Remain constant
- Q.56 In a certain process, 400J of heat energy is supplied to a system and at the same time 150J of work is done by the system. The increase in internal energy of system is (NUMS 2020)  
 A. 150J  
 B. 300J  
 C. 250J  
 D. 500J
- Q.57 The rapid escape of air from a burst tyre is an example of:  
 A. Isothermal  
 B. Adiabatic  
 C. Isothermal  
 D. Isochoric
- Q.58  $C_p - C_v$  value, find if 3moles of gas is gives (NMDCAT 2021)  
 A. 3R  
 B. 2R  
 C. 5R  
 D. 7R

## ANSWER KEY

### TOPIC-WISE MCQs & PAST PAPER MCQs

1	C	11	B	21	A	31	D	41	D	51	D
2	D	12	B	22	C	32	C	42	B	52	B
3	A	13	A	23	B	33	B	43	B	53	A
4	C	14	A	24	B	34	D	44	B	54	A
5	C	15	D	25	C	35	A	45	D	55	D
6	D	16	D	26	B	36	C	46	A	56	C
7	A	17	B	27	A	37	B	47	A	57	B
8	B	18	D	28	C	38	B	48	B	58	A
9	D	19	A	29	B	39	B	49	A		
10	C	20	B	30	A	40	C	50	A		

- Heat required to raise the temperature of a body through  $1^\circ\text{C}$  is called thermal capacity.
- Thermal capacity =  $mc$   
 Thermal capacity =  $40\text{g} \times 0.2\text{cal g}^{-1} \text{C}^{-1} = 8\text{ cal}^\circ\text{C}^{-1}$ .
- Heat used in melting  $m$  gram of ice = Heat lost by 80g water  
 $m\text{L} = 80 \times 1 \times 30$   
 $m = (80 \times 1 \times 30) / 80 = 30\text{g}$ .
- Mud is bad conductor of heat. It does not allow heat to come in from outside in summer and it does not allow heat to go out from a house in winter.
- Whatever amount of heat is taken in or given out by a substance of infinite heat capacity, there is no change in its temperature.
- $U = nC_v T$   
 $\Delta U/U \times 100 = \Delta T/T \times 100$   
 $\Delta U/U \times 100 = (1200 - 300) / 1200 \times 100 = 9/12 \times 100 = 75\%$ .
- For gas in cylinder A,  $Q = nC_p \Delta T_1$   
 For gas in cylinder B,  $Q = nC_v \Delta T_2$   
 $\Delta T_2 = C_p/C_v \Delta T_1 = 7/5 \times 20 = 28\text{K}$ .
- Thermodynamic equation  $\Delta Q = \Delta U + \Delta W$
- Examples of 1<sup>st</sup> law of thermodynamics.
- Fact.
- Statement of 1<sup>st</sup> law of thermodynamics.
- As the gas expands,  $W$  is negative, also  $\Delta U$  decreases and gas expand. There is no heat supplied to the system. The cooling occurs because of expansion.
- Ideal gas possess only kinetic energy.
- Ideal gas molecules only possess K.E.
- All options are related to internal energy.
- Internal energy is same and  $Q_p > Q_v$  so  $Q_v > Q_p$
- Internal energy  $\propto$  translational  $\langle \text{K.E} \rangle \propto T$
- $U \propto T$
- $\Delta U = Q - W$   
 $= -40 - (-20) = -20\text{J}$  ,  $\Delta U = U_2 - U_1$   
 $U_2 = \Delta U + U_1$  ,  $= -20 + 70 = 50\text{J}$
- As internal energy is a state function therefore change in internal energy does not depends upon the path followed i.e.  $\Delta U_1 = \Delta U_2$
- By rubbing objects, temperature increases hence internal energy increases.
- $U \propto T$
- $\Delta W = P\Delta V$ ; here  $\Delta V$  is negative so  $\Delta W$  will be negative
- $W = P\Delta V = P(0) = 0$
- $W = \Delta P\Delta V = 1 \times 10^5 \times 2 \times 10^{-3} = 2 \times 10^2 = 200\text{J}$   $W =$  positive because work done by yz path is more than xw path.

## Topic-5

26.  $Q = \Delta U + W$   
In cyclic process i.e.  $\Delta U = 0$   
 $Q = W = \text{Area of } P-V \text{ graph} = \pi r^2$   
 $= \pi (P_r)(V_r)$   
 $= 3.14(100 \times 10^3)(100) \Rightarrow 31.4 \times 10^6 J$
27.  $W = P\Delta V = 2 \text{ atm} \times (150 - 50) \text{ litre} = 2 \times 1.01 \times 10^5 \text{ Pa} \times 100 \times \frac{1}{1000} \text{ m}^3 \Rightarrow W = 2 \times 10^4 J$
28. Area of PV graph = W  
Area under process 1 > area under process 2 > area under process 3
29.  $W = p\Delta V = 3 \times 10^5 \times 15 \times 10^{-3} = 4500 J$
30. Work done = Area enclosed by triangle  $ABC = \frac{1}{2} AC \times BC = \frac{1}{2} \times (3V - V) \times (3P - P) = 2 PV$
31.  $W = P\Delta V = P(V_2 - V_1)$ . This equation is for work done by gas at constant pressure.
32.  $\Delta Q = 2 \text{ k cal} = 2 \times 10^3 \times 4.2 J = 8400 J$  and  $\Delta W = 500 J$ . Hence from  
 $\Delta Q = \Delta U + \Delta W$ ,  $\Delta W = \Delta Q - \Delta U = 8400 - 500 = 7900 J$
33. The cyclic process 1 is clockwise whereas process 2 is anticlockwise. Clockwise area represents positive work and anticlockwise area represents negative work. Since negative area (2) > positive area (1), hence net work done is negative.
34. Thermodynamics variables.
35. Zeroth law of thermodynamics describes temperature.
36. U and S are state functions.
37. In thermodynamics we deal with conversion of heat energy into other form of energy.
38. Adiabatic process.
39. Isochoric process.
40. Properties of isobaric process.
41.  $\therefore P \propto \frac{1}{V} \Rightarrow p_1 V_1 = p_2 V_2 \Rightarrow V_2 = V - 10\% = \frac{9V}{10}$   
 $p_1 = p_2 \frac{9}{10} \Rightarrow p_2 = \frac{10p_1}{9}$   
 $\%P = \frac{p_2 - p_1}{p_1} \times 100 = \frac{\frac{10p_1}{9} - p_1}{p_1} \times 100 = \frac{1}{9} \times 100 = 11\%$
42. Definition of isochoric process.
43. Graph of isochoric, isothermal and isobaric process.
44.  $\Delta Q = \Delta U + W \Rightarrow 0 = \Delta U + W \Rightarrow -\Delta U = W \Rightarrow \Delta U = -10 J$
45. During  $X \rightarrow Y$ ,  $V \downarrow, P \uparrow$       During  $Y \rightarrow Z$        $P \propto \frac{1}{V}$   
During  $Z \rightarrow X$        $P = \text{constant}$        $V \downarrow$

## Topic-5

46.  $Q = \Delta U + W$   
 $0 = -2 + W \Rightarrow W = +2J$
47.  $W_{\text{isobaric}} > W_{\text{isothermal}} > W_{\text{adiabatic}}$
48. Change in internal energy an ideal gas is depend on temperature according to given options.
49. Definition.
50.  $Q_p = \Delta U + W \Rightarrow Q_p \Delta T = C_v \Delta T + P\Delta V$
51.  $C_p - C_v = R \Rightarrow C_p = C_v + R$   
 $= \frac{5R}{2} + R = \frac{5R + 2R}{2} = \frac{7R}{2}$
52.  $Q = nC\Delta T = 10 \times 10 \times 75.24 = 7524 J$
53. Definition of internal energy.
54. In isochoric process  $\Delta V = 0$   
 $Q = \Delta U + P\Delta V \Rightarrow Q = \Delta U$
55.  $T \propto U$ ,  $T = \text{constant}$ , So  $U = \text{constant}$
56.  $Q = \Delta U + W$
57. The rapid escape of air from a burst tyre is an example of adiabatic.
58.  $C_p - C_v = nR$

# 6 TOPIC


# ELECTROSTATICS

## PRACTICE EXERCISE

- Q.1 Two charges  $1C$  and  $-4C$  exist in air. What is the direction of force?  
 A. Away from  $1C$   
 B. Away from  $-4C$   
 C. From  $1C$  to  $-4C$   
 D. From  $-4C$  to  $1C$
- Q.2 A point charge  $2nC$  is located at origin. What is the potential at  $(1,0,0)$ ?  
 A. 12  
 B. 14  
 C. 16  
 D. 18
- Q.3 A point charge  $0.4nC$  is located at  $(2, 3, 3)$ . Find the potential differences between  $(2, 3, 3)$  m and  $(-2, 3, 3)$  m due to the charge.  
 A. 2.5  
 B. 2.6  
 C. 2.7  
 D. 2.8
- Q.4 What is the dielectric constant of the medium if the capacitance of a parallel plate capacitor increases from  $40F$  to  $80F$  on introducing a dielectric medium between the plates?  
 A. 20  
 B. 0.5  
 C. 2  
 D. 5
- Q.5 What is the maximum energy that can be stored in a capacitor?  
 A. Depends on the maximum area of the capacitor  
 B. Depends on the maximum resistance offered  
 C. Depends on the maximum electric field  
 D. Depends on the maximum current passing through the capacitor
- Q.6 What is the voltage across the  $2F$  capacitor?



- A. 240V  
 B. 200V  
 C. 220V  
 D. 120V
- Q.7 What is the initial current while charging a capacitor?  
 A. High  
 B. Low  
 C. 0  
 D. Cannot be determined
- Q.8 Calculate the total charge of the system.



- A. 200C  
 B. 100C  
 C. 300C  
 D. 400C


## Topic-6

## Electrostatics

- Q.9 If the distance between two-point charges becomes double then the coulomb's force will be  
 A.  $\frac{F}{2}$   
 B.  $4F$   
 C.  $2F$   
 D.  $\frac{F}{4}$
- Q.10 The force between two-point charges placed in air is  $F$ . If air is replaced by a medium of relative permittivity  $\epsilon_r$ , the force is reduced to  
 A.  $\epsilon_r F$   
 B.  $\frac{\epsilon_r}{F}$   
 C.  $\frac{F}{\epsilon_r}$   
 D.  $\epsilon_r$
- Q.11 Two identical metal balls with charges  $+2Q$  and  $-Q$  are separated by some distance, and exert a force  $F$  on each other. They are joined by a conducting wire, which is then removed. The force between them will now be.  
 A.  $F$   
 B.  $F/2$   
 C.  $F/4$   
 D.  $F/8$
- Q.12 When  $10^{19}$  electrons are removed from a neutral metal plate, the electric charge on it is in (coulomb)  
 A.  $10^{-19}$   
 B.  $-1.6$   
 C.  $+1.6$   
 D.  $10^{-19}$
- Q.13 Two point charges  $+3\mu C$  and  $+8\mu C$  repel each other with a force of  $40N$ . If a charge of  $-5\mu C$  is added to each of them, then the force between them will become:  
 A.  $-10N$   
 B.  $+20N$   
 C.  $+10N$   
 D.  $-20N$
- Q.14 Two point charges  $+2C$  and  $+6C$  repel each other with a force of  $12N$ . If a charge of  $-2C$  is given to each of these charges, the force will now be  
 A. Zero  
 B.  $8N$  (repulsive)  
 C.  $8N$  (attractive)  
 D.  $16N$  (attractive)
- Q.15 Two electrons are removed from a conductor the charge on it is  
 A.  $1.6 \times 10^{-19}C$   
 B.  $-3.2 \times 10^{-19}C$   
 C.  $3.2 \times 10^{-19}C$   
 D. neutral
- Q.16 The force between two charges is  $120N$ . If the distance between the charges is doubled, the force will be  
 A.  $60N$   
 B.  $40N$   
 C.  $30N$   
 D.  $15N$
- Q.17 A positive charge is moved against an electric field. Its P.E  
 A. Increases  
 B. Remains same  
 C. Decreases  
 D. May increase or decrease depending upon magnitude of charge



Topic-6


- Q.18 An electric field can deflect  
 A. Neutrons  
 B. X-rays  
 C.  $\gamma$ -rays  
 D. None
- Q.19 Which of given is not the unit of electric intensity  
 A.  $NC^{-1}$   
 B.  $NV^{-1}$   
 C.  $Vm^{-1}$   
 D. None of these
- Q.20 The magnitude of electric intensity  $E$  is such that an electron placed in it would experience an electrical force equal to its weight.  $E$  is given by  
 A.  $mge$   
 B.  $\frac{mg}{e}$   
 C.  $\frac{e}{mg}$   
 D.  $\frac{e^2g}{m^2}$
- Q.21 The electric field intensity at a point 20 cm away from a charge of  $2 \times 10^{-3} C$  is  
 A.  $4.5 \times 10^5 N/C$   
 B.  $3.5 \times 10^5 N/C$   
 C.  $3.5 \times 10^6 N/C$   
 D.  $4.5 \times 10^5 N/C$
- Q.22 The weight of proton (mass =  $1.67 \times 10^{-27} kg$ ) on entering in a vertical electric field  $E$  is balanced by electric force. Then the electric field strength is  
 A.  $10^{-9} V m^{-1}$   
 B.  $10^{-7} V m^{-1}$   
 C.  $10^{-7} V m^{-1}$   
 D.  $10^{-8} V m^{-1}$
- Q.23 Figure shows the electric lines of force emerging from a charged body. If the electric field at A and B are  $E_A$  and  $E_B$  respectively and if the displacement between A and B is  $r$  then  
  
 A.  $E_A < E_B$   
 B.  $E_A = E_B$   
 C.  $E_A > E_B$   
 D. none of these
- Q.24 Two metal plates have potential difference of 300 V and are 0.01 m apart. A charged particle of mass  $1.96 \times 10^{-15} kg$  is held in equilibrium between the plates of the capacitor. Then the electric field is  
 A.  $3 \times 10^2 V m^{-1}$   
 B.  $3 \times 10^4 V m^{-1}$   
 C.  $3 V m^{-1}$   
 D.  $3 \times 10^{-4} V m^{-1}$
- Q.25 The electric field strength between a pair of parallel plates is  $E$ . The separation of plates is doubled and the potential difference between the plates is increased by a factor of four. What is the new electric field strength?  
 A.  $E$   
 B.  $4E$   
 C.  $2E$   
 D.  $8E$
- Q.26 The electric field intensity at a point situated 4 meters from a point charge is 200 N/C. If the distance is reduced to 2 meters, the field intensity will be  
 A. 400 N/C  
 B. 800 N/C  
 C. 600 N/C  
 D. 1200 N/C

Topic-6

- Q.27 The lines of force due to charged particles are  
 A. Always straight  
 B. Sometimes curved  
 C. Always curved  
 D. None of the above
- Q.28 Two thin infinite parallel plates have uniform charge densities  $+\sigma$  and  $-\sigma$ . The electric field in the space between them is  
 A.  $\sigma/2\epsilon_0$   
 B.  $\sigma$   
 C.  $\sigma/\epsilon_0$   
 D. zero
- Q.29 Gauss's law is applied to calculate the  
 A. Electric intensity due to different charge configuration  
 B. Electric intensity due to positive charges only  
 C. Electric intensity due to negative charges only  
 D. None of these
- Q.30 Electric intensity due to an infinite sheet of positive charge is given by  
 A.  $\vec{E} = \frac{\sigma}{2\epsilon_0} \hat{r}$   
 B.  $\vec{E} = \frac{2\sigma}{\epsilon_0} \hat{r}$   
 C.  $\vec{E} = \frac{\sigma}{\epsilon_0} \hat{r}$   
 D.  $\vec{E} = \frac{q}{\epsilon_0} \hat{r}$
- Q.31 The capacity of a parallel plate capacitor is  $5\mu F$ . When a glass plate is placed between the plates of the capacitor, its potential becomes 1/8 of the original value. The value of dielectric constant will be  
 A. 1.6  
 B. 8  
 C. 5  
 D. 40
- Q.32 Two charged spheres of radii 10 cm and 15 cm are connected by a thin wire. No current will flow, if they have:  
 A. The same charge on each  
 B. The same energy  
 C. The same potential  
 D. The same field on their surface
- Q.33 A and B are two points in an electric field. If the work done in carrying 4.0 coulomb of electric charge from A to B is 16.0 joule the potential difference between A and B is  
 A. Zero  
 B. 4 V  
 C. 2.0 V  
 D. 16 V
- Q.34 Potential due to charge  $q$  at distance 1m is 5V, at distance 3m will be  
 A.  $\frac{5}{3} V$   
 B.  $\frac{7}{3} V$   
 C.  $\frac{3}{5} V$   
 D.  $\frac{3}{7} V$
- Q.35 The electric potential at the surface of an atomic nucleus ( $Z = 50$ ) of radius  $9.0 \times 10^{-15} m$  is  
 A. 9 V  
 B.  $8 \times 10^6 V$   
 C.  $9 \times 10^5 V$   
 D. 80 V
- Q.36 A capacitor with air as the dielectric is charged to a potential of 100 volts. If the space between the plate is now filled with a dielectric of dielectric constant 10, the potential difference between the plates will be  
 A. 1000 V  
 B. 10 V  
 C. 100 V  
 D. 0 V

pic-6

Q.37 Figure shows three points A, B and C in a region of uniform electric field E. The line AB is perpendicular and BC is parallel to the field lines. Then which of the following holds good. Where  $V_A$ ,  $V_B$  and  $V_C$  represent the electric potential at points A, B and C respectively



- B.  $V_A = V_B < V_C$
- D.  $V_A > V_B = V_C$

- A.  $V_A = V_B = V_C$
- C.  $V_A = V_B > V_C$

Q.38 Value of potential at a point due to a point charge is

- A. Inversely proportional to square of the distance
- B. Inversely proportional to the distance
- C. Directly proportional to square of the distance
- D. Directly proportional to the distance

Q.39 Equal amount of charge is given to two sphere A and B of radii 2cm and 3cm respectively. The potential  $V_A$  and  $V_B$

- A.  $V_A = V_B$
- B.  $V_A > V_B$
- C.  $V_A < V_B$
- D. Depend upon number of material of sphere

Q.40 Two charge  $+q$  and  $-q$  are situated at a certain distance. At the point exactly midway between them

- A. Electric field and potential both are zero
- B. Electric field is not zero but potential is zero
- C. Electric field is zero but potential is not zero
- D. Electric field is not zero but potential is zero


Q.41 An- $\alpha$ -particle is accelerated through a potential difference of  $10^6$  V. Its K.E will be

- A. 1 MeV
- B. 4 MeV
- C. 2MeV
- D. 8MeV

Q.42 A capacitor C "has a charge Q". The actual charges on its plates are

- A. Q, -Q
- B. Q/2, -Q/2
- C. Q, Q
- D. Q, 0

Q.43 As in figure shown, if a capacitor C is charged by connecting it with resistance R, then energy is given by the battery will be



- A.  $\frac{1}{2} CV^2$
- B. Less than  $\frac{1}{2} CV^2$
- C. More than  $\frac{1}{2} CV^2$
- D. Zero

Q.44 A capacitor of capacitance  $2\mu\text{F}$  is connected with a battery of 12 volt, the charge stored is equal to:

- A.  $2.5 \times 10^{-5}$  C
- B.  $2.4 \times 10^{-6}$  C
- C.  $2.4 \times 10^{-5}$  C
- D.  $2.5 \times 10^5$  C

Q.45 If a  $2\mu\text{F}$  capacitor has a charge of  $20\mu\text{C}$ , the potential difference between the plates is:

- A. 10 V
- B. 20 V
- C. 40 V
- D. 50 V

Q.46 If a  $6\mu\text{F}$  capacitor is charged to 200 V, the charge in coulombs will be

- A.  $800\mu\text{C}$
- B.  $1200\mu\text{C}$
- C.  $900\mu\text{C}$
- D.  $1600\mu\text{C}$

Q.47 What is the area of the plates of a  $3\text{F}$  parallel plate capacitor, if the separation between the plates is 5mm?

- A.  $1.694 \times 10^9 \text{ m}^2$
- B.  $9.281 \times 10^9 \text{ m}^2$
- C.  $4.529 \times 10^9 \text{ m}^2$
- D.  $12.981 \times 10^9 \text{ m}^2$

Q.48 The capacity of a condenser in which a dielectric of dielectric constant 5 has been used, is C. If the dielectric is replaced by another with dielectric constant 20, the capacity will become

- A.  $\frac{C}{4}$
- B.  $\frac{C}{2}$
- C.  $4C$
- D.  $2C$

Q.49 The space between the plates of a capacitor is filled by a dielectric constant k. The capacitance of the capacitor

- A. Increases by a factor k
- B. Decreases by a factor k
- C. Increases by a factor  $k^2$
- D. Decreases by a factor  $k^2$

Q.50 If an insulating material called dielectric is introduced between the plates, the capacitance of capacitor is:

- A.  $\frac{A\epsilon_0\epsilon_r}{2d}$
- B.  $\frac{A\epsilon_0\epsilon_r}{d}$
- C.  $\frac{Ad}{\epsilon_0\epsilon_r}$
- D.  $\frac{2Ad\epsilon_0\epsilon_r}{d}$

Q.51 Capacitance with air is 10F, if a dielectric of  $\epsilon_r = 100$  is inserted then new capacitance

- A. 1000 F
- B.  $10\mu\text{F}$
- C. 10 F
- D. 100 F

Q.52 If the sheet of a bakelite is inserted between the plates of an air capacitor, the capacitance will

- A. Decrease
- B. REMAINS unchanged
- C. Increase
- D. become zero

Q.53 A dielectric material must be

- A. Resistor
- B. Good conductor
- C. Insulator
- D. Semi-conductor

Q.54 The capacitance of a capacitor is not affected by

- A. Distance between plates
- B. Thickness of plates
- C. Area of plates
- D. All of the above

Q.55 The quantity  $\frac{1}{2}\epsilon_0 E^2$  has the significance of:

- A. Energy/farad
- B. Energy/coulomb
- C. Energy/volume
- D. Energy

Q.56 The energy stored between the plates of a capacitor is not represented by

- A.  $U = \frac{CV^2}{2}$
- B.  $U = \frac{q^2}{2C}$
- C.  $U = 2qV$
- D.  $U = \frac{qV}{2}$

Topic-6

- Q.57 If the potential difference across the two plates of a parallel plate capacitor is doubled then its energy stored in it will be:  
 A. 2 times  
 B. 4 times  
 C. 16 times  
 D. Remains same
- Q.58 If a 10 F capacitor is to have an energy content of 20 J, it must be placed across a potential difference of  
 A. 4 volts  
 B. 2 volts  
 C. 9 volts  
 D. 1 volt
- Q.59 Energy stored in the electric field of a capacitor C when charged from a D.C source of voltage V is equal to joules  
 A.  $0.5CV^2$   
 B.  $CV^2$   
 C.  $0.5 CV^2$   
 D.  $0.5 CV$
- Q.60 When 4 volts e.m.f. is applied across a 1 farad capacitor, it will store energy of  
 A. 2 joules  
 B. 6 joules  
 C. 4 joules  
 D. 8 joules
- Q.61 A capacitor charging and discharging  
 A. Rapidly  
 B. Exponentially  
 C. Linearly  
 D. Logarithmically
- Q.62 If RC is small, then capacitor will be charged and discharged  
 A. Slowly  
 B. Quickly  
 C. With medium speed  
 D. Both 'A' and 'C'
- Q.63 Which of the following statement is true?  
 A. The current in the discharging capacitor grows linearly  
 B. The current in the discharging capacitor decays exponentially  
 C. The current in the discharging capacitor grows exponentially  
 D. The current in the discharging capacitor decreases constantly


## PAST PAPER MCQs

- Q.64 A particle carrying charge of  $2e$  falls through a potential difference of 3.0 V. Calculate the energy required by it: (MCAT 2009)  
 A.  $9.6 \times 10^{-19}$  J  
 B.  $9.1 \times 10^{-19}$  J  
 C.  $1.6 \times 10^{-19}$  J  
 D.  $6.0 \times 10^{-19}$  J
- Q.65 The work done in moving a unit positive charge from one point to another point against the electric field is a measure of: (MCAT 2009)  
 A. Capacitance  
 B. Potential difference between two points  
 C. Intensity of electric field  
 D. Resistance between two points
- Q.66 Electric intensity is a vector quantity and its direction is (MCAT 2010)  
 A. Perpendicular to the direction of field  
 B. Opposite to the direction of force  
 C. At a certain angle  
 D. Along the direction of force
- Q.67 The magnitude of an electric field between two separated plates can be calculated by the relation (MCAT 2010)  
 A.  $\Delta V = Ed$   
 B.  $\Delta V = E/d$   
 C.  $\Delta V = qd$   
 D.  $E = \Delta V$
- Q.68 The capacitor which charges and discharges quickly will have (ETEA 2011)  
 A. small value of RC  
 B. Large value of RC  
 C. Large value of time constant  
 D. None of these

Topic-6

Electrostatics


- Q.69 10 V potential difference is applied across the plates of 1 F capacitor. What is the energy stored in capacitor? (MCAT 2012)  
 A. 0.5 mJ  
 B. 0.05 mJ  
 C. 5 J  
 D. 50 J
- Q.70 What will be the effect on the capacitance of a capacitor if area of each plate is doubled with separation between the plates is halved? (MCAT 2012)  
 A. Capacitance remains same  
 B. Capacitance becomes double  
 C. Capacitance becomes four times  
 D. Capacitance reduces of half
- Q.71 What is the charge stored on a  $5\mu\text{F}$  capacitor charged to the potential difference of 12V? (MCAT 2013)  
 A.  $60\mu\text{C}$   
 B. 2.4C  
 C.  $2.4\mu\text{C}$   
 D. 60.C
- Q.72 The distance between the plates of a parallel plate capacitor is 2.0 mm and area of each plates is  $2.0 \text{ m}^2$ . A potential difference of  $1.0 \times 10^{-4}$  V is applied across the plates. Find the capacitance. (MDCAT 2014)  
 A.  $4 \times 10^4 \text{ F}$   
 B.  $3.54 \times 10^9 \text{ F}$   
 C.  $8.85 \times 10^{-9} \text{ F}$   
 D.  $9.0 \times 10^{-4} \text{ F}$
- Q.73 In the direction indicated by an electric field line: (ETEA 2014)  
 A. The potential must increase  
 B. The potential must decrease  
 C. The electric field strength must increase  
 D. The electric field strength must decrease
- Q.74 The potential difference between a pair of similar parallel conducting plates is known. What additional information is needed in order to find the electric field strength between the plates? (ETEA 2014)  
 A. Separation of the plates  
 B. Separation and decrease of the plates.  
 C. Permittivity of the medium separation of the plates.  
 D. Permittivity of the medium separation and area of plates.
- Q.75 The unit of the electric field is: (ETEA 2015)  
 A. N/C  
 B. V/m  
 C. J/C.m  
 D. All of the above
- Q.76 If the length, width and separation between the plates of a parallel plate capacitor is doubled then its capacitance becomes (MDCAT 2016)  
 A. Double  
 B. Half  
 C. Four time  
 D. Eight time
- Q.77 A charged capacitor stores 10C at 40V. Its stored energy is: (ETEA 2016)  
 A. 400J  
 B. 4J  
 C. 0.2 J  
 D. 200 J
- Q.78 A point charge at a distance 'x' from another point charge experiences a force of repulsion, which one of the following graphs shows. How the force is related to 'x': (MDCAT 2017)



6  
 The Coulomb force between two charges  $q_1 = 2C$  and  $q_2$  is  $2N$ , the distance between charges is  $3m$ . What is the charge of  $q_2$ ?  
 A.  $1 \times 10^9 C$   
 B.  $1 \times 10^{-9} C$   
 C.  $2 \times 10^9 C$   
 D.  $4 \times 10^9 C$

Q.80 The electric field strength at the position  $\vec{r} = (4\hat{i} + 3\hat{j})m$  caused by a point charge of  $5\mu C$  placed at origin is  
 A.  $1240\hat{i} + 1280\hat{j} N/C$   
 B.  $1440\hat{i} + 1080\hat{j} V/m$   
 C.  $1440\hat{i} + 1080\hat{j} N/m$   
 D.  $1240\hat{i} + 1080\hat{j} N/C$

Q.81 If a charge particle is placed one by one at point A, B and C then at which point it will experience a large force:



A. At point "A"  
 B. At point "C"  
 C. At point "B"  
 D. Same at all point

Q.82 The potential of the two plates of a capacitor are  $+10V$  and  $-10V$ . The charge on one of the plates is  $40C$ . The capacitance of the capacitor is:  
 A.  $2F$   
 B.  $4F$   
 C.  $0.5F$   
 D.  $0.25F$

Q.83 The ratio of the electric force between two protons to that between two electrons is of the order of:  
 A.  $10^{42}$   
 B.  $10^{39}$   
 C.  $10^{36}$   
 D.  $1$

Q.84 Coulomb's law is given by the formula  $F = k q_1 q_2 / r^2$ . The magnitude of  $k$  having the unit of  $Nm^2C^{-2}$  for free space is equal to  
 A.  $9 \times 10^7$   
 B.  $6 \times 10^7$   
 C.  $10 \times 10^9$   
 D.  $9 \times 10^9$

Q.85 Force experienced per unit positive test charge at a point in an electric field is the definition of:  
 A. Electric potential energy  
 B. Electric field strength  
 C. Electric potential  
 D. Electric field

Q.86 A torch is rated  $2.2V$ ,  $0.25A$ . Calculate the charge passing through the bulb in one second and energy transferred by the passage of each coulomb of charge.  
 A.  $2.5C$  and  $0.55J$   
 B.  $0.25C$  and  $2.2J$   
 C.  $0.25C$  and  $2.2V$   
 D.  $0.25C$  and  $0.55J$

Q.87 Electric potential due to  $2\mu C$  charge at distance of one meter is equal to  
 A.  $18 \times 10^4 \text{ volt}$   
 B.  $1.8 \times 10^6 \text{ volt}$   
 C.  $1.8 \times 10^9 \text{ volt}$   
 D.  $1.8 \times 10^4 \text{ volt}$

Q.88 Two determine the resistance of a voltmeter by discharging a capacitor through it, the instantaneous voltage is then given by the relation:  
 A.  $V_0 e^{-t/RC}$   
 B.  $V_0 e^{t/RC}$   
 C.  $V_0/2$   
 D.  $V_0/\sqrt{2}$

Q.89 There are two charges  $+3\mu C$  and  $+8\mu C$  the ratio of the force acting on them will be:  
 A.  $3:1$   
 B.  $1:1$   
 C.  $11:8$   
 D.  $3:8$

Q.90 What is the magnitude of a point charge which produces an electric field of  $2NC^{-1}$  at a distance of  $60cm$ ?  
 A.  $8 \times 10^{-11}C$   
 B.  $2 \times 10^{-12}C$   
 C.  $3 \times 10^{-11}C$   
 D.  $6 \times 10^{-10}C$

Q.91 The force between two charged bodies is "F". if one of the charge is doubled and the distance between them is halved, the force acting on each charged body is:  
 A.  $2F$   
 B.  $4F$   
 C.  $8F$   
 D.  $16F$

Q.92 Electric field strength of a point charge is  $E$  and electric potential is  $V$  at a distance  $r$  from the point charge. What is the electric potential at a point for the same point charge where electric field strength is  $E/4$ ?  
 A.  $V/4$   
 B.  $4V$   
 C.  $V/2$   
 D.  $2V$

Q.93 A particle carrying a charge of  $5e$  falls through a potential difference of  $25V$ . What would be energy acquired by the particle in 'J'.  
 A.  $125 \times 10^{-19}J$   
 B.  $1.6 \times 10^{-19}J$   
 C.  $125 \times 1.6 \times 10^{-19}J$   
 D.  $125J$

Q.94 Electric field strength at a point between oppositely charge plates is  $E$ . If the distance between plates is reduced to half, what will be the new value of electric intensity?  
 A.  $4E$   
 B.  $E/2$   
 C.  $E/4$   
 D.  $2E$

Q.95 If the potential at a point which is  $1m$  from a charge is  $1\text{volt}$ , then the potential at a point which is  $2m$  from the same charge will be:  
 A.  $2v$   
 B.  $1v$   
 C.  $0.5v$   
 D.  $3v$

Q.96 The values of electric intensity will \_\_\_\_\_ due to the presence of dielectric medium:  
 A. Increase  
 B. Increase exponentially  
 C. Decrease  
 D. Remains same

Q.97 In capacitors, energy is stored in the form of:  
 A. Gravitational energy  
 B. Kinetic energy  
 C. Electric intensity  
 D. Magnetic energy

Q.98 Ohm time's farad is equivalent to:  
 A. Time  
 B. Charge  
 C. Distance  
 D. Capacitor

# EXPLANATORY NOTES

Q.99 By increasing area of the plates and decreasing distance between them the capacitance of capacitor:  
 A. Increases  
 B. Decreases  
 C. Remains unchanged  
 D. Depending upon temperature  
 (NUMS 2020)

Q.100 If we double the separation between two charges, then coulomb's force will become?  
 A. Doubled  
 B. Half  
 C. 4-times  
 D.  $1/4^{\text{th}}$   
 (NMDCAT 2021)

Q.101 Electric force and electric field intensity for a charge are  
 A. Parallel to each other  
 B. Opposite to each other  
 C. Perpendicular to each other  
 D. At any orientation  
 (NMDCAT 2021)

Q.102 A charge Q placed at the centre of two charges +q and +q. the system is in equilibrium when net force experienced by Q.  
 A. Zero  
 B. F  
 C. At equilibrium, net force will be zero  
 D. F/2  
 (NMDCAT 2021)

Q.103 In series combination of capacitors  
 A.  $C_1 + C_2 + C_3$   
 B.  $\frac{C_1 C_2}{C_1 + C_2}$   
 C.  $\frac{C_1 + C_2}{C_1 C_2}$   
 D.  $\frac{1}{C_1 + C_2 + C_3}$   
 (NMDCAT 2021)

Q.104 Work done per unit coulomb is equal to  
 A. Electric field  
 B. Magnetic field  
 C. Electric potential  
 D. None  
 (NMDCAT 2021)

Q.105 Work on charge is equipotential surface is  
 A. +ve  
 B. -ve  
 C. Zero  
 D. None  
 (NMDCAT 2021)

Q.106 Capacitance of spherical capacitor is  
 A.  $4\pi Eab$   
 B.  $4\pi E(b-a)$   
 C.  $4\pi E(b-a)$   
 D. None  
 (NMDCAT 2021)

Q.107 For n numbers of Capacitors, each of the capacitance Is "C" what will be the ratio between maximum and minimum capacitor  
 A. n  
 B.  $n^2$   
 C.  $n^3$   
 D.  $n^4$   
 (NMDCAT 2021)

## ANSWER KEY

### TOPIC-WISE MCQs & PAST PAPER MCQs

1	C	11	D	21	A	31	B	41	C	51	A	61	B	71	A	81	D	91	C	101	A
2	D	12	C	22	B	32	C	42	A	52	C	62	B	72	C	82	A	92	C	102	C
3	C	13	A	23	C	33	B	43	C	53	C	63	C	73	B	83	D	93	C	103	B
4	C	14	A	24	B	34	A	44	C	54	B	64	A	74	A	84	D	94	D	104	C
5	C	15	C	25	C	35	B	45	A	55	C	65	B	75	D	85	B	95	C	105	C
6	D	16	C	26	B	36	B	46	B	56	C	66	D	76	A	86	D	96	C	106	D
7	A	17	A	27	C	37	C	47	A	57	B	67	A	77	D	87	D	97	C	107	B
8	C	18	D	28	C	38	B	48	C	58	B	68	A	78	B	88	A	98	A		
9	D	19	B	29	A	39	B	49	A	59	A	69	D	79	B	89	B	99	A		
10	C	20	B	30	A	40	D	50	B	60	D	70	C	80	B	90	A	100	D		

- Since the charges are unlike, the force will be attractive. Thus, the force directs from 1C to -4C.
- $V = Q/(4\pi\epsilon_0 r)$ , where  $r = 1\text{m}$   
 $V = (2 \times 10^{-9})/(4\pi\epsilon_0 \times 1) = 18 \text{ volts.}$
- $V_{AB} = (Q/4\pi\epsilon_0)(1/r_A) + (1/r_B)$ , where  $r_A$  and  $r_B$  are position vectors  $r_A = 1\text{m}$  and  $r_B = 4\text{m}$ . Thus  $V_{AB} = 2.7 \text{ volts.}$
- Capacitance without dielectric = 40 F.  
 Capacitance with dielectric = 80 F.  
 $k = 80/40$   
 $k = 2.$
- The maximum energy that can be desirably stored in a capacitor is depends on the maximum electric field that the dielectric can withstand without breaking down. Therefore, capacitors of the same type have about the same maximum energy density, i.e. joules of energy per cubic meter.
- Capacitors are in series.  
 $1/C = 1/2 + 1/4 + 1/6$ , therefore,  $C = (12/11) \text{ F.}$   
 $Q = C \times V = 220 \times (12/11) = 240\text{C.}$   
 $V \text{ across } 2\text{F capacitor} = Q/C = 240/2 = 120\text{V.}$
- The initial current of a capacitor is very high because the voltage source will transport charges from one plate of the capacitor to the other plate.
- The equivalent capacitance when capacitors are connected in parallel is the sum of all the capacitors = 1 + 2 = 3F.  $Q = CV = 3 \times 100 = 300\text{V.}$
- If  $r = 2r$

$$F = \frac{kq^2}{r^2} \Rightarrow F' = \frac{F}{4}$$

$$10. F = \frac{kq_1 q_2}{r^2}$$

$$F' = \frac{1}{4\pi\epsilon_0 \epsilon_r} \times \frac{q_1 q_2}{r^2} \Rightarrow F' = \frac{F}{\epsilon_r}$$

11. When two balls are connected by a conducting wire the net charge is  $(2Q - Q)$  i.e Q and Q is divided equally between two balls.

$$F \propto 2Q^2 \text{ --- (i)}$$

$$F' \propto \left(\frac{Q}{2}\right)^2 \text{ --- (ii)}$$

Dividing equation (ii) by equation (i)

$$\frac{F'}{F} = \frac{Q^2/4}{2Q^2} = \frac{1}{8}$$

$$F' = \frac{1}{8} F$$

Topic-6

12. As electric charge on electron is  $Q = 1.6 \times 10^{-19} \text{ C}$   
 If  $10^{19}$  electrons removed from a neutral plate, then electric charge is  $+1.6 \text{ C}$ .  
 In second case, charges will be  $-2 \mu\text{C}$  and  $+3 \mu\text{C}$ . Since  $F \propto Q_1 Q_2$ , i.e.  

$$\frac{F}{F'} = \frac{Q_1 Q_2}{Q'_1 Q'_2} = \frac{40}{-2 \times 3} = -\frac{3 \times 8}{2 \times 3} = -4 \Rightarrow F' = 10 \text{ N (Attractive)}$$
  
 14. Because  $+2 \text{ C}$  charge will become neutral with  $-2 \text{ C}$  charge. So  $F = 0$   
 15. If two electrons are removed from a conductor the charge on it is equal to 2 protons  
 $q = 2e = 2 \times 1.6 \times 10^{-19}$ ,  $C = 3.2 \times 10^{-19} \text{ C}$

16. 
$$F = \frac{kq_1 q_2}{r^2} \Rightarrow F \propto \frac{1}{r^2}$$
  
 $r' = 2r \Rightarrow F' = \frac{F}{4} = \frac{120}{4} = 30 \text{ N}$

17. When positive charge is moved against electric field its P.E increases. When negative charge is moved along electric field its P.E increases.  
 18. Neutron, x-rays and  $\gamma$ -rays are neutral.

19.  $E = \frac{F}{q} \Rightarrow \text{NC}^{-1}$   
 $E = \frac{\Delta V}{\Delta r} \Rightarrow \text{Vm}^{-1}$   
 $\text{NC}^{-1}, \text{Vm}^{-1}$  both are units of electric field.

20.  $F_e = F_g$   
 $eE = mg$   
 $E = \frac{mg}{e}$

21.  $E = \frac{kq}{r^2} \Rightarrow \frac{9 \times 10^9 \times 2 \times 10^{-3}}{(20 \times 10^{-2})^2}$   
 $E = 4.5 \times 10^6 \text{ N/C}$

22.  $E = \frac{F}{q} = \frac{mg}{q} = \frac{1.67 \times 10^{-27} \times 10}{1.6 \times 10^{-19}} = 10^{-7} \text{ Vm}^{-1}$

23. Field will be stronger where field lines are closer to each other i.e.  $E_A > E_B$

24. As  $E = \frac{\Delta V}{\Delta r}$   
 $E = \frac{300}{0.01} = 30000 = 3 \times 10^4 \text{ Vm}^{-1}$

Topic-6

25.  $E = \frac{\Delta V}{\Delta r} = \frac{4\Delta V}{2\Delta r} = 2$

26.  $E = \frac{kq}{r^2} \Rightarrow r' = \frac{r}{2}$   
 $E' = 4E = 4 \times 200 = 800 \text{ N/C}$

27. Properties of electric field lines  
 28. According to 3<sup>rd</sup> application of Gauss's law, the electric field between two equal and opposite charged plates is  $E = \frac{\sigma}{\epsilon_0}$

29. Gauss's law is used to determine electric field intensity due to different charge configuration.

30.  $E = \frac{\sigma}{2\epsilon_0} \hat{r}$

31.  $V_{med} = \frac{V}{\epsilon_r}$

$\epsilon_r = \frac{V}{V_{med}} = \frac{V}{V/8} = 8$

32. Because current flows from higher potential to lower potential.

33.  $\Delta V = \frac{W}{q} = \frac{16}{4} = 4 \text{ V}$

34. As we know,

$V = \frac{kq}{r} \Rightarrow \frac{V_1}{V_2} = \frac{q_1/r_1}{q_2/r_2}$

$V_2 = \frac{q_2}{r_2} \times \frac{r_1}{q_1} \times V_1 \Rightarrow V_2 = \frac{q}{3} \times \frac{1}{q} \times 5 \Rightarrow V_2 = \frac{5}{3} \text{ volt}$

35.  $V = \frac{1}{4\pi\epsilon_0} \frac{Ze}{r}$

$V = \frac{9 \times 10^9 \times 50 \times 1.6 \times 10^{-19}}{9 \times 10^{-15}} \Rightarrow V = 8.0 \times 10^6 \text{ V}$

36.  $C_{med} = \epsilon_r \times C_{vac}$

$\frac{\phi}{V_{med}} = \epsilon_r \times \frac{\phi}{V_{vac}}$


$V_{med} = \frac{V_{vac}}{\epsilon_r} = \frac{100}{10} = 10 \text{ V}$

37. In the direction of electric field potential decreases.

38.  $V = \frac{1}{4\pi\epsilon_0} \frac{Q}{r} \Rightarrow V \propto \frac{1}{r}$

39.  $I' = \frac{kq}{r} \Rightarrow I' \propto \frac{1}{r}$

40. At O,  $E \neq 0, V = 0$



41.  $K.E = q\Delta V$

where  $q_a = 2e$

$K.E = (2e)(10^6 V) = 2 \times 10^6 eV = 2 MeV$

42. Both plates has opposite charge having magnitude "Q".

43. Energy stored in a capacitor and some loss of energy in the form of heat in resistance.

44.  $Q = CV$

$Q = 2 \times 10^{-9} \times 12 \Rightarrow Q = 2.4 \times 10^{-5} C$

45.  $Q = CV$

$V = \frac{Q}{C} = \frac{20 \times 10^{-6}}{2 \times 10^{-6}} = 10 \text{ volt}$

46.  $Q = CV = (6 \times 10^{-9}) \times 200 = 1200 \mu C$

47. We have  $C = \frac{\epsilon_0 A}{d} \Rightarrow A = \frac{Cd}{\epsilon_0} = \frac{3 \times 5 \times 10^{-3}}{8.85 \times 10^{-12}} = 1.69 \times 10^9 m^2$

48.  $C = \frac{K\epsilon_0 A}{d}; \frac{C_1}{C_2} = \frac{K_1}{K_2} \Rightarrow \frac{C}{20} = \frac{5}{2} \Rightarrow C_2 = 4C$

49.  $C_{med} = \frac{A\epsilon_0 k}{d} \Rightarrow C_{med} \propto k$

50.  $C = \frac{A\epsilon_0 \epsilon_r}{d}$

51. As,  $C_{med} = \epsilon_r \times C_{vac} \Rightarrow C = 100 \times 10 \Rightarrow C = 1000 F$

52. When dielectric is inserted between plates then capacitance will increase

53. information

54.  $C_{vac} = \frac{A\epsilon_0}{d}$ , Thickness of plates has no effect on the capacitance of a capacitor.

55. Energy density =  $\frac{\text{energy}}{\text{volume}} = \frac{1}{2} \epsilon_0 E^2$

56. As we know,

$U_m = \frac{1}{2} CV^2 = \frac{Q^2}{2C} = \frac{qV}{2}$  but  $U_m \neq 2qV$

57.  $E = \frac{1}{2} CV^2 \Rightarrow E \propto V^2$

$V' = 2V \Rightarrow E' = 4E$

58. As we know,

$E = \frac{1}{2} CV^2 \Rightarrow V = \sqrt{\frac{2E}{C}}$

$V = \sqrt{\frac{2 \times 20}{10}} \Rightarrow V = \sqrt{4} = 2 \text{ volt}$

59. Energy =  $\frac{1}{2} CV^2$

60.  $U = \frac{1}{2} CV^2 = \frac{1}{2} \times 1 \times (4)^2 = 8J$

61. A capacitor charging and discharging exponentially.

62.  $RC =$  time constant

If  $RC$  is small then time constant will also small then capacitor will be charged and discharged quickly.

63. In discharging of capacitor, current increases exponentially.

64.  $\Delta K.E = q\Delta V$

$= (2e)(3V) = 6eV$

$= 6 \times 1.6 \times 10^{-19} = 9.6 \times 10^{-19} J$

65. Definition of potential difference

66.  $\vec{E} = \frac{\vec{F}}{q_0} \therefore \hat{E} = \hat{F}$

67.  $\Delta V = Ed$

68. Smaller value of time constant,  $RC$  leads to more discharge.

69. Energy stored =  $\frac{1}{2} CV^2 = \frac{1}{2} (1)(10)^2 = 50J$

70.  $C = \frac{A\epsilon_0}{d} \Rightarrow C' = \frac{(2A)\epsilon_0}{\frac{1}{2}d} = 4C$

71.  $Q = CV = (5 \times 10^{-6})(12) = 60 \mu C$

72.  $C = \frac{A\epsilon_0}{d} = \frac{2 \times 8.85 \times 10^{-12}}{2 \times 10^{-3}} = 8.85 \times 10^{-9} F$

73.  $E = \frac{-\Delta V}{\Delta r}$

74.  $E = \frac{V}{d}$

75.  $E = \frac{\Delta V}{\Delta r} = \frac{V}{m} = \frac{J}{Cm} \Rightarrow E = \frac{F}{q} = \frac{N}{C}$

76. Length =  $L' = 2L$ , Width =  $W' = 2W \Rightarrow A' = 2L \times 2W = 4A \Rightarrow C' = \frac{(4A)\epsilon_0}{2d} = 2C$

77.  $U = \frac{1}{2} QV = \frac{1}{2} \times 10 \times 40 = 200J$

## Topic-6

78.  $F = \frac{kq_1q_2}{r^2} \Rightarrow F \propto \frac{1}{r^2}$
79.  $F = \frac{kq_1q_2}{r^2} \Rightarrow 2 = \frac{(9 \times 10^9)(2)q_2}{3^2} \Rightarrow q_2 = 1 \times 10^{-9} \text{ C}$
80.  $\vec{r} = (4\hat{i} + 3\hat{j}) \text{ m} \Rightarrow |\vec{r}| = \sqrt{4^2 + 3^2} = 5 \text{ m}$   
 $\vec{E} = \frac{kq}{r^2} \vec{r} = \frac{kq}{r^2} \frac{\vec{r}}{r} = \frac{(9 \times 10^9)(5 \times 10^{-8})(4\hat{i} + 3\hat{j})}{(5)^2 \times 5} = (1440\hat{i} + 1080\hat{j}) \text{ V/m}$
81. In uniform electric field then at point A, B, C a charge particle will experience same force.
82.  $C = \frac{Q}{\Delta V}$   $\Delta V = V_2 - (-V_1)$   $\Delta V = 20 \text{ V}$   $C = \frac{40 \text{ C}}{20 \text{ V}} = 2 \text{ F}$
83.  $c = P$   $Fc = \frac{kc^2}{R^2}$   $Fp = \frac{kc^2}{R^2}$   $\frac{Fc}{Fp} = \frac{kc^2}{R^2} \times \frac{R^2}{kc^2} = 1$
84.  $k = 9 \times 10^9 \text{ Nm}^2\text{C}^{-2}$
85.  $\vec{E} = \frac{\vec{F}}{q_0}$
86.  $P = VI \Rightarrow \frac{E}{l} = VI \Rightarrow E = VI \times l = 2.2 \times 0.25 \times 1 = 0.55 \text{ V}$   
 $V = \frac{E}{q} \Rightarrow q = \frac{E}{V} = \frac{0.55}{2.2} = 0.25 \text{ C}$
87.  $V = \frac{kq}{r} = \frac{(9 \times 10^9)(2 \times 10^{-6})}{1^2} = 1.8 \times 10^4 \text{ V}$
88. For discharging  $q = q_0 e^{-t/RC}$  (Dividing by C) so:  $q/C = q_0/C(e^{-t/RC})$ ,  $V = V_0 e^{-t/RC}$
89. Electrostatic force is mutual force
90.  $E = Kq/r^2$
91.  $F = \frac{kq_1q_2}{r^2}$
92.  $E \propto \frac{1}{r^2}$  for  $r' = 2r \Rightarrow E' = \frac{E}{4}$   
 $V \propto \frac{1}{r} \Rightarrow V' \propto \frac{1}{2r} \Rightarrow V' = \frac{V}{2}$
93. Energy  $= qV = 5e \times 25 = 125 \text{ eV} = 125 \times 1.6 \times 10^{-19} \text{ J}$
94.  $E = -\frac{\Delta V}{\Delta r} \Rightarrow E \propto \frac{1}{\Delta r}$

95.  $V \propto \frac{1}{r} \rightarrow \frac{V_1}{V_2} = \frac{r_2}{r_1}$   
 $\frac{1}{V_2} = 2 \rightarrow V_2 = \frac{1}{2} = 0.5 \text{ Volt}$
96.  $E = \frac{q}{4\pi\epsilon_0\epsilon_r r^2} \rightarrow E \propto \frac{1}{\epsilon_r}$
97. In capacitor electrical energy is stored in the form of electric intensity  
 $E_d = \frac{1}{2} \epsilon_0 E^2$
98.  $RC = t \Rightarrow (\text{ohm})(\text{farad}) = \text{time}$
99.  $C = \frac{A\epsilon_0}{d}$
100.  $F \propto \frac{1}{r^2}$
101.  $\hat{E} = \hat{F}$
102. At equilibrium  $\Sigma F = 0$
103. For two capacitor of series  $C_s = \frac{\text{Product}}{\text{Sum}} = \frac{C_1 C_2}{C_1 + C_2}$
104.  $V = \frac{W}{q}$
105. On equipotential surface  $\Delta V = 0$   
 $W = q\Delta V = 0$
106.  $C = 4\pi\epsilon_0 \left( \frac{ba}{b-a} \right)$
107.  $\frac{C_{\max}}{C_{\min}} = \frac{nc}{\frac{c}{n}} = n^2$



- Q.1 Give the number of electrons passing through a wire per minute. The current flowing through it is 500mA.  
 A.  $1.875 \times 10^{20}$   
 C.  $1.875 \times 10^{-20}$   
 B.  $6.875 \times 10^{20}$   
 D.  $6.875 \times 10^{-20}$
- Q.2 Kabir bought 5 new light bulbs of 50 W each in addition to the 7 bulbs he already had in his house which were also 50 W each. Calculate his electricity bill, if he keeps the new bulbs on for 5 hours and the older bulbs on only for 3 hours, and the cost of one unit of electricity is Rs. 60.  
 A. Rs.136  
 C. Rs.138  
 B. Rs.137  
 D. Rs.139
- Q.3 Which one of the following is the practical unit of power?  
 A. Watt (W)  
 C. Horse power (hp)  
 B. Kilowatt hour (kWh)  
 D. Kilojoule (kJ)
- Q.4 An engine uses 30 A of current. The resistance offered is 15 ohms. Calculate the power consumed by the engine in horse power.  
 A. 18 hp  
 C. 17 hp  
 B. 19 hp  
 D. 13500 hp
- Q.5 A current of 3 A passes through an electric circuit for 5 minutes and does a work of 900J. What is the emf of the source?  
 A. 3V  
 C. 5V  
 B. 1V  
 D. 10V
- Q.6 A cooler of 1500 W, 200 volt and a fan of 500 W, 200 volts are to be used from a household supply. The rating of fuse to be used is  
 A. 2.5 A  
 C. 7.5 A  
 B. 5.0 A  
 D. 10 A
- Q.7 An electric current is passed through a circuit containing two wires of the same material, connected in parallel. If the lengths and radii of the wires are in the ratio of 4/3 and 2/3, then the ratio of the currents passing through the wire will be  
 A. 3  
 C. 1/3  
 B. 8/9  
 D. 2
- Q.8 Graph between V and I for non-ohmic devices  
 A. Straight line  
 C. Usually not straight-line  
 B. Always not straight line  
 D. Sometime straight line
- Q.9 The current in a resistor is 8.0 mA. What charge flows through the resistor in 0.020 s?  
 A. 0.16 mC  
 C. 1.6 mC  
 B. 4.0 mC  
 D. 0.40 mC
- Q.10 Which of the following statements is not true?  
 A. Conductance is the reciprocal of resistance and is measured in siemens  
 B. Semiconductors diode are ohmic devices  
 C. Ohm's law is not applicable at very low and very high temperatures  
 D. Ohm's law is not applicable to electron tubes, discharge tubes


- Q.11 Ohm's law establishes a relation between  
 A. Current and voltage  
 C. Charge and voltage  
 B. Resistance and voltage  
 D. Current and resistance
- Q.12 For an ohmic conductor, doubling the voltage without changing the resistance will cause the current to  
 A. Decrease by a factor of 4  
 C. Remain unchanged  
 B. Decrease by a factor of 2  
 D. Increase by a factor of 2
- Q.13 Which equation is used to define resistance?  
 A. Energy = (current)<sup>2</sup> × resistance × time  
 B. Power = (current)<sup>2</sup> × resistance  
 C. Potential difference = current × resistance  
 D. Resistivity = resistance × area ÷ length
- Q.14 The charge of an electron is  $1.6 \times 10^{-19}$  (c) How many electrons strike the screen of a cathode ray tube each second when the beam current is 16 mA  
 A.  $10^{19}$   
 C.  $10^{17}$   
 B.  $10^{-19}$   
 D.  $10^{-17}$
- Q.15 The following four wires are made of the same material and are at the same temperature. Which one of them has highest electrical resistance  
 A. length = 100 cm, diameter = 1 mm  
 C. length = 50 cm, diameter = 0.5 mm  
 B. length = 200 cm, diameter = 2 mm  
 D. length = 300 cm, diameter = 3 mm
- Q.16 The resistance of a wire of uniform diameter d and length l is R. The resistance of another wire of the same material but diameter 2d and length 4l will be  
 A. 2R  
 C. R  
 B.  $\frac{R}{2}$   
 D.  $\frac{R}{4}$
- Q.17 When the length and area of cross-section both are doubled, then its resistance  
 A. Will become half  
 C. Will be doubled  
 B. Will remain the same  
 D. Will become four times
- Q.18 The example for non-ohmic resistance is  
 A. Copper wire  
 C. Carbon resistance  
 B. Diode  
 D. Tungston wire
- Q.19 A wire 100cm long and 2.0 mm diameter has a resistance of 0.7 ohm, the electrical resistivity of the material is  
 A.  $4.4 \times 10^{-6}$  ohm m  
 C.  $2.2 \times 10^{-6}$  ohm m  
 B.  $1.1 \times 10^{-6}$  ohm m  
 D.  $0.22 \times 10^{-6}$  ohm m
- Q.20 Which of the following have the same temperature coefficient of resistivity?  
 A. iron and silver  
 C. iron and platinum  
 B. platinum and silver  
 D. silver and gold
- Q.21 When a wire is stretched double of its length, then its resistance will be  
 A. 16 R  
 C. 2 R  
 B. 4 R  
 D. 8 R
- Q.22 A wire of resistance R is cut into two equal parts, its resistance becomes R/2, what happens to resistivity:  
 A. Double  
 C. Half  
 B. Same  
 D. One fourth

Topic-7

- Q.23 A wire of resistance  $R$  is stretched four times its length uniformly. Its new resistance will be  
 A.  $16R$   
 B.  $4R$   
 C.  $\frac{R}{4}$   
 D.  $\frac{R}{16}$
- Q.24 The resistance of a wire is  $1\ \Omega$ . Which of the following is new resistance if length of wire is doubled?  
 A.  $2\ \Omega$   
 B.  $4\ \Omega$   
 C.  $\frac{1}{2}\ \Omega$   
 D.  $\frac{1}{4}\ \Omega$
- Q.25 When Cu and Ge are cooled to  $-150\ ^\circ\text{C}$  Then resistance of Cu \_\_\_\_\_ and that of Ge \_\_\_\_\_  
 A. Increases, increases  
 B. Decreases, decreases  
 C. Increases, decreases  
 D. Decreases, increases
- Q.26 If a wire conductor of  $0.2\ \text{ohm}$  resistance is doubled in length, its resistance becomes  
 A.  $0.4\ \text{ohm}$   
 B.  $0.8\ \text{ohm}$   
 C.  $0.6\ \text{ohm}$   
 D.  $1.0\ \text{ohm}$
- Q.27 If  $r = 0$ ,  $R = \infty$ , ( $V = \epsilon - Ir$ ) then  
 A.  $V > \epsilon$   
 B.  $V = \epsilon$   
 C.  $V < \epsilon$   
 D.  $V = 0$
- Q.28 An  $8\ \Omega$  resistance connected to a battery with internal resistance draws  $1.6\ \text{A}$  and if a  $30\ \Omega$  resistance is connected to the same battery if draws  $0.5\ \text{A}$ . What is the current drawn by a  $6\ \Omega$  resistance from this battery?  
 A.  $2\ \text{A}$   
 B.  $2.5\ \text{A}$   
 C.  $2.2\ \text{A}$   
 D. None of these
- Q.29 Internal resistance of ideal current source is  
 A. Infinite  
 B. Very low  
 C. Zero  
 D. Very high
- Q.30 Internal resistance is the resistance offered by \_\_\_\_\_  
 A. Source of e.m.f.  
 B. Resistor  
 C. Conductor  
 D. Capacitor
- Q.31 An electric current source is actually source of  
 A. Current  
 B. Energy  
 C. Charge  
 D. Power
- Q.32 A new flashlight cell of emf  $1.5\ \text{volts}$  gives a current of  $15\ \text{A}$ , when connected directly to an ammeter of resistance  $0.04\ \Omega$ . The internal resistance of cell is  
 A.  $0.04\ \Omega$   
 B.  $0.10\ \Omega$   
 C.  $0.06\ \Omega$   
 D.  $10\ \Omega$
- Q.33 If the current in electric bulb decreases by  $0.5\%$ , then the power in the bulb decreases by approximately  
 A.  $1\%$   
 B.  $0.5\%$   
 C.  $2\%$   
 D.  $0.25\%$
- Q.34 What will be energy used by the battery if the battery has to drive  $6.28 \times 10^{18}$  electrons with potential difference of  $20\ \text{V}$  across the terminal?  
 A.  $5\ \text{joules}$   
 B.  $15\ \text{joules}$   
 C.  $10\ \text{joules}$   
 D.  $20\ \text{joules}$
- Q.35 1 horse power = \_\_\_\_\_  
 A.  $746\ \text{kW}$   
 B.  $746\ \text{W}$   
 C.  $746\ \text{mW}$   
 D.  $746\ \text{MW}$

Topic-7

- Q.36 If  $R_1$  and  $R_2$  are respectively the filament resistances of a  $200\text{-watt}$  bulb and  $100\text{-watt}$  bulb designed to operate on the same voltage, then  
 A.  $R_1$  is two times  $R_2$   
 B.  $R_2$  is four times  $R_1$   
 C.  $R_2$  is two times  $R_1$   
 D.  $R_1$  is four times  $R_2$
- Q.37 Two electric bulbs, one of  $200\ \text{volt}$   $40\ \text{watt}$  and the other  $200\ \text{volt}$   $100\ \text{watt}$  are connected in a house wiring circuit  
 A. They have equal currents through them  
 B. The resistance of the filaments in both the bulbs is same  
 C. The resistance of the filament in  $40\text{-watt}$  bulb is more than the resistance in  $100\text{-watt}$  bulb  
 D. The resistance of the filament in  $100\text{-watt}$  bulb is more than the resistance in  $40\text{-watt}$  bulb
- Q.38 A heater coil is cut into two equal parts and only one part is now used in the heater. The heat generated will now be  
 A. halved  
 B. four times  
 C. one-fourth  
 D. doubled
- Q.39 A battery is used to light a  $24\ \text{W}$  electric lamp. The battery provides a charge of  $120\ \text{C}$  in  $60\ \text{s}$ .



- What is the potential difference across the bulb?  
 A.  $5\ \text{V}$   
 B.  $12\ \text{V}$   
 C.  $24\ \text{V}$   
 D.  $120\ \text{V}$
- Q.40 A  $100\ \text{W}$ ,  $200\ \text{V}$  bulb is connected to a  $160\ \text{V}$  supply. The actual power consumption would be  
 A.  $64\ \text{W}$   
 B.  $100\ \text{W}$   
 C.  $72\ \text{W}$   
 D.  $90\ \text{W}$
- Q.41 Electrical energy is converted to heat at the rate of \_\_\_\_\_  
 A.  $IRt$   
 B.  $I^2Rt$   
 C.  $I^2R$   
 D.  $VIt$
- Q.42 A  $40\ \text{W}$  lamp turns half the electrical energy to give light. How much light energy does it give out in  $10\ \text{s}$ ?  
 A.  $200\ \text{J}$   
 B.  $800\ \text{J}$   
 C.  $400\ \text{J}$   
 D.  $40\ \text{J}$
- Q.43 An electrical bulb marked  $100\ \text{W}$ ,  $200\ \text{V}$  would mean the resistance is  
 A.  $200\text{-ohm}$   
 B.  $50\ \text{ohm}$   
 C.  $400\ \text{ohm}$   
 D.  $50\ \text{ohm}$
- Q.44 You are given four bulbs of  $25\ \text{W}$ ,  $40\ \text{W}$ ,  $50\ \text{W}$  and  $60\ \text{W}$ . Which bulb has the lowest resistance?  
 A.  $25\ \text{W}$   
 B.  $60\ \text{W}$   
 C.  $50\ \text{W}$   
 D.  $40\ \text{W}$
- Q.45 SI unit of resistivity is \_\_\_\_\_  
 A.  $\Omega\text{-m}^2$   
 B.  $(\Omega\text{-m})^{-1}$   
 C.  $\Omega\text{-m}$   
 D.  $(\Omega\text{-m})^{-1}$
- Q.46 The following four wires are made of the same material and are at the same temperature. Which one of them has highest electrical resistance  
 A. length =  $100\ \text{cm}$ , diameter =  $1\ \text{mm}$   
 B. length =  $50\ \text{cm}$ , diameter =  $0.5\ \text{mm}$   
 C. length =  $200\ \text{cm}$ , diameter =  $2\ \text{mm}$   
 D. length =  $300\ \text{cm}$ , diameter =  $3\ \text{mm}$

Q.47 A wire has resistance 100 Ohm at 0 °C and 200 Ohm at 100 °C. What is its temperature coefficient in K<sup>-1</sup>?  
 A. -0.01  
 B. 0.01  
 C. -1/273  
 D. 1/273  
 (MCAT 2008)

Q.48 The heat produced by a current I in the wire of resistance R during time interval t is  
 A. I<sup>2</sup>Rt  
 B. I<sup>2</sup>/Rt  
 C. I<sup>2</sup>Rt  
 D. IR<sup>2</sup>t  
 (MCAT 2009)

Q.49 The fractional change in resistance per Kelvin is known as:  
 A. Temperature coefficient of resistance  
 B. Linear coefficient of expansion  
 C. Thermal coefficient  
 D. Volumetric coefficient of expansion  
 (MCAT 2009)

Q.50 The energy supplied by the cell to the charge carriers is derived from the conversion of:  
 A. Heat energy into Electrical energy  
 B. Solar energy into Electrical energy  
 C. Chemical energy into Electrical energy  
 D. Mechanical energy into Electrical energy  
 (MCAT 2009)


Q.51 The deviation of I-V graph from the straight line is due to:  
 A. Decrease in temperature and decrease in resistance  
 B. Decrease in temperature and increase in resistance  
 C. Increase in temperature and increase in resistance  
 D. Increase in temperature and decrease in resistance

Q.52 The equivalent current which passes from a point at higher potential to a point at a lower potential as if it represented a movement of positive charges is  
 A. Electronic current  
 B. Magnetic lines  
 C. Electric current  
 D. Conventional current  
 (MCAT 2010)

Q.53 The substances like germanium and silicon have  
 A. Negative temperature coefficient  
 B. Positive temperature coefficients  
 C. Both A and B  
 D. None of the above  
 (MCAT 2010)

Q.54 If 2 A current passes through a resistor when connected to a certain battery. If the resistance is replaced by the double resistance, then the current will become  
 A. 2A  
 B. 6A  
 C. 4A  
 D. 1A  
 (MCAT 2011)

Q.55 Which one of the following is I-V curve of a junction diode?  
 (MCAT 2012)



Q.56 Electric charge on an object is measured as 5 micro coulombs. How the value of this charge can be expressed in terms of base unit:  
 A. 5×10<sup>-3</sup> ampere second  
 B. 5×10<sup>-6</sup> ampere second  
 C. 5×10 ampere second  
 D. 5×10<sup>-2</sup> ampere second  
 (MCAT 2012)

Q.57 12 volt battery is applied across 6Ω resistance to have a steady flow of current. What must be the required potential difference across the same resistance to have a steady current of one ampere?  
 A. 12V  
 B. 1V  
 C. 3.V  
 D. 6.V  
 (MCAT 2013)

Q.58 Resistance between two opposite faces of square thin film of area 1mm<sup>2</sup> having thickness of 1μm if resistivity of material is 10<sup>-6</sup>Ωm will be:  
 A. 1000 Ω  
 B. 1 Ω  
 C. 100 Ω  
 D. 10 Ω  
 (MCAT 2016)

Q.59 If the potential difference across a resistor is doubled:  
 A. Only the current is doubled  
 B. Only the current is halved  
 C. Only the resistance is doubled  
 D. Only the resistance is halved  
 (ETEA 2016)

Q.60 A total charge of 100 C flows through a 12W bulb in a time of 50 second. What is the potential difference across the bulb during this time?  
 A. 0.12V  
 B. 2.0V  
 C. 6.0V  
 D. 24V  
 (ETEA 2016)

Q.61 The temperature coefficient of resistance of a semiconductor is:  
 A. Positive  
 B. Negative  
 C. Imaginary  
 D. Zero  
 (ETEA 2016)

Q.62 2 × 10<sup>6</sup> electrons passing through a conductor in 1 ms. Find electric current flowing through conductor:  
 A. 32 × 10<sup>-9</sup> A  
 B. 3.2 × 10<sup>-10</sup> A  
 C. 320 × 10<sup>-10</sup> A  
 D. 0.32 × 10<sup>-10</sup> A  
 (MDCAT 2017)

Q.63 A carbon resistor is connected to a battery of 50volt and 2ampere current is passing through it. If voltage is increased to 75volt, the current will become:  
 A. 3Amp  
 B. 4.5Amp  
 C. 1.5Amp  
 D. 6Amp  
 (MDCAT 2017)

Q.64 When the current is neither drawn from a source nor given to it then:  
 A. E = V<sub>t</sub>  
 B. E > V<sub>t</sub>  
 C. V<sub>t</sub> > E  
 D. Both "B" & "C"  
 (MDCAT 2017)

Q.65 When potential difference is applied across the ends of uniform wire of length l and radius r, a current I flow in the wire. If same potential difference is applied to the ends of another wire of the same material but of length 2l and radius 2r, the current in the wire is  
 A. I/4  
 B. I  
 C. 2I  
 D. I/2  
 (MDCAT 2018)

Q.66 The reciprocal of the conductance is called  
 A. conductivity  
 B. Resistivity  
 C. Resistance  
 D. Inductance  
 (ETEA 2018)

Q.67 A typical mobile phone of 5.0 V and an internal resistance of 200 m Ω. What is the terminal P.D of the battery when it supports a current of 500 mA?  
 A. 4.8V  
 B. 4.9V  
 C. 5.0 V  
 D. 5.1V  
 (ETEA 2018)

Q.68 A metal cube with sides of length "a" has electrical resistance R between opposite faces. What is the resistance between the opposite faces of a cube of the same metal with sides of length 3a?  
 A. 9R  
 B. 3R  
 C. R/3  
 D. R/9  
 (ETEA 2018)

Topic-7

Current Electricity

- Q.69 A filament lamp has a resistance of  $180\Omega$  when the current in it is  $500\text{mA}$ . What is the power dissipated in the lamp? (ETEA 2018)  
 A.  $45\text{W}$  B.  $90\text{W}$   
 C.  $290\text{W}$  D.  $360\text{W}$
- Q.70 A cell of internal resistance  $2.0\Omega$  and electromotive force (e.m.f)  $1.5\text{V}$  is connected to a resistor of resistance  $3.0\Omega$  what is the potential difference across  $3\Omega$  resistor. (ETEA 2018)  
 A.  $5\text{V}$  B.  $1.2\text{V}$   
 C.  $0.9\text{V}$  D.  $0.6\text{V}$
- Q.71 Calculate the rate at which energy is transferred by  $220\text{V}$  mains supply which provides a current of  $0.1\text{A}$  to a LED? (MDCAT 2019)  
 A.  $22\text{kW}$  B.  $22\text{W}$   
 C.  $2.2\text{kW}$  D.  $2.2\text{W}$
- Q.72 A copper wire has length  $L$  and cross-sectional area  $A$ . Its resistance is  $R$ . If we halved the length and halved the diameter of wire, then what will be the resistance of this wire? (MDCAT 2019)  
 A.  $R$  B.  $2R$   
 C.  $3R$  D.  $4R$
- Q.73 A car battery has EMF of  $12\text{Volts}$  and internal resistance  $5 \times 10^{-2}\text{ohm}$ . If it draws  $60$  ampere current, then terminal voltage of the battery will be (ETEA 2019)  
 A.  $5\text{volts}$  B.  $3\text{volts}$   
 C.  $15\text{volts}$  D.  $9\text{volts}$
- Q.74 One kilowatt-hour is commonly termed as one commercial unit of electric energy which is equal to (NMDCAT 2020)  
 A.  $3.6 \times 10^6\text{j}$  B.  $3.6 \times 10^6\text{j}$   
 C.  $3.6 \times 10^4\text{j}$  D.  $3.6 \times 10^3\text{j}$
- Q.75 When a wire is compressed and its radius becomes  $2R$  then its resistance will be: (NMDCAT 2020)  
 A.  $16R$  B.  $4R$   
 C.  $1/16 R$  D.  $1/4 R$
- Q.76 One of the following is an ohmic device (NMDCAT 2020)  
 A. Filament bulb B. Semiconductor diode  
 C. Transistor D. Copper wire
- Q.77 The change in a resistance of metallic conductor below  $0^\circ\text{C}$ ? (NMDCAT 2020)  
 A. Nonlinear B. Curve  
 C. Linear D. Curvilinear
- Q.78 The power of an electric bulb is  $100\text{W}$ . It is connected to  $110\text{V}$  power is supply. The resistance of electric bulb will be? (NUMS 2020)  
 A.  $11\Omega$  B.  $121\Omega$   
 C.  $20\Omega$  D.  $200\Omega$
- Q.79 If length of the wire becomes two time to the original value and area becomes one half to its original value, then resistance of the wire becomes: (NUMS 2020)  
 A. Double B. Four times  
 C. One half D. One fourth

Topic-7

Current Electricity

- Q.80 Conductivity depend on (NMDCAT 2021)  
 A. Temperature and nature of material B. Length  
 C. Area D. All
- Q.81 A load of resistance  $0.04\Omega$  is attached to the cell having E.M.F  $1.5\text{V}$  and  $15\text{A}$  is drawn in the circuit. The internal resistance of cell will be (NMDCAT 2021)  
 A.  $0.6\Omega$  B.  $0.06\Omega$   
 C.  $6\Omega$  D.  $0.3\Omega$
- Q.82 Bulb Having Current  $200\text{A}$ , and Voltage  $220$  find Power (NMDCAT 2021)  
 A.  $22\text{kW}$  B.  $44\text{kW}$   
 C.  $44\text{watt}$  D.  $40\text{kw}$
- Q.83 A heater of  $400\text{Watt}$  was on for  $5$  hours what is electrical Consumption (NMDCAT 2021)  
 A.  $20\text{KwH}$  B.  $12\text{KWH}$   
 C.  $2\text{KWH}$  D. None
- Q.84 Which of the following bulb has least resistance: (NMDCAT 2021)  
 A.  $50\text{W}$  B.  $100\text{W}$   
 C.  $200\text{W}$  D.  $300\text{W}$

# ANSWER KEY

## TOPIC-WISE MCQs & PAST PAPER MCQs


1	A	11	A	21	B	31	B	41	B	51	C	61	B	71	B	81	B
2	C	12	D	22	B	32	C	42	A	52	D	62	B	72	B	82	B
3	C	13	C	23	A	33	A	43	C	53	A	63	A	73	D	83	C
4	A	14	C	24	A	34	D	44	B	54	D	64	A	74	B	84	D
5	B	15	C	25	D	35	B	45	C	55	C	65	C	75	C		
6	D	16	C	26	A	36	C	46	B	56	B	66	C	76	D		
7	C	17	B	27	B	37	C	47	B	57	D	67	B	77	C		
8	B	18	B	28	A	38	D	48	C	58	B	68	C	78	B		
9	A	19	C	29	C	39	B	49	A	59	A	69	A	79	B		
10	B	20	C	30	A	40	A	50	C	60	C	70	C	80	A		

# EXPLANATORY NOTES

- Number of electrons,  $I = \frac{ne}{t} \Rightarrow n = \frac{It}{e}$   
 $I = (500 \times 10^{-3} \times 60) / (1.6 \times 10^{-19}) = 1.875 \times 10^{20}$
- New bulbs = 5; Old bulbs = 7; Total wattage of new bulbs =  $5 \times 50$ ;  
 Total wattage of old bulbs =  $7 \times 50$ ; Time the new bulbs are kept on = 5 hours;  
 Time the old bulbs are kept on = 3 hours  
 The number of units of electricity consumed by new bulbs ( $n_1$ ) = (total wattage  $\times$  time in hours) / 1000 =  $50 \times 5 \times 5 / 1000 = 1.25$   
 The number of units of electricity consumed by old bulbs ( $n_2$ ) =  $50 \times 7 \times 3 / 1000 = 1.05$  Total bill of electricity = number of units of electricity consumed  $\times$  amount for one unit of electricity  
 $= (n_1 + n_2) \times 60 = (1.25 + 1.05) \times 60 = 138$   
 Therefore, his electricity bill will be Rs. 138.
- The practical unit of power is horse power (hp). Kilo watt is also another practical unit of power. 1 kilowatt = 1000 watt; 1 hp = 746 watt. It is usually used in reference to the output of engines or motors.
- Current used = 30 A; Resistance = 15 ohms  
 The required equation is: Power = current<sup>2</sup>  $\times$  resistance =  $30 \times 30 \times 15 = 13,500$  Watts  
 We know that, 1 Watt = 746 horse power (hp). So, 13,500 Watts = 18.096 hp, which can approximately be equal to 18 hp.  
 Therefore, the engine consumes 18 hp power.
- Current = 3 A; Time taken = 5 minutes = 300 seconds  
 Work done = 900 J; Power = Work done / Time taken =  $900/300 = 3$  W  
 Power = Voltage (emf)  $\times$  Current  $\rightarrow$  Emf = Power/Current =  $3/3 = 1$  V  
 Therefore, the emf of the source is 1 volt.
- Total power used,  $P = P_1 + P_2 = 1500 + 500 = 2000$  W.  
 Current drawn from the supply.  
 $I = \frac{P}{V} = \frac{2000}{100} = 20$  A
- $\frac{i_1}{i_2} = \frac{R_2}{R_1} = \frac{r_2}{r_1} \times \left(\frac{r_1}{r_2}\right)^2 = \frac{3}{4} \left(\frac{2}{3}\right)^2 = \frac{1}{3}$
- For non-ohmic devices, graph is not straight line.
- $Q = It = 8 \times 10^{-3} \times 0.020 = 0.16$  mC
- Semiconductor diode is non-ohmic device.
- $I = V \Rightarrow I = \frac{V}{R}$
- $I = \frac{V}{R} \Rightarrow I \propto V$  ( $\therefore R$  - constant)
- $V = IR$
- $I = \frac{Q}{t} = \frac{ne}{t} \Rightarrow 16 \times 10^{-3} = \frac{n \times 1.6 \times 10^{-19}}{1} \Rightarrow n = 10^{17}$
- $R = \rho \frac{l}{A} \propto \frac{l}{d^2}$ . For highest resistance  $\frac{l}{d^2}$  should be maximum, which is correct for option.

- $R = \frac{4\rho L}{\pi d^2} \Rightarrow R' = \frac{4\rho(4L)}{\pi(4d)^2} = \frac{4\rho L}{\pi d^2} \Rightarrow R' = R$
- $R_1 \propto \frac{l}{A} \Rightarrow R_2 \propto \frac{2l}{2A}$  i.e.  $R_2 \propto \frac{l}{A} \therefore R_1 = R_2$
- Because  $V-i$  graph of diode is non-linear.
- $\rho = \frac{RA}{L} = \frac{R\pi r^2}{L} \Rightarrow \rho = \frac{R\pi d^2}{4L} = \frac{0.7 \times 3.14 \times 4 \times 10^{-6}}{4 \times 1} = 2.2 \times 10^{-6} \Omega m$
- Iron and platinum (information from text book).
- $R' = \frac{\rho L'}{A'} = \frac{\rho(2L)}{A/2} = \frac{4 \times \rho L}{A} \Rightarrow R' = 4R$
- Resistivity does not depend upon dimension ( $L, W$ ) of wire. It only depends upon nature & temperature of wire.
- $R = \frac{\rho L}{A}$   
 $L' = 4L \Rightarrow A' = \frac{A}{4} \Rightarrow \boxed{R' = 16R}$
- $R = \frac{\rho L}{A} \Rightarrow \boxed{R \propto L}$
- Temperature coefficient of resistance =  $\alpha_{-ve \text{ for Ge}}^{+ve \text{ for Cu}}$   
 $\Rightarrow \alpha = +ve$  for Cu  $\Rightarrow T \uparrow \Rightarrow R \uparrow$  or  $T \downarrow \Rightarrow R \downarrow$   
 &  $\alpha = -ve$  for Ge  $\Rightarrow T \uparrow \Rightarrow R \downarrow$  or  $T \downarrow \Rightarrow R \uparrow$   
 So resistance of Cu decreases & that of Ge increases by cooling or decreasing temperature.
- $R \propto L$
- $v_t = \epsilon - Ir$
- $v = \epsilon - I_1 r \Rightarrow I_1 R_1 = \epsilon - I_1 r$   
 $(1.6)(8) = \epsilon - 1.6r \Rightarrow \boxed{12.8 = \epsilon - 1.6r} \rightarrow (i)$   
 $(15 - 12.8) = (\epsilon - 1.6r) - (\epsilon - 0.5r)$   
 $2.2 = -1.1r \Rightarrow \boxed{r = 2}$   
 $\epsilon = 16V$
- $I_2 R_2 = \epsilon - I_2 r \Rightarrow \boxed{15 = \epsilon - 0.5r} \rightarrow (ii)$   
 $\epsilon = I(R + r) \Rightarrow 16 = I(6 + 2) \Rightarrow \boxed{I = 2A}$
- Internal resistance of ideal current source is zero.
- Internal resistance is the resistance offered by source of e.m.f.
- Source of emf provides energy.
- $V_t = \epsilon - Ir \Rightarrow IR = \epsilon - Ir$   
 $(15)(0.04) = 1.5 - (15)(r)$   
 $616 = 1.5 - 15r \Rightarrow 15r = \frac{0.9}{15} = 0.06 \Omega$

33. %age increase in  $P = 2(\text{%age increase in } I) \Rightarrow \text{%age increase in } P = 2(0.5\%) = 1\%$
34.  $Q = ne = 6.28 \times 10^{18} \times 1.6 \times 10^{-19} = 10 \times 10^{-1} = 1C \Rightarrow \Delta V = \frac{W}{Q} \Rightarrow W = Q\Delta V = 1 \times 20 = 20J$
35.  $I_{hp} = 746 \text{ Watt}$
36.  $P \propto \frac{I}{R} \Rightarrow \frac{P_1}{R_1} = \frac{P_2}{R_2} \Rightarrow \frac{200}{100} = \frac{R_2}{R_1} \Rightarrow R_2 = 2R_1$
37.  $P = \frac{V^2}{R} \Rightarrow R_1 = \frac{V_1^2}{P_1} = \frac{(200)^2}{40} = 1000\Omega$  and  $R_2 = \frac{V_2^2}{P_2} = \frac{(200)^2}{100} = 400\Omega$
- So,  $R_1 > R_2$
38.  $H = I^2 R t \Rightarrow H' = (2I)^2 \left(\frac{R}{2}\right) t \Rightarrow H' = 2H$
39.  $P = VI = V \frac{Q}{t}$   
 $V = \frac{Pt}{Q} = \frac{24 \times 60}{120} = 12V$
40.  $R = \frac{V^2}{P} = \frac{(200)^2}{100} = 400 \Omega \Rightarrow P = \frac{V^2}{R} = \frac{(160)^2}{400} = 64 \text{ Watt}$
41.  $H = I^2 R t$
42.  $E = P \times t = (20)(10) = 200J$
43.  $P = \frac{V^2}{R} \Rightarrow R = \frac{V^2}{P} = \frac{200^2}{100} = 400\Omega$
44. The resistance of high power devices is smaller than the low power ones. The resistance of 60 watt bulb is smaller than 40 watt bulb and so on  $P = \frac{V^2}{R}$  For a given voltage,  $P \propto \frac{1}{R}$ . So, 60 W bulb has the lowest resistance.
45.  $\rho = \frac{RA}{L} = \frac{\Omega m^2}{m} = \Omega m$
46.  $R \propto \frac{l}{r^2}$ . For highest resistance  $\frac{l}{r^2}$  should be maximum, which is correct for option.
47.  $\alpha = \frac{R_1 - R_2}{R_2 t} = \frac{200 - 100}{100 \times 100} = \frac{100}{100 \times 100} = 0.01$
48. According to Joule's Law  $H = I^2 R t$
49. Definition of temperature co-efficient of resistance.
50. The energy supplied by the cell to the charge carrier is derived from conversion of chemical energy into electrical energy.
51. The deviation of I-V graph means, Temperature is not constant. So  $\Delta R \propto \Delta t$

52. Conventional current.
53. The substances like germanium and silicon have negative temperature co-efficient of resistance.
54.  $V = \text{same}$   
 $I \propto \frac{1}{R} \therefore I = 2A$  if  $R' = 2R$   $I' = \frac{I}{2} = \frac{2}{2} = 1A$
55. 
56.  $Q = 5\mu C \Rightarrow Q = 5 \times 10^{-6} \text{ As}$
57.  $\frac{V_2}{V_1} = \frac{I_2 R_2}{I_1 R_1} \therefore I_1 = \frac{V}{R} = \frac{12}{6} = 2A$   
 $V_2 = \frac{I_2 R_2}{I_1 R_1} \times V_1 \Rightarrow V_2 = \frac{1 \times 6}{2 \times 6} \times 12 = 6V$
58.  $R = \rho \frac{L}{A} \Rightarrow \frac{\rho L}{L t} = \frac{\rho}{t} = \frac{10^{-6}}{10^{-6}} = 1\Omega$
59. Ohm law  $V \propto I$
60.  $V = \frac{P}{I} = \frac{P}{\frac{Q}{t}} = \frac{P \times t}{Q}$
61.  $\alpha = \frac{\Delta R}{R_0 \Delta T}$
62.  $I = \frac{Q}{t} = \frac{ne}{t} = \frac{2 \times 10^6 \times 1.6 \times 10^{-19}}{10^{-1}} = 3.2 \times 10^{-10} A$
63.  $V \propto I, \frac{V_2}{V_1} = \frac{I_2}{I_1} \Rightarrow I_2 = \frac{V_2}{V_1} \times I_1 = \frac{75}{50} \times 2 \Rightarrow I_2 = 3A$
64.  $vt = \epsilon - Ir \Rightarrow I = 0 \Rightarrow vt = \epsilon$
65.  $I = \frac{V}{R}, R = \frac{\rho L}{A} = \frac{\rho L}{\pi r^2}$   
 $R' = \frac{\rho(2L)}{\pi(2r)^2} = \frac{2\rho L}{4A} = \frac{R}{2} \Rightarrow I' = \frac{V}{R/2} = 2 \frac{V}{R} = 2I$
66.  $G = \frac{1}{R}$
67.  $E = I(R+r)$
68. Parallel combination of resistance
69.  $P = I^2 R$
70.  $I = 1.5/3 + 2 = 1.5/5$  now  $V = IR = 1.5/5 \times 3 = 0.9V$

$$71. R = \frac{V}{I} = \frac{220}{0.1} = 2200\Omega \Rightarrow P = \frac{V^2}{R} = \frac{(220)^2}{2200} = 22W$$

$$72. R = \frac{\rho L}{A} = \frac{\rho L}{\pi r^2} = \frac{\rho L}{\pi \frac{D^2}{4}} \Rightarrow R = \frac{4\rho L}{\pi D^2}$$

$$R' = \frac{4\rho L/2}{\pi \left(\frac{D}{2}\right)^2} = \frac{4\rho L}{2\pi D^2/4} = \frac{2(4\rho L)}{\pi D^2} = 2R$$

$$73. E = V + Ir$$

$$74. P = \frac{E}{t} \Rightarrow E = P \times t = 1000 \times 3600 \Rightarrow P = 3.6 \times 10^6 J$$

$$75. R = \rho \frac{L}{A} = \frac{\rho \times l \times A}{A^2} = \rho \frac{V}{A^2}$$

$$R = \rho \frac{V}{\pi^2 R^4} \Rightarrow R \propto \frac{1}{R^4}$$

76. For copper wire there is straight line graph of I-V

77. This is linear behavior between change in resistance w.r.t temperature below 0°C

$$78. P = \frac{V^2}{R}$$

$$79. R = \frac{\rho L}{A}$$

80. Conductivity depends on temperature and nature of material.

$$81. r = \frac{\epsilon - vt}{I} = \frac{1.5 - 15 \times 0.04}{15} = 0.06\Omega$$

$$82. P = IV = 200 \times 220 = 44 kW$$

$$83. \text{Electrical consumption} = \frac{P(W) \times t(\text{hours})}{1000} = \frac{400 \times 5}{1000} = 2 kWh$$

84.

$$P = \frac{V^2}{R}$$



$$P \propto \frac{1}{R}$$


# 8 TOPIC

# ELECTROMAGNETISM

## PRACTICE EXERCISE

- Q.1 Which of the following will experience a maximum force, when projected with the same velocity perpendicular to the magnetic field : (i)  $\alpha$ -particle, and (ii)  $\beta$ -particle?  
 A. Both  $\alpha$ -particle and  $\beta$ -particle  
 B. None  
 C.  $\beta$ -particle  
 D.  $\alpha$ -particle
- Q.2 Two parallel wires carrying currents in the opposite directions  
 A. Repel each other  
 B. Attract each other  
 C. Have no effect upon each other  
 D. They cancel out their individual magnetic fields
- Q.3 The direction of the magnetic lines of force depends upon:  
 A. Nature of the material of the conducting wire  
 B. Amount of the current  
 C. Area of the conducting wire  
 D. Direction of the current
- Q.4 The direction of magnetic field at a point on the magnetic lines of force can be taken along:  
 A. Normal at that point  
 B. Axis of the magnetic line of force at that point  
 C. The tangent at that point  
 D. Can't be taken
- Q.5 The direction of the magnetic lines of force can be found by using:  
 A. Right hand rule  
 B. Left hand rule  
 C. Henry's law  
 D. Faraday's law
- Q.6 A uniform magnetic field is represented by a set of lines of force which are  
 A. Parallel  
 B. Convergent  
 C. Divergent  
 D. None of these
- Q.7 When a current carrying conductor is placed in a magnetic field. It moves from a region of  
 A. Stronger to weak field  
 B. Weak to strong field  
 C. Strong to weak if current is large  
 D. Weak to strong if current is large
- Q.8 Magnetic lines of force  
 A. Always intersect  
 B. Tend to crowd far away from the poles of a magnet  
 C. Are always in closed shape  
 D. Do not pass through vacuum
- Q.9 A magnetic needle is kept in a non-uniform magnetic field. It experiences  
 A. Neither a force on a torque  
 B. A force but not a torque  
 C. A torque but not a force  
 D. A force and a torque
- Q.10 Which of the following quantity is not affected by a magnetic field?  
 A. Stationary charge  
 B. Change in magnetic flux  
 C. Moving charge  
 D. Current flowing in a conductor


- Q.11 A straight current carrying conductor is shown in the figure. A person observes it from below at point "B". What must be direction of magnetic field observed?  
 A. Clock-wise  
 B. Out of paper  
 C. Anti-clock wise  
 D. Into the paper
- 
- Q.12 Magnetic field has its application in \_\_\_\_\_  
 A. Motors  
 B. Current detector  
 C. Generators  
 D. All
- Q.13 A conductor is shown in the figure connected with the terminals of a source as shown. What is true about this?  
 A.  $\vec{B}$  is present inside conductor  
 B.  $\vec{B}$  and  $\vec{E}$  both are present inside conductor  
 C.  $\vec{B}$  is present outside conductor  
 D.  $\vec{B}$  is outside while  $\vec{E}$  is inside conductor
- 
- Q.14 A square coil  $10^{-2}m^2$  area is placed perpendicular to a uniform magnetic field of strength  $10^3$  weber/ $m^2$ . The magnetic flux through the coil.  
 A. 10 weber  
 B.  $10^5$  weber  
 C.  $10^3$  weber  
 D.  $10^6$  weber
- Q.15 A magnetic field of 2.5 T passes perpendicular to a disc of radius 2cm. Find the magnetic flux associated with the disc.  
 A.  $3.14 \times 10^{-3}$  Wb  
 B.  $3.14 \times 10^{-4}$  Wb  
 C.  $3.14 \times 10^{-2}$  Wb  
 D.  $3.14 \times 10^{-5}$  Wb
- Q.16 A coil of area of cross section  $10^{-2}m^2$  and 100 turns is placed in a magnetic field of 1T, with its axis making an angle  $60^\circ$  with the field. Find the total flux associated with the field.  
 A. 0.5 Wb  
 B. 0.3 Wb  
 C. 0.4 Wb  
 D. 0.2 Wb
- Q.17 When the coil is rotated between the pole pieces of a magnet, during one complete rotation of the coil, how often will the magnetic flux linked with coil be maximum and minimum?  
 A. Maximum and minimum once each  
 B. Maximum once minimum twice  
 C. Maximum and minimum twice each  
 D. Maximum twice, minimum once
- Q.18 When area is placed perpendicular to external magnetic field the magnetic flux will be?  
 A. Minimum  
 B. In between maximum and minimum  
 C. Maximum  
 D. None
- Q.19 Direction of magnetic flux is  
 A. Normal to the surface  
 B. At any angle  
 C. Parallel to the surface  
 D. No direction
- Q.20 If 0.5 T magnetic field is perpendicular to the  $0.5 m^2$  area then the magnetic flux would be  
 A. 0.25 Wb  
 B. 1.25 Wb  
 C. 6.25 Wb  
 D. Zero

- Q.21 20 Wb magnetic flux passes through the  $5m^2$  area of certain sheet, the magnetic flux density would be  
 A.  $2 Wb m^{-2}$   
 B.  $6Wb m^{-2}$   
 C.  $4 Wb m^{-2}$   
 D.  $8Wb m^{-2}$
- Q.22 The Relationship between tesla(T) and smaller unit gauss(G) of magnetic induction is given by  
 A.  $1T = 10^3 G$   
 B.  $1T = 10^4 G$   
 C.  $1T = 10^{-2} G$   
 D.  $1T = 10^8 G$
- Q.23 Weber ampere per metre is equal to  
 A. Joule  
 B. Newton  
 C. Watt  
 D. Henry
- Q.24 The unit of magnetic flux density is  
 A.  $Wb m^{-2}$   
 B.  $NA^{-1} m^{-1}$   
 C. Tesla  
 D. All of these
- Q.25 The magnetic induction B is also called the  
 A. Flux  
 B. Flux density  
 C. Density  
 D. Tesla
- Q.26 A 2 MeV proton is moving perpendicular to a uniform magnetic field of 2.5 T. The force on the proton is:  
 A.  $2.5 \times 10^{-10} N$   
 B.  $2.5 \times 10^{-11} N$   
 C.  $8 \times 10^{-11} N$   
 D.  $8 \times 10^{-12} N$
- Q.27 The magnitude of the force on a moving charge is maximum when angle between the velocity of the charge and the magnetic field is,  
 A.  $0^\circ$   
 B.  $90^\circ$   
 C.  $180^\circ$   
 D.  $45^\circ$
- Q.28 If a charge particle enters a uniform magnetic field, there is a change in its  
 A. Kinetic energy  
 B. Direction of velocity  
 C. Magnitude of velocity  
 D. All of these
- Q.29 The fig shows a uniform magnetic field  $\vec{B}$  directed into the plane of paper. A particle with negative charge moves in the plane, which of four paths 1, 2, 3 or 4 does the particle follow
- 
- A. Path 1  
 B. Path 3  
 C. Path 2  
 D. Path 4
- Q.30 Which of the following Statements is false?  
 A. A stationary charge produces a constant electric field  
 B. An accelerated charge produces combination of varying electric and magnetic field  
 C. A moving charge with uniform speed produces a constant magnetic field.  
 D. A conductor carrying steady current has no electric field in it




c-8

In the given figure the electron enters into the magnetic field. It deflects in.....direction







- B. -ve X direction
- D. -ve Y direction

- A. +ve X direction
  - C. +ve Y direction
- Q.32 In the formula  $F = q(v \times C)$ .
- A. F must be perpendicular to v but not necessarily to B
  - B. v must be perpendicular to B but not necessarily to F
  - C. F must be perpendicular to both v and B
  - D. All three vectors must be mutually perpendicular
- Q.33 In the figure below, what is the direction of the magnetic force  $F_B$ ?



- A. To the right
  - C. Upward, in the plane of the page
- Q.34 Which of the following cannot be deflected by a magnetic field?
- A. Alpha rays
  - B. Gamma rays
  - C. Beta rays
  - D. Cosmic rays

Q.35 The following diagrams shows an electron passing through a magnetic field. Which diagram shows the possible path of the electrons as they pass through the field?




- Q.36 Work done by magnetic force on a moving charge particle is
- A. Positive
  - B. Zero
  - C. Negative
  - D. Infinite

- Q.37 A positively charged particle moving due east enters a region of uniform magnetic field directed vertically upwards. The particle will
- A. Continue to move due east
  - B. Move in a circular orbit with its speed increases
  - C. Move in a circular orbit with its speed unchanged
  - D. Get deflected vertically upwards


Topic-8

Electromagnetism

- Q.38 A strong magnetic field is applied to a stationary electron, then
- A. Electron move in the direction of field
  - B. Electron start spinning
  - C. Electron move opposite to field
  - D. Electron remain stationary
- Q.39 The magnitude of force on a moving charge is zero then angle between the velocity of the charge carrier and magnetic field is
- A.  $0^\circ$
  - B.  $45^\circ$
  - C.  $90^\circ$
  - D.  $120^\circ$
- Q.40 An electron enters in a uniform magnetic field making an angle  $60^\circ$  with field. The shape of its trajectory in magnetic field is
- A. Circle
  - B. Straight line
  - C. Parabola
  - D. Helix
- Q.41 Which of the following in motion cannot be deflected by magnetic field?
- A. Electron
  - B. Proton
  - C. Neutron
  - D. Sodium ion
- Q.42 When a charged particle is projected perpendicularly in a magnetic field its trajectory is
- A. Hyperbola
  - B. Parabola
  - C. Helix
  - D. Circular
- Q.43 When a charged particle moves through a magnetic field, it suffers a change in
- A. Direction
  - B. Energy
  - C. Speed
  - D. No change
- Q.44 Magnetic field is caused by
- A. Stationary charge
  - B. A moving negative charge only
  - C. A moving positive charge only
  - D. Moving positive and negative charges both
- Q.45 A charged particle moves through a magnetic field in a direction perpendicular to it. Which of the following remain unchanged for the particle?
- A. Velocity
  - B. Speed
  - C. Acceleration
  - D. Direction
- Q.46 The diagram shows an electron as it enters magnetic field. The path of the electron and the magnetic field are in the plane of the paper. In which direction is the electron initially deflected?



- A. into the plane of the paper
  - B. to the left of its incoming path
  - C. out of the plane of the paper
  - D. to the right of its incoming path
- Q.47 An electron moves in the negative x direction, through a uniform magnetic field in the negative y direction. The magnetic force on the electron is



- A. In the negative z direction
- B. In the negative y direction
- C. In the positive z direction
- D. In the positive y direction

PAST PAPER MCQs


Q.48 An electric charge in uniform motion produces:  
 A. An electric field  
 B. A magnetic field  
 C. Both magnetic and electric fields  
 D. Neither magnetic nor electric fields  
 (MCAT 2008)

Q.49 A charge of two micro coulombs ( $2 \mu\text{C}$ ) moves with velocity of two meter per second ( $2 \text{ m/sec}$ ) in the direction of two Tesla magnetic field. The force that will act on it will be:  
 A. 2N  
 B. Zero  
 C. 8N  
 D. 4N  
 (MCAT 2009)

Q.50 Force experienced by a moving charge in a magnetic field is:  
 A.  $F = BA \cos\theta$   
 B.  $F = \mu_0 NI$   
 C.  $F = q(v \times B)$   
 D.  $F = I(L \times B)$   
 (MCAT 2010)

Q.51 Which one of the following relations is correct?  
 A.  $1 \text{ wb m}^{-2} = \text{N m}^{-1} \text{ A}^{-1}$   
 B.  $1 \text{ Tesla} = 10^4 \text{ Gauss}$   
 C.  $1 \text{ wb m}^{-2} = 1 \text{ Tesla}$   
 D. All of these  
 (MCAT 2010)


Q.52 Two long parallel wires Wire 1 and Wire 2 repel each other as shown in the figure. What could be the reasons?  
 (MCAT 2011)



A. Both carry current in same direction  
 B. Both carry current in opposite direction  
 C. Wire 1 has current, but Wire 2 has no current  
 D. Wire 2 has current, wire 1 has no current  
 (ETEA 2011)

Q.53 The SI unit of magnetic flux is weber which is equal to:  
 A.  $\text{NmA}^{-1}$   
 B.  $\text{Nm}^2\text{A}^{-1}$   
 C.  $\text{Nm}^{-1}$   
 D.  $\text{AmA}^{-2}$

Q.54 Two long straight parallel wires held vertically, have equal but opposite currents as shown in the figure.  
 (MCAT 2012)



Which of the following effect will be observed?  
 A. Magnetic field at 'X' is stronger than that at 'Y' and 'Z'  
 B. Magnetic field at 'X', and 'Z' are same  
 C. Magnetic field at 'X' is weaker than that at Z  
 D. Magnetic field at 'X' is weaker than that at out, stronger than that at 'Z'

Q.55 A long straight current carrying conductor has current direction from bottom to top when held vertically. What will be the direction of magnetic field lines when observed from below the conductor?  
 (MCAT 2013)


A. Clockwise  
 B. Vertically upward  
 C. Anticlockwise  
 D. Vertically downward

Q.56 Due to current in straight conductor the distance between magnetic field lines.  
 A. Increases away from conductor  
 B. Increases towards conductor  
 C. Decreases away from conductor  
 D. Decreases and then increases towards conductor  
 (MCAT 2014)

Q.57 If the value of magnetic flux is  $10 \text{ Wb}$ , when magnetic lines of force containing magnetic field strength of  $1 \text{ Tesla}$  passing through unit area of  $10 \text{ m}^2$  then the angle between magnetic field and unit area is:  
 A.  $180^\circ$   
 B.  $360^\circ$   
 C.  $90^\circ$   
 D.  $45^\circ$   
 (MDCAT 2017)

Q.58 A charge is projected with velocity of  $10 \text{ m/s}$  in a magnetic field of  $10 \text{ T}$  at angle of  $60^\circ$ . If force of  $2.78 \times 10^{-17} \text{ N}$  is exerted on the charge then value of charge will be:  
 A.  $1.60 \times 10^{-19} \text{ C}$   
 B.  $2.70 \times 10^{-19} \text{ C}$   
 C.  $4.80 \times 10^{-19} \text{ C}$   
 D.  $3.20 \times 10^{-19} \text{ C}$   
 (MDCAT 2017)

Q.59 Electric current is flowing through a straight conductor as shown in figure given below. The direction of magnetic lines of force will be  
 (MDCAT 2017)



A. Anticlockwise  
 B. Clockwise  
 C. From Bottom to Top  
 D. From Top to Bottom

Q.60 Two long, parallel conductors which are free to move are arranged  $1.0 \text{ cm}$  apart. A steady current of  $20 \text{ A}$  flows in each of the conductor in the same direction. The conductors  
 A. remain stationary  
 B. move towards each other  
 C. move away from each other  
 D. move at right angles to each other  
 (MDCAT 2018)

Q.61 A neutron having mass equal to a proton ( $m_p = 1.6 \times 10^{-27} \text{ kg}$ ) is moving in a magnetic field of intensity  $1.20 \times 10^{-3} \text{ T}$  with a speed of  $2.0 \times 10^{-7} \text{ ms}^{-1}$  what is the Maximum force experienced by the neutron.  
 A.  $3.84 \times 10^{-15} \text{ N}$   
 B.  $0 \text{ N}$   
 C.  $3.84 \times 10^{-12} \text{ N}$   
 D.  $38.4 \times 10^{-15} \text{ N}$   
 (MDCAT 2018)

Q.62  $e/m$  of an electron is given by the relationship,  
 A.  $e/m = 2(V/B^2 r^2)$   
 B.  $e/m = (V/Br)^2$   
 C.  $e/m = Vr/B$   
 D.  $e/m = VB/r$   
 (MDCAT 2018)

Q.63 Magnetic flux is maximum when angle between magnetic field and vector area is  
 A.  $0^\circ$   
 B.  $90^\circ$   
 C.  $180^\circ$   
 D.  $45^\circ$   
 (NMDCAT 2020)

Q.64 Formula of toroid for magnetic field is  
 A.  $\mu_0 NI$   
 B.  $\frac{\mu_0 NI}{2\pi r}$   
 C.  $\frac{\mu_0 NI}{2\pi}$   
 D.  $\mu_0 I$   
 (NMDCAT 2021)

- Q.65 Unit of Magnetic flux  
 A.  $\text{Tm}^2$   
 C. Both A and B
- Q.66 Magnitude of magnetic field of Earth  
 A. 0.6 G  
 C. 6T
- Q.67 Magnetic force depends upon  
 A. Current  
 C. Magnetic field
- B. Weber  
 D. Tesla
- B. 0.4 T  
 D. 4T
- B. Length  
 D. All

(NMDCAT 2021)

(NMDCAT 2021)

(NMDCAT 2021)

# ANSWER KEY

## TOPIC-WISE MCQs & PAST PAPER MCQs

1	D	11	C	21	C	31	D	41	C	51	D	61	B
2	A	12	D	22	D	32	C	42	D	52	B	62	A
3	D	13	D	23	B	33	D	43	A	53	A	63	A
4	C	14	A	24	D	34	B	44	D	54	A	64	B
5	A	15	A	25	B	35	B	45	B	55	A	65	C
6	A	16	A	26	D	36	B	46	C	56	A	66	A
7	A	17	C	27	B	37	C	47	A	57	A	67	D
8	C	18	C	28	B	38	D	48	C	58	D		
9	D	19	D	29	B	39	A	49	B	59	B		
10	A	20	A	30	D	40	D	50	C	60	B		

# EXPLANATORY NOTES

- $F = qvB$ .  
 For  $\alpha$ -particle,  $q = 2e$ ,  $F_\alpha = 2evB$   
 For  $\beta$ -particle,  $q = e$ ,  $F_\beta = evB$ .  
 Thus, the  $\alpha$ -particle will experience maximum force.
- Two parallel wires having current in opposite direction always repel each other because magnetic field between the wires become stronger.
- Direction of magnetic field depends upon direction of current according to right hand rule.
- Direction of magnetic field is along the tangent on a curve.
- Right hand rule indicates the direction of magnetic lines of force
- Parallel lines of forces produce uniform field.
- Current carrying conductor will move from stronger to weaker magnetic field.
- Because magnetic monopoles do not exist in nature.
- Magnetic force produces force and torque on magnetic needle due to position of magnetic needle.
- $F_m = qvB \Rightarrow v = 0$ ,  $F_m = 0$  for stationary charge.
- According to right hand rule the direction of magnetic field line will be anti-clock-wise.
- Motors, generators and current detectors all are applications of magnetic field.
- Magnetic field is always produced around the electric field.
- $\phi = BA = 10^{-2} \times 10^3 = 10$  weber
- $\phi = BA \cos\theta = B (\pi r^2) \cos\theta = 2.5 \times \pi (2 \times 10^{-2})^2 \cos 0^\circ = 3.14 \times 10^{-3}$  wb
- $\phi = nBA \cos\theta = 100 \times 1 \times 10^{-2} \times \cos 60^\circ = 0.5$  wb
- In one complete rotation of coil, magnetic flux linked with coil two times maximum and two times minimum.
- $\phi = BA \cos\theta = BA \cos 0^\circ = BA$  (maximum)
- Magnetic flux is a scalar quantity.
- $\phi = B.A = 0.5 \times 0.5 = 0.25$  Wb
- $B = \frac{\phi}{A} = \frac{20}{5} = 4$  Wb  $\text{m}^{-2}$
- $1T = 10^4$ G
- $\frac{\text{weber} \times \text{ampere}}{\text{meter}} = \frac{\text{NmA}^{-1} \cdot \text{A}}{\text{m}} = \text{N}$
- Basic from unit conversion.
- Book information.
- $K.E = \frac{1}{2}mv^2 \Rightarrow v = \sqrt{\frac{2K.E}{m}}$  ;  $F = qvB$
- $F = qvB \sin \theta$ ,  $F_{\max} = qvB$   $\because \sin 90^\circ = 1$
- Magnetic force is just a deflecting force.

29. According to right hand rule the electron will be deflected downward when magnetic field is into the plane.
30. When an electron charge will move through a conductor it will carry its electric field.
31. According to right hand rule the electron will be deflected downward when magnetic field is into the plane.
32. By using the concept of cross product the resultant force will be perpendicular to both  $\vec{v}$  and  $\vec{B}$ .
33. By using right hand rule the magnetic force will be out of the plane of the page.
34.  $F = qvB \sin \theta$ ,  $F = 0$   $\therefore q = 0$
35. Application of right hand palm rule or Fleming's left hand rule.
36. Magnetic force is perpendicular to velocity of the charge particle so no work is done.
37. According to  $\vec{F} = q(\vec{v} \times \vec{B})$   
In perpendicular magnetic field, the path of a charged particle is a circle, and the magnetic field does not cause any change in speed and energy.
38.  $F = qvB \sin \theta$ ,  $F = 0$   $\therefore v = 0$
39.  $F = qvB \sin \theta \Rightarrow F = 0$  when  $\theta = 0^\circ$
40. If angle equal to  $90^\circ$ , then trajectory will be circle and  $90^\circ > \theta > 0^\circ$  trajectory will be helix.
41. Only charge particle can deflect in a magnetic field.
42. When  $\theta = 90^\circ$  path is circular
43. Magnetic field is deflecting field which only change the direction.
44. Moving charge (may +ve or -ve) produce magnetic field
45. Magnetic field does no work so, K.E of charge remain same, so speed also remain same.
46. Electron will be deflected out of plane of paper due to magnetic force by right hand rule.
47. By using the concept of cross product the direction of force on electron will be in negative z direction.
48. When an electronic charge is at rest, it produces only electric field but when electric charge is in uniform motion, it produces both magnetic and electric field.
49.  $F = qvbB \sin \theta$   $\therefore \theta = 0^\circ \Rightarrow F = qvB \sin 0^\circ = 0$
50. Force on moving charge is  $\vec{F} = q(\vec{v} \times \vec{B})$
51. All relations are correct.
52. Unlike currents in wires repel each other.
53.  $\phi_b = B \cdot A = \frac{F}{IL} A$
54. At "X" two fields reinforce each other by R.H.R. So magnetic field at "X" is stronger than that at "Y" and "Z".
55. By R.H.R the direction of magnetic field lines in a current carrying wire is clock-wise.

56.  $B \propto \frac{1}{r}$  Near the conductor, magnetic field lines are closer while away from conductor, magnetic field lines are wider.
57.  $\phi = BA \cos \theta$   
 $\theta = \cos^{-1} \left( \frac{\phi}{BA} \right) = \cos^{-1} \left( \frac{10}{1 \times 10} \right) = \cos^{-1}(1) = 0^\circ \text{ or } 180^\circ$
58.  $F = qvB \sin \theta$   
 $q = \frac{F}{vB \sin \theta} = \frac{2.78 \times 10^{-17}}{10 \times 10 \times \sin 60} = \frac{2.78 \times 10^{-19} \times 2}{\sqrt{3}} = \frac{5.56 \times 10^{-19}}{1.73} = 3.2 \times 10^{-19} \text{ C}$
59. Use Conventional Current and R.H.R
60. When current flows in same direction in the wires, wires attract each other.
61.  $F = qvB \sin \theta \Rightarrow q = 0$  for neutron  $\Rightarrow F = 0$
62.  $\frac{e}{m} = \frac{2V}{B^2 r^2}$
63.  $\phi m = \vec{B} \cdot \vec{A} = BA \cos 0^\circ$  so, flux will be maximum.
64.  $B = \frac{\mu_o NI}{2\pi r}$
65. Unit of magnetic flux  $\phi = BA = Tm^2 = Wb$
66. Magnetic field at its surface ranges from 25 to 65  $\mu T$  (0.25+0.65G)
67.  $F = ILB \sin \theta$

# 9 TOPIC

## ELECTROMAGNETIC INDUCTION PRACTICE EXERCISE

Q.1 A step-down transformer reduces the supply voltage from 220 V to 11 V and increase the current from 6 A to 100 A. Then its efficiency is  
 A. 1.2  
 B. 0.83  
 C. 0.12  
 D. 0.9


Q.2 In a transformer, the number of turns in the primary and the secondary are 410 and 1250 respectively. If the current in primary is 6A, then that in the secondary coil is  
 A. 2 A  
 B. 18 A  
 C. 12 A  
 D. 1 A

Q.3 Which of the following will not generate emf?  
 A. Holding a magnet stationary inside a coil  
 B. Rotating a coil around a stationary coil  
 C. Rotating a coil in a magnetic field  
 D. Moving a bar magnet across a flat piece of metal

Q.4 Which one of the following does not affect the magnitude of the induced emf in electromagnetic induction?  
 A. The strength of the magnetic field linking to the coil  
 B. The speed with which the coil cuts the magnetic field  
 C. The resistance of the coil cutting the magnetic field  
 D. The number of turns in the coil

Q.5 Relative motion between a \_\_\_\_\_ and a conducting coil produces current in the coil  
 A. Magnet  
 B. Insulator  
 C. Iron bar  
 D. All of these

Q.6 In a closed ring (a) and in an open ring (b) magnets are falling along the axis of the ring. The current generated in a and b have directions




A. Clockwise, Zero  
 C. Anticlockwise, zero

B. Anticlockwise, clockwise  
 D. Zero, zero

Q.7 The direction of induced current in a coil or circuit is such that it opposes every cause of its production. This law is given by

A. Faraday  
 C. Kirchhoff  
 B. Lenz  
 D. Ampere

Q.8 A metallic circular ring is suspended by a string and is kept in a vertical plane. When a magnet is approached towards the ring then it will



A. Remain stationary  
 C. Get displaced away from the magnet

B. Get displaced towards the magnet  
 D. Nothing can be said

## Electromagnetic Induction


Q.9 In the equation  $\epsilon = -N \frac{\Delta\phi}{\Delta t}$  the negative sign indicates that  
 A. The self-induced current opposes its cause  
 B. The induced e.m.f opposes the change which produces it.  
 C. The accordance of equation with the Lenz's Law  
 D. All of the above

Q.10 The e.m.f. induced in a coil is ..... the rate of change in flux linkages.  
 A. Directly proportional to  
 C. Inversely proportional to  
 B. Independent of  
 D. None of the above

Q.11 A coil having 500 square loops each of side 10 cm is placed normal to a magnetic induction which increases at the rate of 1.0 tesla/second. The induced e.m.f. in volts is  
 A. 5  
 C. 0.1  
 B. 1  
 D. 0.5

Q.12 In a circuit with a coil of resistance 2 ohms, the magnetic flux changes from 2.0 Wb to 10.0 Wb in 0.2 second. The charge that flows in the coil during this time is  
 A. 5.0 coulomb  
 C. 4.0 coulomb  
 B. 1.0 coulomb  
 D. 0.8 coulomb

Q.13 A magnet falls with its S-pole along the axis of ring. The current generated is .... And acceleration is



A. Clockwise, > g  
 C. Clockwise, < g

B. Anticlockwise, > g  
 D. Anticlockwise, < g


Q.14 Lenz's law is a manifestation of the conservation of  
 A. Current  
 C. Voltage  
 B. Energy  
 D. All of these

Q.15 When a coil of cross-sectional area  $A$  and number of turns  $N$  is rotated in a uniform magnetic field  $B$  with angular velocity  $\omega$ , then the maximum emf induced in the coil will be

A.  $BNA$   
 C.  $\frac{Ba\omega}{N}$

B.  $BNA\omega$   
 D. Zero

Q.16 The graph shows how an alternating current  $I$  of peak value  $I_0$  varies with time  $t$ .



Which expression gives the alternating current  $I$ ?


A.  $I = I_0 \sin(5\pi t)$

B.  $I = I_0 \sin\left(\frac{\pi t}{0.0025}\right)$

C.  $I = I_0 \sin\left(\frac{2\pi t}{2.5}\right)$

D.  $I = I_0 \sin(800\pi t)$

Q. The diagram shows how the e.m.f. of a simple generator varies with time. What is the frequency and the maximum value of the e.m.f.?



	Frequency / Hz	Maximum e.m.f. / V
A.	200	2.0
B.	200	4.0
C.	400	2.0
D.	400	4.0

Q.3 An electric generator is a device that transforms \_\_\_\_\_ energy into electrical energy

- A. Sound
- B. Heat
- C. Mechanical
- D. Magnetic

Q.19 The output of an a.c. generator has a

- A. Sinusoidal shape
- B. Triangular shape
- C. Square shape
- D. Straight line shape


Q.20 The principle of an electric generator is based on

- A. Faraday's law
- B. Ampere's law
- C. Coulomb's law
- D. Lenz's law

Q.21 The armature of a generator consists of a flat square coil of side 4 cm and 200 turns. The coil rotates in a magnetic field of 0.75T. The angular speed so that a maximum emf of 1.6V is generated is \_\_\_\_\_

- A.  $\frac{20}{3} \text{ rads}^{-1}$
- B.  $\frac{20}{3} \text{ rpm}$
- C.  $\frac{20}{3} \text{ rotations/s}$
- D. None

Q.22 The graph shows the variation with time t of a low-frequency alternating voltage V.



Which expression is a representation of this voltage?

- A.  $0.5\sin(0.4\pi t)$
- B.  $1.0\sin(0.2\pi t)$
- C.  $0.5\sin(10\pi t)$
- D.  $1.0\sin(10\pi t)$

Q.23 Which quantity is increased in step-down transformer?

- A. Current
- B. Power
- C. Voltage
- D. Frequency

Q.24 Which of the remain constant in a transformer?

- A. Current
- B. Potential
- C. Power
- D. Frequency

Q.25 The voltage in the primary and the secondary coils of a step-up transformer are 200 V and 4 kV respectively. If the current in the primary is 1 ampere, then the current in the secondary coil will be

- A. 50 mA
- B. 500 mA
- C. 5A
- D. 5 mA

Q.26 The turn ratio of a transformer is 2:3. If the current through primary is 3A, then current through load resistance is

- A. 1A
- B. 4.5 A
- C. 2A
- D. 1.5 A

Q.27 A transformer is used to

- A. Convert alternating current to direct current
- B. Convert mechanical energy to electrical energy
- C. Convert direct current to alternating current
- D. Change the level of alternating voltage

Q.28 A transformer is used to light a 100 W and 110 V lamp from 220 V mains. If the main current is 0.5 A, the efficiency of the transformer is approximately.

- A. 10%
- B. 30%
- C. 50%
- D. 90%

Q.29 If the current in the primary coil and number of turns in it are  $I_p$  and  $N_p$  respectively and the number of turns and current in the secondary are  $N_s$  and  $I_s$  respectively then the value the of  $N_s : N_p$

- A.  $I_s : I_p$
- B.  $I_s^2 : I_p^2$
- C.  $I_p : I_s$
- D.  $I_p^2 : I_s^2$

Q.30 The ratio of emf across primary coil to the emf across secondary coil is

- A.  $\frac{N_s}{N_p}$
- B.  $\frac{I_s}{I_p}$
- C.  $\frac{I_s^2}{I_p^2}$
- D. None of these

Q.31 In step up transformer, voltage in the secondary increases and power in secondary

- A. Remain same
- B. Increases because current decreases
- C. Decreases because voltage increases
- D. May increases if voltage remain same

Q.32 In a transformer heat is produced due to eddy current in

- A. Primary coil
- B. Secondary coil
- C. Iron core
- D. All of these

Q.33 \_\_\_\_\_ is the energy expended to magnetize and demagnetize the core material in each cycle of the A.C.

- A. Power loss
- B. Heat loss
- C. Hysteresis loss
- D. All of the above

Q.34 Laminated core in a transformer is used to reduce

- A. Eddy current losses
- B. Hysteresis losses
- C. Iron losses
- D. Heat losses due to resistance

pic-9

- Q.35 The loss of power in transformer is due to  
 A. Eddy current  
 B. Magnetic hysteresis  
 C. Resistance of coils  
 D. All
- Q.36 A step-down transformer, transforms 220 volts to 11 volts. If the current in primary and secondary coil are 5A and 90 A respectively, efficiency of transformer is  
 A. 70%  
 B. 20%  
 C. 40%  
 D. 90%
- Q.37 If turns in primary = 50, secondary = 200, primary voltage = 120V, primary current 3A, then output power will be about  
 A. 360 W  
 B. 460 W  
 C. 300 W  
 D. Zero
- Q.38 To improve efficiency of transformer the flux coupling between primary & secondary coils should be  
 A. Small  
 B. Maximum  
 C. May be small or may be maximum  
 D. Zero
- Q.39 Primary and secondary coils of a transformer have 50 and 200 turns respectively. When primary is connected to 9-volt battery secondary voltage is  
 A. 90  
 B. 18  
 C. 36  
 D. Zero
- Q.40 The ratio of secondary to the primary turns in a transformer is 3 : 2. If the power output be  $P$ , then the input power neglecting all losses must be equal to  
 A.  $5P$   
 B.  $1.5P$   
 C.  $P$   
 D.  $\frac{2}{5}P$
- Q.41 The efficiency of transformer is very high because  
 A. There is no moving part in a transformer  
 B. It produces very high voltage  
 C. It produces very low voltage  
 D. None of the above
- Q.42 A step-down transformer is used on a 1000 V line to deliver 20 A at 120 V at the secondary coil. If the efficiency of the transformer is 80% the current drawn from the line is.  
 A. 3 A  
 B. 30 A  
 C. 0.3 A  
 D. 2.4 A
- Q.43 A step-up transformer  
 A. Increases power-level  
 B. Decreases current-level  
 C. Increases voltage-level  
 D. Both B and C
- Q.44 The number of turns in the primary and secondary coil of a step-up transformer are 200 and 500 respectively. If the power in the input is 100 Watt and current 1A then the output power and current will respectively  
 A. 100 W, 2 A  
 B. 200 W, 0.2 A  
 C. 400 W, 4 A  
 D. 100 W, 0.4 A

PAST PAPER MCQs

- Q.45 A transformer changes 12V to 18000 V and there are 6000 turns in the secondary coil. The number of turns in the primary coil are: (ETEA 2009)  
 A. 40  
 B. 20  
 C. 30  
 D. 4
- Q.46 The phenomenon of mutual induction is practically used is (ETEA 2016)  
 A. Transformer  
 B. Generator  
 C. Galvanometer  
 D. Avometer

TOPIC

- Q.47 The phenomenon used for producing emf in coil of generator is;  
 A. Mutual induction  
 B. Self-induction  
 C. Electrostatic induction  
 D. Electromagnetic inductions (ETEA 2017)
- Q.48 The function of main transformer is to convert:  
 A. One direct voltage to another direct voltage of different magnitude  
 B. One alternating voltage to another alternating voltage of different magnitude  
 C. A high value alternating voltage to low value direct voltage  
 D. A high value alternating current to low value direct voltage (ETEA 2018)
- Q.49 In transmission from grid station, power losses are minimized by:  
 A. Increasing current  
 B. Decreasing current  
 C. Increasing resistance  
 D. Increasing voltage (NMDCAT 2020)
- Q.50 The domestic electricity supply has a frequency of:  
 A. 150 Hz  
 B. 100 Hz  
 C. 50 Hz  
 D. 25 Hz (NMDCAT 2020)
- Q.51 Transformer is a device which steps up or steps down the input:  
 A. Current  
 B. Voltage  
 C. Energy  
 D. Power (NMDCAT 2020)
- Q.52 If a stationary bar magnet is placed near a coil at rest so maximum lines of force passes through the coil, the galvanometer shows:  
 A. Maximum current  
 B. Minimum current  
 C. No current  
 D. Intermediate value of current (NMDCAT 2020)
- Q.53 Which is the correct relation:  
 A.  $N_s/N_p = V_s/V_p$   
 B.  $N_s N_p = V_s/V_p$   
 C.  $N_s/N_p = I_s/I_p$   
 D. None (NMDCAT 2021)
- Q.54 In a step-up transformer, current across secondary is \_\_\_\_\_  
 A. More  
 B. Less  
 C. Equal  
 D. None (NMDCAT 2021)
- Q.55 Find Turn ratio if  $V_s/V_p=2$   
 A. 2 : 1  
 B. 1 : 2  
 C. 4 : 8  
 D. None (NMDCAT 2021)

ANSWER KEY

TOPIC-WISE MCQs & PAST PAPER MCQs


1	B	12	C	23	A	34	A	45	D
2	A	13	C	24	D	35	D	46	A
3	A	14	B	25	A	36	D	47	D
4	C	15	B	26	B	37	A	48	B
5	A	16	D	27	D	38	B	49	D
6	A	17	A	28	D	39	D	50	C
7	B	18	C	29	C	40	C	51	B
8	C	19	A	30	B	41	A	52	C
9	D	20	A	31	A	42	A	53	A
10	A	21	A	32	C	43	D	54	B
11	A	22	C	33	C	44	D	55	A

# EXPLANATORY NOTES

1. Primary power =  $220 \times 6 = 1320 \text{ W}$   
 Secondary power =  $11 \times 100 = 1100 \text{ W}$   
 Efficiency =  $\frac{\text{secondary power}}{\text{Primary power}} = \frac{1100}{1320} = 0.833$

2.  $\frac{N_s}{N_p} = \frac{I_p}{I_s}$   
 $\frac{1230}{410} = \frac{6}{I_s}$   
 $I_s = \frac{6 \times 410}{1230} = 2 \text{ A}$

3. Holding a magnet inside solenoid will not change magnetic flux. Hence no emf is generated.  
 4. Resistance of coil has no effect on emf.  
 5. Induce current will produce when there is relative motion between magnet and coil.  
 6. (i) A magnet falls with S-pole along the axis of ring. So according to Lenz's law S-pole form along the face of magnet. As shown in Fig. the current clock wise.  
 (ii) The ring is open. So, no current induced in the ring i.e.  $I = 0$   
 7. It is the statement of Lenz's law.  
 8. According to Lenz's law, the current induced in the ring oppose the cause that produce it




9. All options are correct by using Lenz's law.

10.  $\epsilon = -N \frac{\Delta\phi}{\Delta t} \Rightarrow \epsilon \propto \frac{\Delta\phi}{\Delta t}$

11.  $\epsilon = N \left( \frac{\Delta B}{\Delta t} \right) \cdot A \cos\theta = 500 \times 1 \times (10 \times 10^{-2})^2 \cos 0^\circ = 5 \text{ V}$

12.  $\epsilon = \frac{\Delta\phi}{\Delta t} \Rightarrow IR = \frac{\Delta\phi}{\Delta t} \Rightarrow \frac{Q}{\Delta t} = \frac{\Delta\phi}{R\Delta t}$   
 $\Rightarrow Q = \frac{\Delta\phi}{R} = \frac{10-2}{2} = 4 \text{ C}$

13. (i) A magnet falls with S-pole along the axis of ring. So according to Lenz's law S-pole form along the face of magnet. As shown in Fig. the current clock wise.  
 (ii) Due to repulsion between magnet and ring, the acceleration of magnet is less than "g".



14. Lenz's law is a manifestation of the conservation of energy.  
 As,  $\epsilon = N\omega AB \sin\theta$

15. For maximum induced emf  $\theta = 90^\circ$   
 $\epsilon = N\omega AB$

$I = I_0 \sin \omega t = I_0 \sin(2\pi f t)$

16. From the graph, the period T is 2.5 ms.

$\therefore$  Frequency,  $f = \frac{1}{2.5 \times 10^{-3}} = 400 \text{ Hz}$

$\therefore I = I_0 \sin(2\pi(400)t) = I_0 \sin(800\pi t)$

17. From graph

$V_0 = 2 \text{ V}$

$T = 5 \text{ ms}$


$= 5 \times 10^{-3} \text{ sec}$

As

$f = \frac{1}{T} = \frac{1}{5 \times 10^{-3}}$

$= \frac{10^3}{5} = \frac{1000}{5}$

$= 200 \text{ Hz}$



18. Generator is a device that converts mechanical energy into electrical energy.  
 19. Output of A.C generator is a sine wave.

20. Electric generator work on the principal of Faraday's law

21.  $\epsilon = N\omega AB$

$\epsilon_{\max} = N\omega AB \Rightarrow \omega = \frac{\epsilon}{NAB} = \frac{1.6}{(200) \left( \frac{4 \times 4}{100 \times 100} \right) (0.75)} = \frac{20}{3} \text{ rad / s}$

22.  $V = V_0 \sin\left(\frac{2\pi}{T}t\right) = 0.5 \sin\left(\frac{2\pi}{0.2}t\right) = 0.5 \sin(10\pi t)$

23. Transformer is a device which is use to increase or decrease the alternating voltage.

$P = VI \Rightarrow P = \text{same} \Rightarrow V \propto \frac{1}{I}$

24. As,  $\frac{N_s}{N_p} = \frac{V_s}{V_p} = \frac{I_p}{I_s}$

From relation current and voltage changes. Only for ideal transformer power remain same otherwise power also changes. But frequency in transformer does not change.



25. As,  $\frac{V_s}{V_p} = \frac{I_p}{I_s}$

$$\frac{4 \times 10^4}{200} = \frac{1}{I_s}$$

$$I_s = \frac{200}{4 \times 10^4} = 50 \times 10^{-4} \text{ A} = 50 \text{ mA}$$

26. As,  $\frac{N_s}{N_p} = \frac{I_p}{I_s}$

$$\frac{2}{3} = \frac{3}{I_s} \Rightarrow I_s = \frac{9}{2} = 4.5 \text{ A}$$

27. Because transformer only works with alternating voltage and it is used increase or decrease the alternating voltages.

28. efficiency =  $\frac{P_{\text{output}}}{P_{\text{input}}} \times 100$

$$= \frac{100}{V_p I_p} \times 100 = \frac{100}{220 \times 0.5} \times 100$$

$$= \frac{100}{110} \times 100 = 90\%$$

29. As,  $\frac{N_s}{N_p} = \frac{V_s}{V_p} = \frac{I_p}{I_s}$

30. As,  $\frac{V_s}{V_p} = \frac{I_p}{I_s} \Rightarrow \frac{V_p}{V_s} = \frac{I_s}{I_p}$

31. In an ideal transformer power remain same.

32. The magnetic flux changes through solid conductor (core material), induce current (i.e eddy current) are setup in closed path in the body of conductor. It results in power dissipation and heating of core material.

33. Hysteresis loss

34. The insulation between lamination sheets should be perfect so as to stop the flow of eddy currents.

35. (i) The eddy current results in power dissipation and heating of the core material.

(ii) Hysteresis loss (magnetic hysteresis) is the energy spent to magnetize and demagnetize the core material

(iii) The power also loss in transformer due to resistance of coil.

36.  $\eta = \frac{P_{\text{output}}}{P_{\text{input}}} \times 100\% = \frac{V_s I_s}{V_p I_p} = \frac{11(90)}{220(5)} = \frac{990}{1100} = 0.9$

$$\% \eta = 0.9 \times 100\% = 90\%$$

$$P_o = P_i = I_1 V_1 = 120 \times 3 = 360 \text{ W}$$

37. Flux coupling between primary and secondary coils must be maximum.

38. Voltage by a battery is D.C but transformer operates on A.C

39. Transformer is a device use to step up or down alternating current and voltage keeping power

40. constant ideally.

41. A machine having moving parts reduce its efficiency due to frictional losses but in transformer there is no moving parts.

42.  $\eta = \frac{P_o}{P_i} = \frac{I_s V_s}{I_p V_p} = 0.8$

$$I_p = \frac{I_s V_s}{V_p (0.8)} = \frac{(20)(120)}{(1000)(0.8)} = 3 \text{ A}$$

43. Step-up transformer increase voltage level hence current level decreases.

44. Power input = power output for an ideal transformer

For current  $\frac{I_s}{I_p} = \frac{N_p}{N_s}$

45.  $\frac{V_s}{V_p} = \frac{N_s}{N_p} \Rightarrow N_p = \frac{N_s}{V_s} \times V_p$

46. Principle of transformer.

47. Principle of ac generator.

48. The function of main transformer is to convert one alternating voltage to another alternating voltage of different magnitude.

49.  $P = VI$

If I decrease and voltage increase

$$\text{So } P = I^2 R t$$

Power losses will be minimum.

50. Frequency is used in Pakistan is 50 Hz.

51. Basically, transformer is used to step up or step-down voltage.

52. Induced emf and induced current flows if there is rate of change of magnetic flux.


$$EMF = -N \frac{\Delta \phi}{\Delta t}$$

53.  $\frac{N_s}{N_p} = \frac{V_s}{V_p}$


54. In step up transformer voltage increases and current decreases but power remains same.

55.  $\frac{N_s}{N_p} = \frac{V_s}{V_p} = 1$

- Q.1 The applied input a.c. power to a half-wave rectifier is 100 watts. The d.c. output power obtained is 40 watts. What is the rectification efficiency?  
 A. 50% B. 80%  
 C. 40% D. 95%
- Q.2 The ripple factor indicates the number of ripples in the  
 A. A.C. output B. D.C. output  
 C. A.C. input D. None of these
- Q.3 For a half wave or full wave rectifier the Peak Inverse Voltage of the rectifier is always  
 A. Greater than the input voltage  
 B. Smaller than the input voltage  
 C. Equal to the input voltage  
 D. Greater than the input voltage for full wave rectifier and smaller for the half wave rectifier
- Q.4 The device which converts A.C into D.C is called  
 A. Oscillator B. Transducer  
 C. Rectifier D. Diode
- Q.5 The types of rectifications are  
 A. 3 B. 4  
 C. 5 D. 2
- Q.6 Rectification is possible by  
 A. Transistor B. Diode  
 C. Amplifier D. Capacitor
- Q.7 The semiconductor diode can be used as a rectifier because \_\_\_\_\_  
 A. It has low resistance to the current flow when forward biased & high resistance when reverse biased  
 B. It has low resistance to the current flow when forward biased  
 C. It has high resistance to the current flow when reverse biased  
 D. Its conductivity increases with rise of temperature
- Q.8 In the half-wave rectifier circuit shown, Which one of the following wave forms is true for diode, the output across C and D?




- Q.9 For the given circuit of PN-junction diode, which of the following statement is correct




- A. In forward biasing the voltage across R is 2V  
 B. In reverse biasing the voltage across R is 2V  
 C. In forward biasing the voltage across R is V  
 D. In reverse biasing the voltage across R is V
- Q.10 The simplest type of rectification known as half wave rectification is obtained by  
 A. Using a transistor  
 B. Suppressing half wave of A.C supply by using diode  
 C. Suppressing the harmonics in A.C voltage  
 D. Using a Coolidge tube
- Q.11 Output of half wave rectifier is suitable only  
 A. To operate radio B. Charging batteries  
 C. For running a D.C motor D. All of these
- Q.12 During the interval  $0 \rightarrow \frac{T}{2}$  the forward biased diode offers  
 A. Very small resistance B. Very high resistance  
 C. Very small current flow through it D. Zero resistance
- Q.13 In a half wave rectifier, the frequency of the input is N, the frequency and form of the output will be  
 A. N/2 and Pulsating B. 2N and steady  
 C. N and Pulsating D. N and continuous
- Q.14 The most common device used as filter is  
 A. Capacitor B. Transformer  
 C. Resistor D. Transistor
- Q.15 The method by which only one half of A.C cycle is converted into direct current is called  
 A. Half wave amplification B. Full wave rectification  
 C. Half wave rectification, D. Full wave amplification
- Q.16 If time period of input T in the full wave bridge rectifier circuit, then time period of the pulsating output of the circuit will be  
 A. T B. T/2  
 C. 2T D. T/4
- Q.17 In a bridge rectifier how, many diodes conduct during each half cycle of input A.C  
 A. 2 B. 1  
 C. 3 D. All
- Q.18 If a full wave rectifier circuit is operating from 50 Hz mains, the fundamental frequency in the ripple will be  
 A. 50 Hz B. 70.7 Hz  
 C. 100 Hz D. 25 Hz

Q.19 In figure the input is across the terminals A and C and the output is across B and D. Then the output is




- A. Half wave rectified
  - B. Full wave rectified
  - C. Zero
  - D. Same as input
- Q.20 To reduce ripples in the output of bridge rectifier we should use
- A. Diodes having low forward resistance
  - B. Low frequency A.C
  - C. Diodes having high forward resistance
  - D. A filter circuit
- Q.21 In full wave rectification, the output D.C. voltage across the load is obtained for
- A. The positive half cycle of input A.C
  - B. The complete cycle of input A.C
  - C. The negative half cycle of input A.C
  - D. All of the above

Q.22 In the following figure



- A. D<sub>1</sub> and D<sub>3</sub> conducts simultaneously
  - B. D<sub>1</sub> and D<sub>2</sub> conducts alternatively
  - C. D<sub>1</sub> and D<sub>2</sub> conducts simultaneously
  - D. Both A and C
- Q.23 Output voltage of rectifier is not smooth. It can be made smooth by using a circuit known as:
- A. Wheat stone circuit
  - B. Ripple circuit
  - C. Bridge circuit
  - D. Filter circuit
- Q.24 In the diagram, diodes are arranged for the full wave rectification where input alternating voltage must be applied



- A. across A and B
  - B. Across B and D
  - C. Across A and C
  - D. Across B and C
- Q.25 The basic reason why a full wave rectifier has a twice the efficiency of a half wave rectifier is that
- A. It make use of transformer
  - B. It utilizes both half cycle of the input
  - C. Its ripple factor is much less
  - D. Its output frequency is double the frequency

- Q.26 If an A.C voltage of rms value of 10V is applied as input of half wave rectifier, then the rms voltage value of D.C output will be
- A. 10 V
  - B. 10.3 V
  - C. 10.7 V
  - D. 9.3 V
- Q.27 If a half-wave rectifier is used to convert 50Hz A.C into D.C then the number of pulses present in rectifier voltage is:
- A. 25
  - B. 100
  - C. 50
  - D. 5
- Q.28 The output from a full wave rectifier is
- A. An ac voltage
  - B. Uni-directional
  - C. Pulsating dc
  - D. Both B and C

PAST PAPER MCQs

- Q.29 Conversion of alternating current into direct current is called (ETEA 2012)
- A. Rectification
  - B. Amplification
  - C. Regeneration
  - D. Oscillation
- Q.30 Output voltage of rectifier is not smooth. It can be made smooth by using a circuit known as: (MDCAT 2017)
- A. Wheat stone circuit
  - B. Ripple circuit
  - C. Bridge circuit
  - D. Filter circuit
- Q.31 In the following figure what happens for the positive half cycle of the input? (MDCAT 2017)
- 
- A. D<sub>1</sub> and D<sub>3</sub> conducts
  - B. D<sub>4</sub> and D<sub>2</sub> conducts
  - C. D<sub>1</sub> and D<sub>2</sub> conducts
  - D. D<sub>4</sub> and D<sub>1</sub> conducts
- Q.32 In case of half wave rectification, resistance of diode during negative half of A.C is: (MDCAT 2018)
- A. Very high
  - B. A few ohms
  - C. Very low
  - D. Negligible
- Q.33 The direction of current through the load resistance of a full-wave rectification circuit: (MDCAT 2018)
- A. Inverts for negative cycle
  - B. Inverts for positive cycle
  - C. Changes for every cycle
  - D. Remains constant
- Q.34 In full wave rectification, the diodes are used (NMDCAT 2020)
- A. 1
  - B. 2
  - C. 3
  - D. 4
- Q.35 Which has less ripple factor (NMDCAT 2021)
- A. Half wave rectifier
  - B. Full wave rectifier
  - C. Both
  - D. None

- Q.36 Ripple factor of full wave rectifier?  
 A. 0.482  
 C. 0.21  
 B. 1.21  
 D. 0.84
- Q.37 If  $f = 50\text{Hz}$  then in half wave no of pulses are  
 A. 50  
 C. 100  
 B. 60  
 D. 150

# ANSWER KEY

## TOPIC-WISE MCQs & PAST PAPER MCQs

1	C	6	B	11	B	16	B	21	B	26	D	31	B	36	A
2	B	7	A	12	A	17	A	22	D	27	C	32	A	37	A
3	B	8	D	13	C	18	C	23	D	28	D	33	D		
4	C	9	C	14	A	19	B	24	C	29	A	34	D		
5	D	10	B	15	C	20	D	25	B	30	D	35	B		

# EXPLANATORY NOTES

- Rectification efficiency =  $\frac{\text{d.c. output power}}{\text{a.c. input power}} = \frac{40}{100} = 0.4 = 40\%$
- The ripple factor indicates the number of ripples in the D.C. output.
- The peak input voltage is smaller than the input voltage due to the presence of diode(s). A single diode reduces the output voltage by approximately 0.7V.
- Definition of rectifier.
- Two types of rectification  
 i) Half wave rectification  
 ii) Full wave rectification
- Diode is used as rectifier.
- Its reverse biased resistance is very high in order of  $M \Omega$  and forward biased resistance is very low in order of few ohms.
- Half wave rectifier, rectifies only the half cycle of input ac signal and it blocks the other half
- In forward biasing, resistance of PN junction diode is zero, so whole voltage appears across the resistance.
- PN junction has low resistance in one direction of potential difference  $+V$ , so a large current flows (forward biasing). It has a high resistance in the opposite potential difference direction  $V$ , so a very small current flows (Reverse biasing)
- Output voltage of half wave rectifier is pulsating, when it made smooth by filter, voltage level is decrease this low voltage suitable for charging batteries.
- Forward biased diodes has low resistance.
- In half wave rectifier
  - Input frequency = output frequency
  - Output wave is pulsating
- Filters made output smooth.
- In half wave rectification only one half of AC is converted into DC.
- As in the full wave rectification  $f_o = 2f_m \Rightarrow T_o = \frac{T}{2}$
- In a bridge rectifier two diodes in forward biased and conduct in each half cycle of input A.C
- In full wave rectifier, frequency of output wave is double of input wave.
- It is the diagram of full wave bridge rectifier circuit.
- We use capacitor as filter circuit.
- Direction of current flow through the load resistance R is the same in both halves cycles of input wave.
- Full wave rectifier circuit opposite diodes work simultaneously
- Output voltage of rectifier is not smooth. It can be made smooth by using a circuit known as filter circuit.
- In full wave bridge rectification circuit, input must be applied at the terminals where opposite ends of two diodes are joined
- It allows both half cycles so its efficiency twice.

26. In general potential barrier of Si is 0.7V  
So, d.c output =  $10 - 0.7 = 9.3$  V
27. For half wave rectifier.  $f' = f \Rightarrow f' = 50\text{Hz}$
28. Rectification is process in which a.c converts into pulsative d.c.
29. Definition of rectification.
30. Output voltage of rectifier is not smooth. It can be made smooth by using a circuit known as filter circuit.
31. During the positive half-cycle the diodes  $D_2$  and  $D_4$  are forward biased and diodes  $D_1$  and  $D_3$  are reversed biased.
32. In case of half rectification, resistance of diode during negative half of A.C is very high.
33. In full wave rectification circuit, direction of current through load resistance remains same during +ve & -ve half cycles.
34. Four diode are used in FWR
35. Ripple factor of H.W.R = 1.21  
Ripple factor of F.W.R = 0.48
36. Form factor  $\frac{I_{rms}}{I_{DC}} = \frac{I_o / \sqrt{2}}{2I_o} = \frac{\pi}{2\sqrt{2}} = 1.11$   
 $\sqrt{R.F} = \sqrt{(F.F)^2 - 1} = \sqrt{(1.1)^2 - 1} = 0.48$
37. In half wave rectification  $f_{output} = f_{input}$

# 11 TOPIC

## DAWN OF MODERN PHYSICS PRACTICE EXERCISE

- Q.1 Threshold wavelength for a metal having work function  $W_0$  is  $X$ . What is the threshold wavelength for the metal having work function  $2W_0$ ?  
A.  $4\lambda$   
C.  $\lambda/2$   
B.  $2\lambda$   
D.  $\lambda/4$
- Q.2 An electric bulb of 100 W converts 3 % of electrical energy into light energy. If the wavelength of light emitted is 6625 Å, the number of photons emitted in 1 s is \_\_\_\_\_.  
( $h=6.625 \times 10^{-34}$  J.sec)  
A.  $10^{17}$   
C.  $10^{21}$   
B.  $10^{19}$   
D.  $10^{15}$
- Q.3 The wavelength of a photon needed to remove a proton from a nucleus which is bound to the nucleus with 1 MeV energy is nearly  
A. 1.2 nm  
C.  $1.2 \times 10^{-6}$  nm  
B.  $1.2 \times 10^{-3}$  nm  
D.  $1.2 \times 10^4$  nm
- Q.4 If K.E. of free electron is doubled, its de Broglie wavelength become  
A.  $\frac{1}{\sqrt{8}}$   
C.  $\frac{1}{\sqrt{2}}$   
B.  $\sqrt{2}$   
D. 2
- Q.5 In a photon-particle collision, the quantity that does not remain conserved is  
A. Total energy  
C. Number of photons  
B. Total momentum  
D. None of these
- Q.6 The momentum of a photon is  $2 \times 10^{-16}$  gm-cm/sec. Its energy is  
A.  $0.61 \times 10^{-26}$  erg  
C.  $6 \times 10^{-6}$  erg  
B.  $2.0 \times 10^{-26}$  erg  
D.  $6 \times 10^{-8}$  erg
- Q.7 Ratio of momentum of photons having wavelength 4000 angstrom and 8000 angstroms is  
A. 2 : 1  
C. 20 : 1  
B. 1 : 2  
D. 1 : 20
- Q.8 A radio station emits 10 kW power of 90.8 MHz. Find the number of photons emitted per second  
A.  $1.6 \times 10^{28}$   
C.  $1.6 \times 10^{30}$   
B.  $1.6 \times 10^{29}$   
D.  $1.6 \times 10^{32}$
- Q.9 The energy of a photon is  $3 \times 10^{-19}$  J. Its momentum is  
A.  $10^{-27}$  kg ms<sup>-1</sup>  
C.  $10^{-11}$  kg ms<sup>-1</sup>  
B.  $9 \times 10^{-11}$  kg ms<sup>-1</sup>  
D.  $3 \times 10^{-7}$  kg ms<sup>-1</sup>
- Q.10 The mass of a photon at rest is  
A. 1 a.m.u.  
C.  $1.67 \times 10^{-35}$  kg  
B.  $9 \times 10^{-31}$  kg  
D. Zero


Topic - 11

Topic - 11

- Q.11 The momentum of a photon is  $p$ . The frequency associated with it is given by  
 A.  $pc/h$  B.  $h/p$   
 C.  $ph/c$  D.  $h/pc$
- Q.12 Photon A has twice the energy of photon B. What is the ratio of the momentum of A to that of B?  
 A. 2 : 1 B. 1 : 2  
 C. 1 : 1 D. None of these
- Q.13 The value and units of the Planck's constant 'h' can be expressed as:  
 A.  $6.63 \times 10^{-34} \text{ Js}^{-1}$  B.  $6.63 \times 10^{-34} \text{ Js}$   
 C.  $6.63 \times 10^{-42} \text{ Js}$  D.  $3.63 \times 10^{-34} \text{ Js}$
- Q.14 Let  $n_r$  and  $n_b$  be respectively the number of photons emitted by a red bulb and a blue bulb of equal power in a given time. Then  
 A.  $n_r = n_b$   
 B.  $n_r < n_b$   
 C.  $n_r > n_b$   
 D. The information is insufficient to get a relation between  $n_r$  and  $n_b$

- Q.15  $\lambda$  is proportional to  
 A.  $\frac{1}{E}$  for both photons and particles B.  $\frac{1}{\sqrt{E}}$  for both photons and particles  
 C.  $\frac{1}{E}$  for photons and  $\frac{1}{\sqrt{E}}$  for particles D.  $\frac{1}{\sqrt{E}}$  for photons and  $\frac{1}{E}$  for particles

Q.16 Which of the following graphs correctly represents the variation of particle momentum with associated de-Broglie wavelength?



- Q.17 A material particle with a rest mass  $m_0$  is moving with speed of light  $c$ . The associated de-Broglie wavelength is given by  
 A.  $\frac{h}{m_0 c}$  B.  $\frac{m_0 c}{h}$   
 C. 0 D.  $\infty$
- Q.18 A photon is considered to have:  
 A. Energy B. Wavelength and frequency  
 C. Momentum D. All of these
- Q.19 If electron and proton have same De-Broglie wavelength, which have greater speed  
 A. Electron B. Proton  
 C. Both have same D. Electron and Proton can't have wavelength
- Q.20 The wavelength of a moving particle is inversely proportional to  
 A. Mass B. Velocity  
 C. Energy D. Momentum

- Q.21 According to De-Broglie, an electron can be regarded as:  
 A. Particle only B. Particle and wave both  
 C. Are negligible D. None of these
- Q.22 Davisson determine the wavelength of scattered electron from the relation:  
 A.  $\lambda = \frac{h}{2mVe}$  B.  $\lambda = \frac{2h}{\sqrt{mVe}}$   
 C.  $\lambda = \frac{h}{2\sqrt{2mVe}}$  D.  $\lambda = \frac{h}{\sqrt{2mVe}}$
- Q.23 If an electron is accelerated through a potential difference of 54 volts, its de-Broglie wavelength will be:  
 A.  $1.66 \times 10^{-8} \text{ m}$  B.  $1.66 \times 10^{-10} \text{ m}$   
 C.  $1.66 \times 10^{-9} \text{ m}$  D.  $1.66 \times 10^{-12} \text{ m}$
- Q.24 In Davisson and Germer experiment, target crystal is made up of  
 A. Copper B. Nickel  
 C. Aluminium D. Silver
- Q.25 A body of mass 200 g moves at the speed of 5 m/hr. So de-Broglie wavelength related to it is of the order ( $h=6.26 \times 10^{-34} \text{ Js}$ )  
 A.  $10^{-10} \text{ m}$  B.  $10^{-20} \text{ m}$   
 C.  $10^{-30} \text{ m}$  D.  $10^{-40} \text{ m}$
- Q.26 In Davisson-Germer experiment, the diffracted electron beam from crystal shows  
 A. Particle property B. Light property  
 C. Wave property D. Quantum property
- Q.27 The electron, accelerated by a potential difference  $V$  has de-Broglie wavelength  $\lambda$ . If the electron is accelerated by a p.d  $4V$ , its de-Broglie wavelength will be  
 A.  $2\lambda$  B.  $\frac{\lambda}{2}$   
 C.  $4\lambda$  D.  $\frac{\lambda}{4}$
- Q.28 A proton and an  $\alpha$ -particle are accelerated through the same potential difference. The ratio of their de-Broglie wavelengths ( $\lambda_p / \lambda_\alpha$ ) is  
 A. 1 B.  $2\sqrt{2}$   
 C. 2 D.  $1\sqrt{2}$
- Q.29 The ratio of momenta of an electron and an  $\alpha$ -particle which are accelerated from rest by a potential difference of 100 V is  
 A. 1 B.  $\sqrt{\frac{m_e}{m_\alpha}}$   
 C.  $\sqrt{\frac{2m_e}{m_\alpha}}$  D.  $\sqrt{\frac{m_e}{2m_\alpha}}$
- Q.30 What is the de Broglie wavelength of a proton whose linear momentum has a magnitude of  $3.3 \times 10^{-23} \text{ kg m/s}$ ?  
 A. 0.0002 nm B. 0.002 nm  
 C. 0.02 nm D. 0.2 nm

Topic - 11

- Q.31 The velocity of a particle of mass  $m$  of de-Broglie wavelength  $\lambda$  is \_\_\_\_\_  
 A.  $\frac{2h}{m\lambda}$  B.  $\frac{m\lambda c^2}{h}$   
 C.  $2m\lambda c^2$  D.  $h/m\lambda$
- Q.32 If  $E_1$ ,  $E_2$  and  $E_3$  are the respective kinetic energies of an electron, an alpha particle and a proton, each having the same de-Broglie wavelength, then  
 A.  $E_1 > E_3 > E_2$  B.  $E_1 > E_2 > E_3$   
 C.  $E_2 > E_3 > E_1$  D.  $E_1 = E_2 = E_3$
- Q.33 If the K.E of a free electron doubles then its de-Broglie wavelength changes by a factor  
 A.  $\frac{1}{2}$  B. 2  
 C.  $\sqrt{2}$  D.  $\sqrt{2}$
- Q.34 The magnitude of de-Broglie wavelength ( $\lambda$ ) of electrons (e), proton (p), neutron (n), and  $\alpha$  particles all have the same energy 1 MeV, in increasing order will follow the sequence.  
 A.  $\lambda_e, \lambda_p, \lambda_n, \lambda_\alpha$  B.  $\lambda_e, \lambda_p, \lambda_n, \lambda_\alpha$   
 C.  $\lambda_\alpha, \lambda_n, \lambda_p, \lambda_e$  D.  $\lambda_\alpha, \lambda_p, \lambda_n, \lambda_e$
- Q.35 The wavelength of matter waves is independent of:  
 A. Mass B. Velocity  
 C. Momentum D. Charge
- Q.36 A particle of mass  $M$  at rest decays into two masses  $m_1$  and  $m_2$  with non-zero velocities. The ratio of de-Broglie wave lengths of the particles  $\frac{\lambda_1}{\lambda_2}$  is  
 A.  $\sqrt{\frac{m_2}{m_1}}$  B.  $\frac{m_1}{m_2}$   
 C.  $\frac{\sqrt{m_1}}{\sqrt{m_2}}$  D. 1 : 1
- Q.37 The Davisson and Garmer experiment indicate  
 A. Interference B. Electron diffraction  
 C. Polarization D. Refraction

PAST PAPER MCQs

- Q.38 Select the correct relation between wave and particle nature of radiation? (ETEA 2014)  
 A.  $E = \frac{hc}{\lambda}$  B.  $E = \frac{hc}{c}$   
 C.  $E = \frac{\lambda c}{h}$  D.  $E = h\lambda c$
- Q.39 Which one of the following has the largest energy content? (MCAT 2015)  
 A.  $\gamma$ -rays B. Infrared rays  
 C. X-rays D. Ultra violet radiations

Topic - 11

- Q.40 Choose the correct relationship, when  $E =$  energy,  $h =$  plank's constant,  $c =$  velocity of light,  $f =$  frequency,  $\lambda =$  wavelength: (ETEA 2015)  
 A.  $E = hfc$  B.  $E = \frac{c}{\lambda}$   
 C.  $E = hf$  D.  $E = \frac{n\lambda}{c}$
- Q.41 Which of the following is the best evidence for the wave nature of matter? (ETEA 2015)  
 A. The photoelectric effect  
 B. The Compton effect  
 C. The spectral radiation form cavity radiation  
 D. The reflection of electrons by crystal
- Q.42 The momentum of wave of wavelength  $1.32 \times 10^{-9}$  is: (MDCAT 2017)  
 A.  $5 \times 10^{-26}$  Ns B.  $5 \times 10^{-43}$  Ns  
 C.  $5 \times 10^{-25}$  Ns D.  $5 \times 10^{-44}$  Ns
- Q.43 Calculate the frequency of a photon having a momentum of  $4.42 \times 10^{-26}$  Ns: (MDCAT 2017)  
 A.  $2 \times 10^{14}$  Hz B.  $5 \times 10^{16}$  Hz  
 C.  $2 \times 10^{16}$  Hz D.  $2 \times 10^{18}$  Hz
- Q.44 The De-Broglie wavelength of particle of mass "m" moving with the kinetic energy "E" can be written as:  
 A.  $\sqrt{\frac{h}{2mE}}$  B.  $\frac{h}{2mE}$   
 C.  $\frac{h}{\sqrt{2mE}}$  D.  $\frac{\sqrt{h}}{2mE}$
- Q.45 The de Broglie wave length of an electron travelling with a speed of  $1.0 \times 10^7$  m/s equal to, ( $h = 6.6 \times 10^{-34}$  Js and  $m_e = 9.1 \times 10^{-31}$  kg): (MDCAT 2018)  
 A.  $7.3 \times 10^{11}$  m B.  $7.3 \times 10^{-11}$  m  
 C.  $7.3 \times 10^8$  m D.  $7.3 \times 10^{-13}$  m
- Q.46 A 5-watt LED bulb converts 80% of the power into light photons of wavelength 660 nm. What is the number of photons emitted from the bulb in one second? (MDCAT 2018)  
 A.  $5.8 \times 10^{34}$  C.  $6.6 \times 10^7$   
 B.  $7.5 \times 10^{18}$  D.  $1.3 \times 10^{19}$
- Q.47 The value and units of the Plank constant 'h' can be expressed as: (MDCAT 2019)  
 A.  $6.63 \times 10^{-34}$  Js<sup>-1</sup> B.  $6.63 \times 10^{-34}$  Js  
 C.  $6.63 \times 10^{-43}$  Js D.  $3.63 \times 10^{-34}$  Js
- Q.48 Calculate the energy of a photon of frequency  $3.0 \times 10^{18}$  Hz. ( $h = 6.63 \times 10^{-34}$  Js) (MDCAT 2019)  
 A.  $19.89 \times 10^{-18}$  J B.  $11.89 \times 10^{-16}$  J  
 C.  $1.89 \times 10^{-16}$  J D.  $19.89 \times 10^{-16}$  J
- Q.49 The wavelength associated with an electron is of the order of: (NMDCAT 2020)  
 A. Visible light B. X-Rays  
 C. Radio waves D. Infrared

- Q.50 Which photon carries the most energy?  
 A. Blue B. Violet  
 C. Red D. Green (NMDCAT 2021)
- Q.51 Photon is made up \_\_\_\_\_ quarks  
 A. Basons B. No quark  
 C. Quarks in hadrons D. Masons (NMDCAT 2021)
- Q.52 If clouds were black body \_\_\_\_\_ what would happen?  
 A. Will receive no light on surface B. Some light would reach surface  
 C. Reflects back D. Absorbs light (NMDCAT 2021)
- Q.53 By Wein's constant the wavelength and temperature are related.  
 A. Inversely B. Directly  
 C. Equal D. None (NMDCAT 2021)
- Q.54 If wavelength = 500 nm of a photon, then its frequency is:  
 A.  $5 \times 10^{14}$  Hz B.  $6 \times 10^{14}$  Hz  
 C.  $7 \times 10^{14}$  Hz D.  $8 \times 10^{14}$  Hz (NMDCAT 2021)
- Q.55 Find E in eV "Photon Wavelength was given 1440nm"  
 A. 0.867 eV B. 18.67 eV  
 C. 9.32 eV D. 0.932 eV (NMDCAT 2021)
- Q.56 If velocity becomes 5 times, what happened to De Broglie Wavelength  
 A. Decrease by 5 times B. Increase by 5 times  
 C. Both D. None (NMDCAT 2021)

# ANSWER KEY

## TOPIC-WISE MCQs & PAST PAPER MCQs

1	C	11	A	21	B	31	D	41	B	51	B
2	B	12	A	22	D	32	A	42	C	52	D
3	B	13	B	23	B	33	C	43	C	53	A
4	C	14	C	24	B	34	C	44	C	54	B
5	C	15	C	25	C	35	D	45	B	55	A
6	D	16	D	26	C	36	A	46	D	56	A
7	A	17	C	27	B	37	B	47	C		
8	B	18	D	28	B	38	A	48	D		
9	A	19	A	29	D	39	A	49	B		
10	D	20	D	30	C	40	C	50	B		

# EXPLANATORY NOTES

- Since  $W_o = h \frac{c}{\lambda}$   
 $2W_o = h \frac{c}{\lambda_1} \Rightarrow 2 = \frac{\lambda}{\lambda_1}$  or  $\lambda_1 = \frac{\lambda}{2}$
- $P = 100 \times \frac{3}{100} = 3 \text{ Watt}, t = 1 \text{ sec}, N = ?$   
 $\lambda = 6625 \text{ \AA} = 6625 \times 10^{-10} \text{ m}$   
 $P = \frac{Nh c}{\lambda t}$   
 $N = \frac{P \lambda t}{h c} = \frac{3 \times 6625 \times 10^{-16} \times 1}{6.625 \times 10^{-34} \times 3 \times 10^8}$   
 $N = 10^{19}$
- $E = \frac{h c}{\lambda}, \lambda = \frac{h c}{E} = \frac{1240}{10^6} \text{ nm}$   
 $= 1.24 \times 10^{-3} \text{ nm}$
- Since  $\lambda = \frac{h}{\sqrt{2m K.E}} \Rightarrow \lambda \propto \frac{1}{\sqrt{K.E}}$
- In a photon - particle collision, the number of photons may not be conserved. The photon may be absorbed or a new photon may be created.
- $E = m c \cdot c = p c = (2 \times 10^{-16})(3 \times 10^8) = 6 \times 10^{-8} \text{ erg}$
- $P = \frac{h}{\lambda} \Rightarrow \frac{P_1}{P_2} = \frac{\lambda_2}{\lambda_1} = \frac{8000 \text{ \AA}}{4000 \text{ \AA}} = 2:1$
- $E = n h f \Rightarrow \frac{E}{t} = \frac{n}{t} h f \Rightarrow \frac{n}{t} = \frac{P}{h f} = 1.6 \times 10^{29}$
- $E = P c \Rightarrow P = \frac{E}{c} = \frac{3 \times 10^{-19}}{3 \times 10^8} = 10^{-27} \text{ kg ms}^{-1}$
- The rest mass of photon is zero.
- De-Broglie wavelength  $= \lambda = \frac{h}{p} \Rightarrow \frac{c}{f} = \frac{h}{P} \Rightarrow f = \frac{p c}{h}$ .
- $E = P c \Rightarrow P \propto E, \frac{P_A}{P_B} = \frac{E_A}{E_B} = \frac{2 E_B}{E_B} = 2$
- Plank's constant  $= h = 6.63 \times 10^{-34} \text{ Js}$
- $E n = n h f \Rightarrow n = \frac{E}{h f} = \frac{E \lambda}{h c} \Rightarrow n \propto \lambda$



## Topic - 11

$$15. \text{ For photon } \Rightarrow E = hf = \frac{hc}{\lambda} \Rightarrow E \propto \frac{1}{\lambda}$$

$$\text{ For particle } \Rightarrow \lambda = \frac{h}{\sqrt{2mE}} \Rightarrow \lambda \propto \frac{1}{\sqrt{E}}$$

$$16. P = \frac{h}{\lambda} \Rightarrow P \propto \frac{1}{\lambda}$$

$$17. m = \frac{m_0}{\sqrt{1 - \frac{c^2}{v^2}}} = \infty \quad \lambda = \frac{h}{mv} \Rightarrow \boxed{\lambda = 0}$$

18. Wavelength, frequency, momentum and energy all are associated with photon.

$$19. \lambda = \frac{h}{P} \Rightarrow \lambda = \frac{h}{mv} \Rightarrow v = \frac{h}{m\lambda} \Rightarrow v \propto \frac{1}{\lambda}$$

$$20. \lambda = \frac{h}{P} \Rightarrow \lambda \propto \frac{1}{P}$$

21. According to de-Broglie relation electron regarded as both wave and particle i.e.  $\lambda = \frac{h}{P}$

$$22. \lambda = \frac{h}{P}, \quad P = \sqrt{2meV}$$

$$\text{ So, } \lambda = \frac{h}{\sqrt{2meV}}$$

$$23. \lambda = \frac{h}{\sqrt{2meV}} = \frac{6.63 \times 10^{-34}}{\sqrt{2(9.1 \times 10^{-31})(1.6 \times 10^{-19})(54)}} = 1.66 \times 10^{-10} \text{ m}$$

24. Davisson and Germer showed that electrons are diffracted from nickel crystal in exactly same manner as X-rays.

$$25. m = 200 \text{ g} = 0.2 \text{ kg}, \quad v = 5 \frac{\text{m}}{\text{hr}} = \frac{5}{3600} \frac{\text{m}}{\text{s}}$$

$$p = \frac{h}{\lambda} = mv$$

$$\therefore \lambda = \frac{h}{mv} = \frac{6.626 \times 10^{-34} \times 3600}{0.2 \times 5}$$

$$= 23.85 \times 10^{-31}$$

$$= 2.385 \times 10^{-30} = 10^{-30} \text{ m}$$

26. Diffracted electron beam from crystal shows wave nature.

$$27. \lambda = \frac{h}{\sqrt{2meV}} \Rightarrow \lambda = \frac{1}{\sqrt{V}}$$

$$28. \lambda = \frac{h}{\sqrt{2mqV}} \propto \frac{1}{\sqrt{mq}} \Rightarrow \frac{\lambda_1}{\lambda_2} = \sqrt{\frac{m_2 q_2}{m_1 q_1}} = \sqrt{\frac{4m_e \times 2e}{m_p \times e}} = \sqrt{8} = 2\sqrt{2}$$

$$29. \lambda = \frac{h}{\sqrt{2meV}}$$

$$P = \sqrt{2meV}$$

$$\frac{P_e}{P_a} = \frac{\sqrt{2m_e eV}}{\sqrt{2m_a (2e)V}} \Rightarrow \frac{P_e}{P_a} = \sqrt{\frac{m_e}{2m_a}}$$

$$30. \lambda = \frac{h}{P} \Rightarrow \lambda = \frac{6.63 \times 10^{-34}}{3.3 \times 10^{-23}} = 2 \times 10^{-11} \text{ m} = 0.02 \times 10^{-9} \text{ m} = 0.02 \text{ nm}$$

$$31. \lambda = \frac{h}{P} = \frac{h}{mv} \Rightarrow v = \frac{h}{m\lambda}$$

$$32. \lambda = \frac{h}{\sqrt{2mE}} \Rightarrow E = \frac{h^2}{2m\lambda^2} \Rightarrow E \propto \frac{1}{\lambda^2}$$

$$33. \lambda = \frac{h}{\sqrt{2mk.E}} \Rightarrow \lambda = \frac{1}{\sqrt{K.E}}$$

$$34. \lambda = \frac{h}{\sqrt{2mE}} \Rightarrow \lambda \propto \frac{1}{\sqrt{m}}$$

35. Wavelength of matter wave is independent of charge.

$$36. \lambda = \frac{h}{\sqrt{2meV}} \Rightarrow \lambda = \frac{1}{\sqrt{m}} \Rightarrow \frac{\lambda_1}{\lambda_2} = \sqrt{\frac{m_2}{m_1}}$$

37. Davission and Germer shows diffraction of electron through metal crystal.

$$38. E = hf = \frac{hc}{\lambda}$$

39. Gamma rays have more energy.

40. Energy of photon =  $E = hf$

41. Compton effect shows wave nature of matter.

$$42. p = \frac{h}{\lambda} = \frac{6.63 \times 10^{-34}}{1.32 \times 10^{-9}} = 5 \times 10^{-25} \text{ N s}$$

$$43. \lambda = \frac{h}{p} \Rightarrow \frac{c}{f} = \frac{h}{p} \quad \because c = f\lambda$$

$$\Rightarrow f = \frac{pc}{h} = \frac{4.42 \times 10^{-25} \times 3 \times 10^8}{6.63 \times 10^{-34}} = 2 \times 10^{18} \text{ Hz}$$

$$44. \lambda = \frac{h}{\sqrt{2mVe}}$$

$$Ve = K.E = E$$

$$\lambda = \frac{h}{\sqrt{2mE}}$$

$$45. \lambda = \frac{h}{mc} = \frac{6.63 \times 10^{-34}}{9.1 \times 10^{-31} \times 3 \times 10^8} = 7.3 \times 10^{-11} \text{ m}$$

Topic

$$46. P = \frac{W}{t} = \frac{nhf}{t} \therefore W = E = nhf$$

$$n = \frac{Pt}{hf} = \frac{Pt\lambda}{hc}$$

$$= \frac{80\% (5) \times 1 \times 660 \times 10^{-9}}{6.63 \times 10^{-34} \times 3 \times 10^8} = 1.3 \times 10^{19}$$

$$47. \text{ Value and units of plank's constant is } h = 6.63 \times 10^{-34} \text{ Js.}$$

$$48. E = hf = 6.63 \times 10^{-34} \times 3 \times 10^{15} = 1.989 \times 10^{-15} \text{ J} = 19.89 \times 10^{-16} \text{ J}$$

$$49. \lambda = \frac{h}{mv} = \frac{6.63 \times 10^{-34}}{9.1 \times 10^{-31} \times 10^6} = 10^{-10}$$

$$50. f_v > f_b > f_r$$

$$E = hf$$

$E_{\text{violet}}$  is greater

51. Photon is made of no quarks

52. Clouds absorb all right.

53.

$$\lambda T = \text{constant}$$

$$\lambda = \frac{1}{T}$$

$$54. f = \frac{3 \times 10^8}{500 \times 10^{-9}} = \frac{3}{500} \times 10^{17} = 0.006 \times 10^{17}$$

$$f = 6 \times 10^{14} \text{ Hz}$$

$$55. E = \frac{(6.63 \times 10^{-34})(3 \times 10^8)}{1440 \times 10^{-9}} \text{ J}$$

$$= 0.014 \times 10^{-34+8+9} \text{ J}$$


$$= \frac{1.4 \times 10^{-19}}{1.6 \times 10^{-19}} \text{ eV} = 0.87 \text{ eV}$$

$$56. \lambda = \frac{h}{mv}$$

12  
TOPICATOMIC SPECTRA  
PRACTICE EXERCISE

- Q.1 When a hydrogen atom is in its first excited level, what is the relation of radius and Bohr radius?  
A. Twice  
B. 4 times  
C. Same  
D. Half
- Q.2 What is the energy required to ionize an H-atom from the third excited state, if ground state ionization energy of H-atom is 13.6 eV?  
A. 0.85 eV  
B. 3.4 eV  
C. 13.6 eV  
D. 12.1 eV
- Q.3  $\frac{1}{\lambda_n} = R \left( \frac{1}{p^2} - \frac{1}{n^2} \right)$  in this relation  $p = 1, 2, 3, \dots$  and  $n$  will be  
A.  $p, p+1, p+2, \dots$   
B.  $p+1, p+2, \dots$   
C.  $p, p-1, p-2, \dots$   
D.  $p-1, p-2, \dots$
- Q.4 Lyman series of H-atoms lie in the \_\_\_\_\_ region of electromagnetic spectrum  
A. visible  
B. ultraviolet  
C. infrared  
D. red
- Q.5 The Balmer series for hydrogen atom corresponds to electronic transitions that terminate in the state of quantum number  $n = 2$ . The longest wavelength of photon emitted is  
A.  $\frac{5R}{36}$   
B.  $\frac{36}{5}$   
C.  $\frac{36}{5R}$   
D.  $\frac{5}{36}$
- Q.6 The hydrogen atoms are excited to the stationary state designated by the principal quantum number  $n = 4$ . The number of maximum spectral lines are observing  
A. 2  
B. 4  
C. 3  
D. 6
- Q.7 Bracket series is obtained when all the transition of electron terminate at  
A. First orbit  
B. Third orbit  
C. Second orbit  
D. Fourth orbit
- Q.8 Number of the emission spectra are  
A. One  
B. Three  
C. Two  
D. Four
- Q.9 When electron jumps from  $n^{\text{th}}$  to the  $p^{\text{th}}$  orbit in an hydrogen atom then the wavelength of the emitted radiation is given by  
A.  $\frac{1}{\lambda} = R_H \left[ \frac{1}{p^2} - \frac{1}{n^2} \right]$   
B.  $\frac{1}{\lambda} = \frac{1}{R_H} \left[ \frac{1}{p^2} - \frac{1}{n^2} \right]$   
C.  $\frac{1}{\lambda} = R_H \left[ \frac{1}{n^2} - \frac{1}{p^2} \right]$   
D.  $\frac{1}{\lambda} = \frac{1}{R_H} \left[ \frac{1}{4^2} - \frac{1}{n^2} \right]$
- Q.10 Photon of highest frequency emitted in Lyman series is given as:  
A.  $f = \frac{c}{R_H}$   
B.  $f = \frac{R_H}{c}$   
C.  $f = R_H c$   
D.  $f = \frac{1}{R_H c}$

## Topic - 12

- Q.11 Which of the following is true for number of spectral lines in going from Lyman series to P-fund series?  
 A. Increases  
 B. Decreases  
 C. Unchanged  
 D. May decrease or increase
- Q.12 Number of spectral lines in hydrogen atom is  
 A. 3  
 B. 6  
 C. 15  
 D. infinite
- Q.13 Figure shows the energy levels P, Q, R, S and G of an atom where G is the ground state. A red line in the emission spectrum of the atom can be obtained by an energy level change from Q to S. A blue line can be obtained by following energy level change
- 
- A. R to S  
 B. Q to R  
 C. R to G  
 D. P to Q
- Q.14 The maximum wavelength of Lyman series is.....  
 A.  $\frac{4}{3R}$   
 B.  $\frac{1}{R^2}$   
 C.  $\frac{C}{R}$   
 D.  $\frac{1}{RC}$
- Q.15 An atom is excited to an energy level  $E_1$  from its ground state energy level  $E_0$ . The wavelength of the radiation emitted is  
 A.  $(E_0 - E_1) / hc$   
 B.  $\frac{E_1 - E_0}{hc}$   
 C.  $(E_1 - E_0) / h$   
 D.  $\frac{hc}{(E_1 - E_0)}$
- Q.16 Which of the transitions in hydrogen atom emits a photon of lowest frequency ( $n =$  quantum number)  
 A.  $n = 2$  to  $n = 1$   
 B.  $n = 4$  to  $n = 3$   
 C.  $n = 3$  to  $n = 1$   
 D.  $n = 4$  to  $n = 2$
- Q.17 During the transition of Electron of Hydrogen atom from higher orbit to a third orbit, a photon of:  
 A. Paschen series is emitted  
 B. Lyman series is emitted  
 C. Balmer series is emitted  
 D. Brackett series is emitted
- Q.18 The radiations emitted from hydrogen filled discharge tube shows:  
 A. Continuous spectrum  
 B. Band spectrum  
 C. Line spectrum  
 D. None of these
- Q.19 An electron jumps from the 4<sup>th</sup> orbit to the 2<sup>nd</sup> orbit of hydrogen atom. Given the Rydberg's constant  $R = 10^5 \text{ cm}^{-1}$ . The frequency in Hz of the emitted radiation will be  
 A.  $\frac{3}{16} \times 10^{15}$   
 B.  $\frac{3}{16} \times 10^{15}$   
 C.  $\frac{9}{16} \times 10^{15}$   
 D.  $\frac{3}{4} \times 10^{15}$

## Topic - 12

- Q.20 If an electron jumps from 1st orbital to 3rd orbital, then it will.  
 A. Absorb energy  
 B. Release energy  
 C. No gain of energy  
 D. None of these
- Q.21 The ratio of the frequencies of the long wavelength limits of Lyman and Balmer series of hydrogen spectrum is  
 A. 27 : 5  
 B. 5 : 27  
 C. 4 : 1  
 D. 1 : 4
- Q.22 The electron in the hydrogen atom makes a transition from  $n = 2$  energy state to the ground state  $n = 1$ . The wavelength of emitted photon is  
 A.  $\frac{3R}{4}$   
 B.  $\frac{3}{4}$   
 C.  $\frac{4}{3R}$   
 D.  $\frac{4}{3}$
- Q.23 Shortest wavelength photon in the Balmer series is  
 A.  $\frac{4}{R}$   
 B.  $\frac{R}{4}$   
 C.  $\frac{1}{4}$   
 D. 4
- Q.24 To find longest wavelength radiation in Balmer series, the value of  $n$  used is  
 A. 2  
 B. 4  
 C. 3  
 D.  $\infty$
- Q.25 Photon of highest frequency will be absorbed when transition takes place from:  
 A. 1<sup>st</sup> to 5<sup>th</sup> orbit  
 B. 5<sup>th</sup> to 1<sup>st</sup> orbit  
 C. 3<sup>rd</sup> to 5<sup>th</sup> orbit  
 D. 4<sup>th</sup> to 5<sup>th</sup> orbit
- Q.26 Balmer series lies in that region of electromagnetic wave spectrum, which is known as:  
 A. Visible region  
 B. Ultraviolet region  
 C. Invisible region  
 D. Infra-red region
- Q.27 The relation for Paschen series is given as  
 A.  $\frac{1}{\lambda} = R_H \left( \frac{1}{2^2} - \frac{1}{n^2} \right)$   
 B.  $\frac{1}{\lambda} = R_H \left( \frac{1}{4^2} - \frac{1}{n^2} \right)$   
 C.  $\frac{1}{\lambda} = R_H \left( \frac{1}{3^2} - \frac{1}{n^2} \right)$   
 D.  $\frac{1}{\lambda} = R_H \left( \frac{1}{5^2} - \frac{1}{n^2} \right)$

## PAST PAPER MCQs

- Q.28 Which of the following series lies in ultraviolet region? (NMDCAT 2021)  
 A. Balmer series  
 B. Paschen series  
 C. Lyman series  
 D. Brackett series

## ANSWER KEY

## TOPIC-WISE MCQs &amp; PAST PAPER MCQs

1	B	11	B	21	A
2	A	12	D	22	C
3	B	13	C	23	A
4	B	14	A	24	C
5	C	15	D	25	A
6	D	16	B	26	A
7	D	17	A	27	C
8	B	18	C	28	C
9	A	19	C		
10	C	20	A		

## EXPLANATORY NOTES

- For the first excited level,  $n = 2$ .
- $r_2 = (2)^2 r_0 = 4r_0$ .  
 So, when a hydrogen atom is in its first excited level, its radius is 4 times of the Bohr radius.
  - Third excited state means fourth energy states so electron have energy  

$$E_n = \frac{-E_0}{n^2} = -\frac{13.6 \text{ eV}}{4^2} = -0.85 \text{ eV}$$
 Hence ionization energy  $= -E_4 = -(-0.85 \text{ eV}) = +0.85 \text{ eV}$
  - We know  $\frac{1}{\lambda_n} = R \left( \frac{1}{p^2} - \frac{1}{n^2} \right)$  where  $p$  is lower state and  $n$  is higher state. So,  $n = p+1$
  - In Hydrogen spectrum Lyman series is strongest and lies in the ultraviolet region.
  - $\frac{1}{\lambda_n} = R \left( \frac{1}{p^2} - \frac{1}{n^2} \right)$  For longest wavelength in Balmer series  $p=2$  &  $n=3$   

$$\frac{1}{\lambda} = R \left( \frac{1}{2^2} - \frac{1}{3^2} \right) = R \left( \frac{1}{4} - \frac{1}{9} \right) = \frac{5R}{36} \Rightarrow \lambda = \frac{36}{5R}$$
  - No of spectral lines  $= \frac{n(n-1)}{2}$  Here  $n = 4$ , So  $\frac{4(4-1)}{2} = 6$
  - Bracket series is obtained when all transitions of electron terminate at  $p=4$
  - There are three emission spectra: Line Spectrum, Band Spectrum and Continuous Spectrum
  - We know  $\frac{1}{\lambda} = R_H \left[ \frac{1}{p^2} - \frac{1}{n^2} \right]$  where  $n=p+1$  ( $n$  is higher state and  $p$  is lower state)
  - Highest frequency in Lyman series when  $n=\infty$   $\frac{1}{\lambda} = R_H \left[ \frac{1}{1^2} - 0 \right] = R_H \Rightarrow R_H \lambda = 1$  Also  $f \lambda = c$
  - For Lyman series electron can transit from infinity to  $p=1$  and for P-fund series electron can transit from infinity to  $p=5$
  - In hydrogen atom only one electron but infinite energy states so there are infinite spectral lines.
  - If  $E$  is the energy radiated in transition then  $E_{p \rightarrow 0} > E_{p \rightarrow 1} > E_{p \rightarrow 2} > E_{p \rightarrow 3}$  For getting blue line energy radiated should be maximum  $\left( E \propto \frac{1}{\lambda} \right)$
  - We know  $\frac{1}{\lambda} = R_H \left[ \frac{1}{p^2} - \frac{1}{n^2} \right]$  For maximum wavelength of Lyman series  $p=1$  &  $n=2$   

$$\frac{1}{\lambda} = R_H \left[ \frac{1}{1^2} - \frac{1}{2^2} \right] \Rightarrow \frac{1}{\lambda} = R_H \left[ \frac{1}{1} - \frac{1}{4} \right] = R_H \frac{3}{4} \Rightarrow \lambda = \frac{4}{3R_H}$$
  - $$\Delta E = E_1 - E_0 = hf = \frac{hc}{\lambda} \Rightarrow \lambda = \frac{hc}{E_1 - E_0}$$

## Topic - 12

16. As we go to higher states the energy difference between the consecutive states decreases so photon of lowest frequency is obtained when it transit from  $n=4$  to  $n=3$
17. For Paschen series  $p=3$  and  $n=4,5,6,\dots$
18. Hydrogen gas gives the line spectrum.

$$19. \frac{1}{\lambda} = R \left( \frac{1}{2^2} - \frac{1}{4^2} \right) = \frac{3R}{16} \Rightarrow \lambda = \frac{16}{3R} = \frac{16}{3} \times 10^{-5} \text{ cm} \Rightarrow f = \frac{c}{\lambda} = \frac{3 \times 10^{10}}{\frac{16}{3} \times 10^{-5}} = \frac{9}{16} \times 10^{15} \text{ Hz}$$

20. When electron jumps from lower to higher state it will absorb the energy because higher states have higher energies.

$$21. \text{For Lyman series } \nu_{\text{Lyman}} = \frac{c}{\lambda_{\text{max}}} = Rc \left[ \frac{1}{(1)^2} - \frac{1}{(2)^2} \right] = \frac{3RC}{4}$$

$$\text{For Balmer series } \nu_{\text{Balmer}} = \frac{c}{\lambda_{\text{max}}} = Rc \left[ \frac{1}{(2)^2} - \frac{1}{(3)^2} \right] = \frac{5RC}{36} \text{ So, } \therefore \frac{\nu_{\text{Lyman}}}{\nu_{\text{Balmer}}} = \frac{27}{5}$$

$$22. \frac{1}{\lambda} = R \left( \frac{1}{1} - \frac{1}{4} \right) \Rightarrow \frac{1}{\lambda} = R \left( \frac{3}{4} \right) \Rightarrow \lambda = \frac{4}{3R}$$

$$23. \frac{1}{\lambda} = R \left( \frac{1}{2^2} - \frac{1}{\infty} \right) \Rightarrow \frac{1}{\lambda} = \frac{R}{4} \Rightarrow \lambda = \frac{4}{R}$$

$$24. \frac{1}{\lambda} = R \left( \frac{1}{2^2} - \frac{1}{n^2} \right), \text{ For longest wavelength } = n=3$$

$$25. \Delta E = hf \Rightarrow \Delta E \propto f$$

26. Balmer series lies in visible region of electromagnetic wave spectrum.

$$27. \frac{1}{\lambda} = R \left( \frac{1}{2^2} - \frac{1}{n^2} \right)$$

28. Lyman series lies in ultraviolet region of electromagnetic wave spectrum.

13  
TOPIC

## NUCLEAR PHYSICS

## PRACTICE EXERCISE


- Q.1 What is the half-time of a radioactive sample (in minutes), if its mean life is 200 s?  
A. 0.69 min  
C. 2.57 min  
B. 2 min  
D. 2.31 min
- Q.2 What will happen in a time of 7 hours, if a radioactive substance has an average life of 7 hours?  
A. Half of the active nuclei decay  
C. More than half of the active nuclei decay  
B. Less than half of the active nuclei decay  
D. All active nuclei decay
- Q.3 The difference between  $U^{235}$  and  $U^{238}$  atom is that  
A.  $U^{238}$  contains 3 more protons  
C.  $U^{238}$  contains 3 more protons and 3 more electrons.  
B.  $U^{238}$  contains 3 more neutrons and 3 more electrons.  
D.  $U^{238}$  contains 3 more neutrons
- Q.4 Which two nuclei contain the same number of neutrons?  
A.  ${}^{12}_6\text{C}$  and  ${}^{14}_6\text{C}$   
C.  ${}^{16}_7\text{N}$  and  ${}^{15}_8\text{O}$   
B.  ${}^{23}_{11}\text{Na}$  and  ${}^{24}_{12}\text{Mg}$   
D.  ${}^{32}_{14}\text{Si}$  and  ${}^{32}_{15}\text{P}$
- Q.5 What is the ratio of the nuclear densities of two nuclei having mass numbers in the ratio 1:4?  
A. 1:1  
C. 1:4  
B. 1:2  
D. 1:3
- Q.6 The radius for an atom is \_\_\_\_\_ times the radius of the nucleus.  
A.  $10^5$   
C.  $10^{15}$   
B.  $10^{10}$   
D.  $10^{20}$
- Q.7  $\alpha$ ,  $\beta$  and  $\gamma$  are emitted from a radioactive substance  
A. Spontaneously  
C. When it interacts with the other particles  
B. When it is heated  
D. When it is exposed to light
- Q.8 The order of penetration power of  $\alpha$  and  $\beta$  and  $\gamma$ -rays is  
A.  $\alpha > \beta > \gamma$   
C.  $\alpha > \beta < \gamma$   
B.  $\alpha < \beta > \gamma$   
D.  $\alpha < \beta < \gamma$
- Q.9 In a given reaction  ${}^A_Z\text{X} \longrightarrow {}^A_{Z+1}\text{Y} \longrightarrow {}^{A-4}_{Z-1}\text{B} \longrightarrow {}^{A-4}_{Z-1}\text{B}$   
Radioactive radiations are emitted in the sequence  
A.  $\gamma, \beta, \gamma$   
C.  $\gamma, \alpha, \beta$   
B.  $\beta, \alpha, \gamma$   
D.  $\alpha, \gamma, \beta$
- Q.10 When boron  ${}^{10}_5\text{B}$  is bombarded by neutrons,  $\alpha$ -particles are emitted. The resulting nucleus has the mass number  
A. 11  
C. 7  
B. 6  
D. 15
- Q.11 In which radioactive disintegration neutron dissociates into proton and electron?  
A.  $\alpha$ -emission  
C.  $\gamma$ -emission  
B.  $\beta$ -emission  
D. None of these

Topic-13

Q.12  ${}_{92}^{238}\text{U}$  nucleus emits two  $\alpha$ -particles and two  $\beta$ -particles and transforms into a thorium nucleus. Which of the following is the mass number and atomic number of the thorium nucleus so produced?

- A. 230, 90  
 B. 230, 88  
 C. 234, 90  
 D. 234, 88

Q.13 Specify the parent, daughter nuclei and emitted particle in given reaction



- Parent nucleus      Daughter nucleus      Emitted particle
- A. Thorium              Uranium  
 B. Uranium              Thorium  
 C. Thorium              Uranium  
 D. Uranium              Thorium

Q.14 The decay of a nucleus is accompanied by the emission of two  $\beta$ -particles and  $\alpha$ -radiation. What effect (if any) does this decay have on the proton number and the nucleon number of the nucleus?

	Proton number	Nucleon number
A	Increases	decreases
B	decreases	Increases
C	unchanged	decreases
D	decreases	unchanged

Q.15  $\alpha$ -particle is bombarded on  ${}^{14}\text{N}$  as a result  ${}^{17}\text{O}$  is formed. The particle emitted is

- A. Neutron              B. Electron  
 C. Proton                D. Positron

Q.16 If  ${}_{92}\text{U}^{233}$  decays twice by  $\alpha$ -emission, the resulting isotopes will be

- A.  ${}^{225}\text{Ra}_{88}$               B.  ${}^{234}\text{Pa}_{88}$   
 C.  ${}^{225}\text{Rn}_{86}$               D.  ${}^{229}\text{Th}_{90}$

Q.17 The following represents a sequence of radioactive decays involving two  $\alpha$ -particles and one  $\beta$ -particle.  ${}_{85}^{217}\text{At} \xrightarrow{\alpha} \text{V} \xrightarrow{\alpha} \text{W} \xrightarrow{\beta} \text{X}$

What is the nuclide X?

- A.  ${}_{85}^{213}\text{At}$               B.  ${}_{82}^{209}\text{Pb}$   
 C.  ${}_{81}^{215}\text{Ir}$                 D.  ${}_{81}^{217}\text{Tl}$

Q.18 Radon-220 is radioactive and decays to Polonium-216 with the emission of an  $\alpha$ -particle. The equation for the radioactive decay is shown.  ${}_{86}^{220}\text{Rn} \rightarrow {}_{84}^{216}\text{Po} + {}_2^4\text{He}$  How many neutrons are in the radon and polonium nuclei?

	Rn	Po
A.	86	84
B.	134	132
C.	220	212
D.	220	216

Q.19 In alpha decay, the ratio of decrease in proton number to the decrease in neutron number is

- A. 2 : 1  
 B. 1 : 1  
 C. 1 : 2  
 D. 4 : 1


Q.20 Radon  ${}_{86}^{222}\text{Rn}$  decays by  $\alpha$ - and  $\beta$ -emission to bismuth  ${}_{83}^{214}\text{Bi}$ . For the decay of each nucleus of radon. How many  $\alpha$ - and  $\beta$ - particles are emitted?

	$\alpha$ -particles	$\beta$ -particles
A.	1	1
B.	2	1
C.	1	2
D.	2	2

Q.21 A radioactive nucleus undergoes a series of decay according to the scheme  $A \xrightarrow{\alpha} A_1 \xrightarrow{\beta} A_2 \xrightarrow{\alpha} A_3 \xrightarrow{\gamma} A_4$ . If the mass number and atomic number of A are 180 and 72 respectively, then what are these number for  $A_4$ .

- A. 172 and 69  
 B. 176 and 69  
 C. 174 and 70  
 D. 176 and 70

Q.22 Neutron decay in the free space is given as follows



Then the parenthesis represents an

- A. Photon                      B. Neutrino  
 C. Graviton                    D. Antineutrino

Q.23 The ratio of the rate of decay of a parent atom to the number of radioactive nuclei present at that time is equal to:

- A. Half-life                      B. Mean life  
 C. Decay constant              D. Activity

Q.24 The masses of two radioactive substances are same and their half-lives are 1 year and 2 years respectively. The ratio of their activities after six years will be:

- A. 1:4                              B. 4:1  
 C. 1:8                              D. 8:1

Q.25 Let T be the mean life of a radioactive sample. 75% of the active nuclei present in the sample initially will decay in time.

- A. 2T                              B. 4T  
 C.  $\frac{1}{2}(\ln 2)T$                   D.  $2(\ln 2)T$

Q.26 In a sample of radioactive material, what fraction of the inertial number of active nuclei will remain undisintegrated after half of a half-life of the sample?

- A. 1/4                              B.  $\frac{1}{\sqrt{2}}$   
 C.  $\frac{1}{2\sqrt{2}}$                           D.  $\sqrt{2}-1$

Topic-13


- Q.27 If the decay constant of an element is smaller then  
 A. More of its atoms will decay in a particular interval  
 B. Its infinite atoms will decay in a particular interval  
 C. Less of its atoms will decay in a particular interval  
 D. Both A. and C. are possible depending upon the half-life of element

- Q.28 Three-fourths of the radioactive nuclei present in a radioactive sample decay in  $\frac{3}{4}s$ . The half-life of the sample is:

- A.  $\frac{3}{4}s$   
 B.  $\frac{3}{8}s$   
 C.  $1s$   
 D.  $\frac{1}{2}s$

- Q.29 The decay constant of radium is  $4.28 \times 10^{-4}$  per year. Its half-life will be  
 A. 1240 years  
 B. 1620 years  
 C. 2000 years  
 D. 63 years

- Q.30 Which of the following graph represents the variation of activity A of a radioactive substance with time t?



- Q.31 The activity of a radioactive sample is 1.6 curie and its half-life is 2.5 days. Its activity after 10 days will be  
 A. 0.8 curie  
 B. 0.1 curie  
 C. 0.4 curie  
 D. 0.16 curie

- Q.32 A radioactive element emits 200 particles per second. After three hours 25 particles per second are emitted. The half-life period of element will be  
 A. 50 minutes  
 B. 70 minutes  
 C. 60 minutes  
 D. 80 minutes

- Q.33 The half-life of  $\text{Bi}^{210}$  is 5 days. If we start with 50000 atoms of this isotope, then which of the following will be the number of atoms left over after ten days?  
 A. 5000  
 B. 12500  
 C. 20000  
 D. 25000

- Q.34 The sample of radioactive element decayed after 5 - half lives  
 A.  $\frac{1}{32}N_0$   
 B.  $\frac{N_0}{31} - \frac{N_0}{32}$   
 C.  $\frac{31}{32}N_0$   
 D.  $\frac{N_0}{32} - \frac{N_0}{31}$

- Q.35 Relation for half-life of any radioactive element is  
 A.  $T_{1/2} = \lambda \ln 2$   
 B.  $\lambda = T_{1/2} \ln 2$   
 C.  $T_{1/2} = \frac{\ln 2}{\lambda}$   
 D.  $T_{1/2} = \frac{\lambda}{\ln 2}$

Topic-13

- Q.36 The half-life period of radium is 1600 years. Its average life time will be  
 A. 3200 years  
 B. 2319 years  
 C. 4800 years  
 D. 4217 years

- Q.37 The decay constant of a radioactive element is 0.61 per second. Its half-life period is  
 A. 0.693 sec  
 B. 6.93 sec  
 C. 69.3 sec  
 D. 692 sec

- Q.38 What is the ratio of un-decayed and decayed nuclei of a radioactive element after 3 half-lives have elapsed?  
 A. 1/7  
 B. 7/8  
 C. 7  
 D. 8/7

- Q.39 Which of the following is the percentage of the original quantity of a radioactive material left after five half-lives approximately?  
 A. 3%  
 B. 5%  
 C. 10%  
 D. 20%

- Q.40 Which one of the following can be preferred for the treatment of skin?  
 A.  $\beta$ -particles  
 B. x-rays  
 C.  $\gamma$ -radiations  
 D.  $\alpha$ -rays

- Q.41 For skin cancer which of following can be used:  
 A. Iodine 131  
 B. Carbon 14  
 C. Phosphorous 32  
 D. Carbon 12

- Q.42 Radio isotope cobalt which emit high energy  $\gamma$ -rays is used for the treatment of  
 A. Temperature  
 B. Liver  
 C. Bones  
 D. Cancer

- Q.43 Cancerous tissue in a thyroid gland can be detected by the intake of  
 A. Radio iodine  
 B. Radio carbon  
 C. Radio sodium  
 D. Radio phosphors

- Q.44 Circulation of blood can be studied by:  
 A. Sodium - 24  
 B. Carbon 14  
 C. Strontium 90  
 D. Iodine 131

- Q.45 Which one of the following is the most useful tracer?  
 A. Strontium - 90  
 B. Carbon - 14  
 C. Iodine - 131  
 D. Cobalt - 60

PAST PAPER MCQs

- Q.46 A certain radioactive mass decays from 64 gm to 2 gm in 20 days. What is its half-life? (MCAT 2008)  
 A. 5 days  
 B. 10 days  
 C. 4 days  
 D. 6 days
- Q.47 When a helium atom loses an electron, it becomes: (MCAT 2008)  
 A. An alpha particle  
 B. A positive helium ion  
 C. Proton  
 D. A negative helium ion

(MCAT 2008)

Topic-13

Beta ray emitted by a radioactive substance is:

- A. An electron which was existing outside the nucleus.
- B. An electron emitted by the nucleus as a result of the decay of neutron inside the nucleus.
- C. An electron which was existing inside the nucleus.
- D. A pulse of electromagnetic wave.

Q.49 The emission of  $\gamma$ -radiations from the nucleus is generally represented by the equation: (MCAT 2009)

- A.  ${}_Z X^A \rightarrow {}_Z X^A + \gamma$ -radiations
- B.  ${}_Z X^A \rightarrow {}_Z X^A + \beta$ -particles
- C.  ${}_Z X^A \rightarrow {}_{Z-1} X^A + \gamma$ -radiations
- D.  ${}_Z X^A \rightarrow {}_Z X^A + \gamma$ -radiations

Q.50 In the half-life of an element, the equation for the number of decaying atoms is given by: (MCAT 2010)

- A.  $\Delta N = N\Delta t$
- B.  $\Delta N \propto -n\Delta t$
- C.  $\Delta N = KN\Delta t$
- D.  $\Delta N = -\Delta N\Delta t$

Q.51 Decay constant ' $\lambda$ ' is given as:

- A.  $\frac{\Delta N / N}{\Delta t}$
- B.  $-\frac{N}{\Delta t}$
- C.  $-\frac{\Delta N}{\Delta t}$
- D.  $\frac{\Delta N / N}{\Delta t}$

Q.52 Ionizing capability of gamma rays is:

- A. Equal to alpha and beta particle
- B. Less than both alpha and beta particles
- C. Less than alpha but greater than beta particles
- D. Less than beta but greater than alpha particles

Q.53 Half-life of a radioactive element is:

- A. Inversely proportional to square of decay constant
- B. Directly proportional to decay constant
- C. Directly proportional to square of decay constant
- D. Inversely proportional to decay constant

Q.54 The transformation of a neutron into proton in the nucleus gives rise to emission of: (MCAT 2011)

- A. Beta particles
- B. Gamma particles
- C. Alpha particles
- D. X-rays

Q.55 The ratio of the rate of decay of a parent atom to the number of radioactive nuclei present at that time is equal to: (MCAT 2011)

- A. Half-life of radioactive element
- B. Decay constant of radioactive element
- C. Mean life
- D. Activity of radioactive element

Q.56 What is the charge on alpha particles emitted during the phenomenon of radioactivity? (MCAT 2012)

- A.  $+e$
- B.  $-2e$
- C.  $-e$
- D.  $+2e$


Q.57 A radioactive nuclide decays by emitting an  $\alpha$ -particle and a  $\gamma$ -ray photon, the change in the nucleon number will be: (MCAT 2012)

- A. -4
- B. -2
- C. -2
- D. -3

Q.58 A half-life of sodium-24 is \_\_\_\_\_ which is used to estimate the volume of blood in a patient: (MCAT 2012)

- A. 6 hours
- B. 8 hours
- C. 15 hours
- D. 15 days

Q.59 In a radioactive phenomenon, observation shown in figure where  $\alpha$  deviates lesser than  $\beta$  in same electric or magnetic field (not shown in the figure). What is the reason of less deviation of  $\alpha$ ? (MCAT 2012)



- A.  $\alpha$  is a lighter particle
- B.  $\alpha$  is heavier particle
- C.  $\alpha$  is very fast moving particle
- D. None of these

Q.60 Which of the following effect is observed due to emission of  $\beta$  during the phenomenon of radioactivity? (MCAT 2012)

- A. A increases by 1 and Z remains same
- B. Z decreases by 1 and A remains same
- C. Z increases by 1 and A remains same
- D. A decreases by 1 and Z remains same

Q.61 Isotopes are those nuclei of an element that have: (MCAT 2013)

- A. Same mass number but different atomic number
- B. Different mass number as well as atomic number
- C. Same mass number as well as atomic number
- D. Same atomic number but different mass number

Q.62 Emission of alpha decay from a radioactive substance cause: (MCAT 2013)

- A. Decrease in 'Z' by 4 and decrease in 'A' by 4
- B. Decrease in 'Z' by 1 and 'A' remains same
- C. Decreases in 'A' by 1 and 'Z' remains same
- D. Decrease in 'A' by 4 and decrease in 'Z' by 2

Q.63 Which one of the following emissions takes place in a nuclear reaction? (MCAT 2013)

- A. Alpha
- B. Beta
- C. Gamma
- D. Photons

Q.64 Among the three types of radioactive radiation, which have strongest penetration power? (MCAT 2013)

- A. Alpha
- B. Beta
- C. Gamma
- D.  $\alpha$ ,  $\beta$  and  $\gamma$  have same penetration

Q.65 Emission of radiation from radioactive substance is (MCAT 2013)

- A. Dependent on both temperature and pressure
- B. Independent of both temperature and pressure
- C. Independent of temperature but dependent on pressure
- D. Independent of pressure but dependent on temperature

Q.66 In the nuclear reaction  ${}_{11}\text{Na}^{24} \rightarrow {}_{12}\text{Mg}^{24} + X$  the particle X is: (ETEA 2013)


- A. Electron
- B. Positron
- C. Proton
- D. Neutron



Topic-13

Nuclear Physics

Q.67 Three points of radioactive radiation are observed as shown in the figure presence of electric field, which type of radiation is shown in the path '1'?



- A. Alpha
- B. Beta
- C. Gamma
- D. Cathode ray

Q.68 A beta particle is a fast-moving electron, During a  $\beta$  decay how the atomic number and mass number of a nucleus change?

	Atomic number	Mass number
A.	Remains the same	Increases by one
B.	Increases by one	Remains the same
C.	Increases by one	Decreases by two
D.	Decreases by two	Decreases by four


Q.69 A uranium isotope  ${}_{92}^{234}\text{U}$  undergoes one  $\alpha$ -decay and one  ${}_{-1}^0\beta$ -decay. What is the atomic number of the final product?

- A. 90
- B. 89
- C. 91
- D. 88

Q.70 A naturally occurring radioactive element decays two alpha particles. Which one of the following represents the status of daughter element with respect to mass number A and charge number Z?

- A. Z decreases by 4 and A decreases by 2
- B. Z decreases by 4 and A decreases by 8
- C. Z decreases by 2 and A decreases by 4
- D. Z decreases by 8 and A decreases by 4

Q.71 A radioactive isotope W decay to x which decay to Y and Y decays to Z as represented by the figure below.



What is the change in the atomic number from W to Z?

- A. Increases by 3
- B. Increases by 5
- C. Decreases by 3
- D. Decreases by 5

Q.72 In the reaction  ${}_{90}^{234}\text{Th} \rightarrow {}_{91}^{234}\text{Pa} + {}_{-1}^0\text{e}$  the electron  ${}_{-1}^0\text{e}$  emits from the

- A. 1<sup>st</sup> orbit
- B. Nucleus
- C. 2<sup>nd</sup> orbit
- D. Valence shell

Q.73 According to the equation  ${}^A_Z\text{X} \rightarrow \text{Y} + 3\alpha$  particles, what are the atomic and mass numbers of 'Y'?

- A. Z-6, A-12
- B. Z+1, A
- C. Z-2, A-4
- D. Z+3, A

Q.74 A certain radioactive nuclide of mass number 'x' decay by  $\beta$ -emission and  $\alpha$ -emission to a second nuclide of mass number 't', which of the following correctly relates 'x' and 't'?

- A.  $x = t - 4$
- B.  $x + 3 = t$
- C.  $x = t + 4$
- D.  $x - 1 = t$

Topic-13

Nuclear Physics

Q.75 During the decay of radioactive isotopes  ${}_{92}^{238}\text{X}$  to a stable isotope, six  $\alpha$ -particles and four  $\beta$ -particles are emitted, what is the atomic number 'Z' and mass number 'A' of the stable isotopes:

- A. Z = 70, A = 220
- B. Z = 82, A = 212
- C. Z = 78, A = 212
- D. Z = 82, A = 208

Q.76 Wavelength of  $\gamma$ -rays is:

- A. Equal to the X-rays
- B. Shorter than X-rays
- C. Longer than X-rays
- D. Broader than X-rays

Q.77 Thorium is transformed after the emission of  $\beta$ -particle into:

- A. Bismuth
- B. Polonium
- C. Protactinium
- D. Palladium

Q.78 Emission of  $\gamma$ -rays from radioactive element result into:

- A. Increase of charge number 1
- B. No change in the charge number
- C. Decrease of mass number by 1
- D. Decrease charge number by 1

Q.79 The relation between decay constant ' $\lambda$ ' and half-life ' $T_{1/2}$ ' of radioactive substance is:

- A.  $\lambda = \frac{1}{T_{1/2}}$
- B.  $\lambda = T_{1/2}$
- C.  $\lambda = 0.693 T_{1/2}$
- D.  $\lambda = \frac{0.693}{T_{1/2}}$

Q.80 Radiation damages living organism is primarily due to:

- A. Excitation phenomena
- B. Ionization
- C. Photo electric effect
- D. Heating

Q.81 Complete the radioactive equation  ${}^A_Z\text{X} \rightarrow {}^A_B\text{Y} + \gamma$

- A.  ${}^A_{B-1}\text{Z}$
- B.  ${}^{A+1}_{B+1}\text{Z}$
- C.  ${}^{A+1}_{B-1}\text{Z}$
- D.  ${}^A_B\text{Z}$

Q.82 The quantity of uranium is 400g. After 3<sup>rd</sup> half-life, how much uranium will be left?

- A. 50g
- B. 100g
- C. 25g
- D. 200g


Q.83 The half-life of radium is about 1600 years. If 100 g radium existing now, 25 g will remain un-decayed after:

- A. 4800 years
- B. 2400 years
- C. 6400 years
- D. 3200 years

Q.84 Which of the following has maximum ionizing power?

- A.  $\alpha$
- B.  $\gamma$
- C.  $\beta$
- D. Neutron

Q.85 A radioactive nucleus X undergoes a series of decay according to the same:



If the mass number and atomic number of X are 180 and 72 respectively, the corresponding number of  $\alpha$  are:

- A. 176, 69
- B. 172, 69
- C. 176, 7
- D. 172, 71

Topic-13

- Q.86 Calculate the activity (decaying atom per unit time) of radioactive strontium-90 having  $6.7 \times 10^{21}$  atoms at  $t=0$  decay constant of strontium-90 is  $8.3 \times 10^{-10} \text{ s}^{-1}$  (MDCAT 2018)  
 A.  $8.01 \times 10^{10} \text{ Bq}$   
 B.  $5.6 \times 10^{12} \text{ Bq}$   
 C.  $5.6 \times 10^{11} \text{ s}^{-1}$   
 D.  $1.2 \times 10^{11} \text{ Bq}$
- Q.87 Calculate the half-life of bismuth-214 which has a decay constant of  $4.3 \times 10^3 \text{ s}^{-1}$  (MDCAT 2018)  
 A.  $2.9 \times 10^{-4} \text{ s}$   
 B.  $3.9 \times 10^3 \text{ s}$   
 C.  $1.6 \times 10^{-4} \text{ s}$   
 D.  $2.9 \times 10^3 \text{ s}$
- Q.88 Two radioactive samples,  $S_1$  and  $S_2$  have half-life 3 hours and 7 hours respectively. If they have the same activity at certain instant  $t$ , what is the ratio of the number of atoms of  $S_1$  to  $S_2$  at instant  $t$ ? (ETEA 2018)  
 A. 9:49  
 B. 49:9  
 C. 3:7  
 D. 7:3
- Q.89 In relation  $\lambda T_{1/2} = 0.693$ , which quantity is represented by  $\lambda$ . (MDCAT 2019)  
 A. Half-life  
 B. Activity  
 C. Wavelength  
 D. Decay constant
- Q.90 The main difference between X-Rays and  $\gamma$ -Rays is: (ETEA 2019)  
 A. Frequency  
 B. Wave length  
 C. Energy  
 D. Origin
- Q.91 There are initially 400 atoms in a radioactive sample. What would be the number of atoms after 3 half-lives? (ETEA 2019)  
 A. 400  
 B. 200  
 C. 50  
 D. 25
- Q.92 While using radiation therapy, cancerous thyroid is treated with \_\_\_\_\_ radioisotope: (ETEA 2019)  
 A. Carbon  
 B.  $^{235}\text{U}$  uranium  
 C. Thorium  
 D. Iodine - 131
- Q.93 Half-life of iodine-131 is 8 days. If 20mg is present initially, how much iodine is left behind after 2 half-lives? (NUMS 2020)  
 A. 10 mg  
 B. 5 mg  
 C. 2.5 mg  
 D. 1.25 mg
- Q.94  $4.5 \times 10^9$  year is the half-life of: (NUMS 2020)  
 A.  $\text{U}^{238}$   
 B.  $\text{U}^{235}$   
 C.  $\text{U}^{236}$   
 D.  $\text{U}^{14}$
- Q.95 Number of Quarks in hydrogen atom (MDCAT 2021)  
 A. 1  
 B. 2  
 C. 3  
 D. 4
- Q.96 Which radiations more ionization (MDCAT 2021)  
 A. Alpha  
 B. Beta  
 C. Gamma  
 D. None
- Q.97 High speed beta rays are (MDCAT 2021)  
 A. Electron  
 B. Positron  
 C. Electron and positron  
 D. Proton

# ANSWER KEY

## TOPIC-WISE MCQs & PAST PAPER MCQs

1	D	11	B	21	A	31	B	41	C	51	A	61	D	71	C	81	D	91	C
2	C	12	A	22	D	32	C	42	B	52	B	62	D	72	B	82	A	92	D
3	D	13	D	23	C	33	B	43	A	53	D	63	B	73	A	83	D	93	B
4	B	14	C	24	C	34	C	44	A	54	A	64	C	74	C	84	A	94	A
5	A	15	C	25	D	35	C	45	B	55	B	65	B	75	D	85	B	95	C
6	A	16	A	26	B	36	B	46	C	56	D	66	A	76	B	86	B	96	C
7	A	17	B	27	C	37	C	47	B	57	A	67	C	77	C	87	C	97	A
8	D	18	B	28	B	38	A	48	C	58	C	68	B	78	B	88	C		
9	B	19	B	29	B	39	A	49	D	59	B	69	C	79	D	89	D		
10	C	20	B	30	D	40	C	50	A	60	C	70	B	80	B	90	D		

# EXPLANATORY NOTES

- $T_{1/2} = 0.693t$   
 $T_{1/2} = 0.693 \times 200s$   
 $T_{1/2} = 138.6 / 60 \text{ min}$   
 $T_{1/2} = 2.31 \text{ minutes}$
- In one average life, i.e. at 7 hours, 63.2 % of the active nuclei will decay. Therefore, in a time of 7 hours, it can be considered that more than half of the active nuclei will decay.
- Isotopes are the atoms of an element having same atomic number Z but different mass number A. So  $U^{238}$  contains 3 more neutrons.
- Number of neutrons in  $Na_{11}^{23} = 23 - 11 = 12$   
 Number of neutrons in  $Mg_{12}^{24} = 24 - 12 = 12$
- As the nuclear density is independent of mass number, so the ratio of nuclear densities of the two given nuclei is 1:1. Its value  $d = 2.3 \times 10^{17} \text{ Kg m}^{-3}$
- $R_{\text{atom}} = 10^{-10} \text{ m}$  &  $R_{\text{nucleus}} = 10^{-15} \text{ m}$
- Emission of  $\alpha$ ,  $\beta$  and  $\gamma$ -radiations is spontaneous and random process.
- The penetrating power of  $\alpha$  is least and that of  $\gamma$ -radiations is the most.
- ${}^A_Z X \xrightarrow{\beta} {}^A_{Z+1} Y \xrightarrow{\alpha} {}^{A-4}_{Z-1} B \xrightarrow{\gamma} {}^{A-4}_{Z-1} B$ 
  - When  $\beta$  is emitted then charge will increase by one.
  - When  $\alpha$  is emitted then mass will reduce by 4 and charge will reduce by 2.
- ${}^{10}_5 B + {}^1_0 n \rightarrow {}^7_3 X + {}^4_2 He$  Resultant nucleus has mass number "7".
- In  $\beta$ -emission  ${}^1_0 n \rightarrow {}^0_{-1} \beta + {}^1_1 H$
- ${}^{238}_{92} U \xrightarrow{2\alpha} {}^{230}_{88} X \xrightarrow{2\beta} {}^{230}_{90} Th$
- Here parent nucleus is  ${}^{238}_{92} U$  uranium (146+92=238), thorium is  ${}^{234}_{90} Th$  (144+90=234) and emitted radiation is He (Alpha).
- $X^A_Z \xrightarrow{2\beta} X^A_{Z+2} \xrightarrow{\alpha} X^{A-4}_Z$
- ${}^{14}_7 N + {}^4_2 He = {}^{17}_8 O + {}^1_1 P$
- ${}^{233}_{92} U \xrightarrow{2\alpha} {}^{225}_{88} Ra$
- ${}^{217}_{85} At \xrightarrow{\alpha} {}^{213}_{83} V \xrightarrow{\alpha} {}^{209}_{81} W \xrightarrow{\beta} {}^{209}_{82} X$
- Number of neutrons in radon = 220-86=134  
 Number of neutrons in polonium = 216-84=132
- By emission of alpha two protons and two neutrons are decreased.
- ${}^{222}_{86} Rn \xrightarrow{\alpha} {}^{218}_{84} X \xrightarrow{\alpha} {}^{214}_{82} Y \xrightarrow{\beta} {}^{214}_{83} Bi$  (2- $\alpha$  particles & 1- $\beta$  particle)
- ${}^{180}_{72} A \xrightarrow{\alpha} {}^{176}_{70} A_1 \xrightarrow{\beta} {}^{176}_{71} A_2 \xrightarrow{\alpha} {}^{172}_{69} A_3 \xrightarrow{\gamma} {}^{172}_{69} A_4$
- The reaction for neutron decay is  ${}^1_0 n \rightarrow {}^1_1 H + {}^0_{-1} e + \bar{\nu}$  Here  $\bar{\nu}$  = anti - neutrino

- By definition of decay constant:  $\frac{dN}{dt} = N\lambda \Rightarrow \lambda = \frac{dN/dt}{N}$
- We know  $\frac{A}{A_0} = \frac{N}{N_0} = \frac{1}{2^n}$ , So  $\frac{A_1}{A_2} = \frac{2^3}{2^6} = \frac{1}{8}$
- When 75% decays, 25% is left un-decayed. This requires a time  $t = 2T_{1/2}$ , where  
 $T_{1/2} = \text{half life} = \frac{\ln 2}{\lambda}$ . Also  $T = \frac{t}{\lambda}$ .  $\therefore t = 2 \left( \frac{\ln 2}{\lambda} \right) = 2(\ln 2)T$
- As  $n = \frac{t}{T_{1/2}} = \frac{\frac{1}{2} T_{1/2}}{T_{1/2}} = \frac{1}{2}$ . Fractional un-decay =  $\frac{1}{2^n} = \frac{1}{2^{1/2}} = \frac{1}{\sqrt{2}}$
- $T_{1/2} = \frac{0.693}{\lambda} \Rightarrow \lambda T_{1/2} = 0.693$  It means for smaller  $\lambda$  half-life will be longer.
- $\frac{3}{4} N_0 = \frac{3}{2^2} N_0 \Rightarrow n = 2$  As  $t = nT_{1/2} \Rightarrow T_{1/2} = \frac{t}{n} = \frac{3/4}{2} = \frac{3}{8} \text{ sec}$
- As  $T_{1/2} = \frac{0.693}{\lambda} = \frac{0.693}{4.28 \times 10^{-4}} = 1620 \text{ year}$
- Activity of a radioactive element decays exponentially.
- Number of half lives  $n = \frac{10}{2.5} = 4 \Rightarrow \frac{A}{A_0} = \frac{N}{N_0} = \left( \frac{1}{2} \right)^n \Rightarrow A = 1.6 \times \left( \frac{1}{2} \right)^4 = 0.1 \text{ curie}$
- $n = 3 \Rightarrow T_{1/2} = \frac{t}{n} = \frac{60 \times 3}{3} = 60 \text{ minutes}$
- Number of half-lives (n = 2) We know  $N_R = N_0 \left( \frac{1}{2} \right)^n \Rightarrow \frac{50000}{4} = 12500$
- $N_D = N_0 \frac{2^n - 1}{2^n} \Rightarrow \frac{N_D}{N_0} = \frac{2^5 - 1}{2^5} = \frac{31}{32}$
- $T_{1/2} = \frac{\ln 2}{\lambda} \Rightarrow \lambda T_{1/2} = \ln 2$
- Average life =  $\frac{1}{\lambda} = \frac{T_{1/2}}{0.693} = \frac{1600}{0.693} = 2308 \approx 2319 \text{ years}$
- $T_{1/2} = \frac{0.693}{\lambda} = \frac{0.693}{0.01} = 69.3 \text{ sec}$
- $N_R = N_0 \left( \frac{1}{2^n} \right)$  &  $N_D = N_0 \left( \frac{2^n - 1}{2^n} \right) \Rightarrow \frac{N_R}{N_D} = \frac{1}{8} \times \frac{8}{7} = \frac{1}{7}$
- After 5 half-lives remaining amount =  $\frac{1}{2^5} = \frac{1}{32}$
- Percentage of remaining amount =  $\frac{1}{32} \times 100 \approx 3\%$
- $\gamma$ -radiations can be preferred for the treatment of skin

For skin cancer phosphorus-32 or strontium-90 may be used.  
 Radiotherapy with  $\gamma$ -rays from cobalt-60 is often used in the treatment of cancer of liver.  
 Radio-iodine is absorbed mostly by the thyroid gland.  
 Study the circulation of blood using radioactive isotope sodium-24.  
 The most useful tracer is Carbon - 14.

$m = 64g$

In

$1T_{1/2} = 32g$

$2T_{1/2} = 16g$

$3T_{1/2} = 8g$

$4T_{1/2} = 4g$

$5T_{1/2} = 2g$

$n = 5$

$T_{1/2} = \frac{t}{n} = \frac{20}{5} = 4 \text{ days.}$


7. When a helium atom loses an electron, it gets positive charge and becomes a positive helium ion.
8. Beta ray emitted by a radioactive substance is an electron which was existing inside the nucleus  ${}_Z X^A \longrightarrow {}_Z Y^A + \beta_{-1}^0 + \bar{\nu}$   
 Emission of  $\beta$ -particles is a nuclear phenomenon.
49.  ${}_Z X^A \longrightarrow {}_Z X^A + \gamma$ -radiation
50.  $\Delta N = -N\Delta t$
51.  $\lambda = -\frac{\Delta N / N}{\Delta t}$
52.  $\gamma$ -rays<sub>IE</sub> <  $\beta$ -rays<sub>IE</sub> <  $\gamma$ -particles<sub>IE</sub>
53.  $T_{1/2} = \frac{0.693}{\lambda} \Rightarrow T_{1/2} \propto \frac{1}{\lambda}$
54.  ${}_0 n^1 \longrightarrow {}_1 H^1 + \beta_{-1}^0 + \bar{\nu}$
55.  $\frac{\Delta N}{N\Delta t}$
56. Alpha particle is also known as Helium nuclide, and Helium nuclide have charge 2e.
57.  $X_Z^A \xrightarrow{\alpha} Y_{Z-2}^{A-4} \xrightarrow{\gamma} Y_{Z-2}^{A-4}$
58. Half-life of sodium-24 is 15 hours.
59.  $\alpha$ = particle has more momentum So, it will less deviate.
60.  $X_Z^A \xrightarrow{\beta_{-1}} Y_{Z+1}^A$
61. Definition
62.  $X_Z^A \xrightarrow{\gamma} Y_{Z-2}^{A-4}$
63.  ${}_{90} Th^{234} \longrightarrow {}_{91} Pa^{234} + \beta_{-1}^0$
64.  $\alpha < \beta < \gamma$
65. Emission of radiation from radioactive substance is a natural phenomenon. So, it is independent of both temperature and pressure.

66.  $\beta$ -decay  $X_Z^A \longrightarrow Y_{Z+1}^A + \beta_{-1}$
67. The electric field does not disturb  $\gamma$ -radiation because gamma radiation no charge.
68. During beta deca, Atomic number increases by 1 and mass number remains same.
69.  ${}_{92} U^{234} \xrightarrow{\alpha} {}_{90} Y^{230} \xrightarrow{\beta} {}_{91} Y^{230}$  So,  $z=91$
70.  ${}_Z X^A \xrightarrow{2\alpha} {}_{Z-4} Y^{A-8}$
71.  $W \xrightarrow{\beta} X \xrightarrow{\alpha} Y \xrightarrow{\alpha} Z$
72.  $\beta$ -particles emit from nucleus.
73.  $X_Z^A \xrightarrow{3\alpha} Y_{Z-6}^{A-12}$
74.  $X_Z^A \longrightarrow Y_{Z+1}^A \xrightarrow{\alpha} Y_{Z-1}^{A-4}$   
 $X - 4 = t$   
 $X = t + 4$
75.  $X_{90}^{232} \xrightarrow{6\alpha} X_{78}^{208} \xrightarrow{4\beta} X_{82}^{208}$
76.  $E = \frac{hc}{\lambda} \Rightarrow E \propto \frac{1}{\lambda} \Rightarrow \lambda_{\gamma\text{-rays}} < \lambda_{x\text{-rays}}$
77.  ${}_{90} Th^{234} \longrightarrow {}_{91} Pa^{234} + \beta_{-1}^0$
78.  $X_Z^A \longrightarrow X_Z^A + \gamma$ -rays
79.  $T_{1/2} \lambda = 0.693 \Rightarrow \lambda = \frac{0.693}{T_{1/2}}$
80. Radiation produced ionization
81.  ${}_b Z^a \longrightarrow {}_b Z + \gamma$
82.  $N = \frac{N_c}{2^n} = \frac{400}{2^3} = 50g$
83.  $N = N_0 \left(\frac{1}{2}\right)^n \Rightarrow 25 = 100 \frac{1}{2^n} \Rightarrow \frac{1}{4} = \frac{1}{2^2} = \frac{1}{2^n} \Rightarrow n = 2$   
 $T = nT_{1/2} = 2 \times 1600 = 3200 \text{ years}$
84.  $\alpha$ -particle has maximum ionizing power.
85.  ${}_{72} X^{180} \xrightarrow{\alpha} {}_{70} X_1^{176} \xrightarrow{\beta} {}_{71} X_2^{176} \xrightarrow{\alpha} {}_{69} X_3^{172} \xrightarrow{\gamma} {}_{69} X_4^{172}$
86.  $A = \lambda N$   
 $= 8.3 \times 10^{-10} \times 6.7 \times 10^{21} = 55.61 \times 10^{11} = 5.6 \times 10^{12} \text{ Bq.}$
87.  $T_{1/2} = \frac{0.693}{\lambda} = \frac{0.693}{4.3 \times 10^{13}} = 0.16 \times 10^{-1} = 1.6 \times 10^{-4} \text{ s}$

88.  $R = \frac{\Delta N}{\Delta t} = \lambda N = \frac{0.693N}{T} \Rightarrow N \propto T$
89. In relation  $\lambda T_{1/2} = 0.693$ ,  $\lambda$  is decay constant.
90. Both are E.M waves. X-Rays produced by Coolidge tube while  $\gamma$ -Rays produced by decelerated of nucleus
91.  $N_n = N_o \cdot \left(\frac{1}{2}\right)^n = \frac{400}{8} = 50$
92. Cancerous thyroid is treated with iodine -131
93.  $L_{cft} = \frac{1}{2^n} N_o$
94. Information of book.
95. Two up quarks and one down quark.
96.  $I.E$  of  $\alpha > I.E$  of  $B > I.E$  of  $\gamma$
97. High speed  $\beta$ -rays are electrons.

1  
UNITFORCE AND MOTION  
SELF ASSESSMENT TEST

- Q.1 A truck requires 3 Hrs to complete a journey of 150 km, what is the average speed?  
A. 50 km/hr  
C. 15 km/h
- Q.2 A car is moving around a tree in a circular path. What can be said about the average velocity?  
A. It is non zero  
C. It is greater than zero
- Q.3 A man covers half of his journey at 6km/hr and the remaining half at 3km/hr. His average speed is  
A. 4 km/hr  
C. 9 km/hr
- Q.4 The relative velocity of a particle moving with a velocity  $v$  w.r.t. itself is  
A.  $v$   
C. zero
- Q.5 A ball is dropped from a height 10m. The ball is embedded in sand 1m and stops, then  
A. Only momentum remains conserved  
C. Both momentum and K.E are conserved
- Q.6 A ball is thrown horizontally from the top of a tower. What happens to the horizontal component of its acceleration?  
A. First increases and then decreases  
C. Decrease
- Q.7 Velocity time graph of body X and Y is shown in fig. The ratio of the acceleration of Y to acceleration of X is



- A. 3:1  
C. 1:3
- B. 1: $\sqrt{3}$   
D.  $\sqrt{3}$ :1
- Q.8 Four projectiles are fired with the same velocity at angle, 25°, 40°, 55° and 70° with the horizontal. The range of projectile will be largest for the one projected at angle  
A. 70°  
C. 55°
- B. 40°  
D. 25°
- Q.9 A fighter plane drops a bomb when it is at the top of enemies target. Bomb misses the target due to  
A. Horizontal component of velocity  
C. Vertical component of velocity
- B. Action of gravity  
D. Bad weather
- Q.10 A projectile is projected with kinetic energy (K.E). If it has the maximum possible horizontal range, then its kinetic energy at the highest point will be:  
A. K.E  
C. 0.5 K.E
- B. 0.75 K.E  
D. 0.25 K.E
- Q.11 A car travels equal distance in the same direction with velocities 60 km h<sup>-1</sup>, 20 km h<sup>-1</sup> and 10 km h<sup>-1</sup> respectively. The average velocity of the car over the whole journey of motion is  
A. 8 ms<sup>-1</sup>  
C. 7 ms<sup>-1</sup>
- B. 6 ms<sup>-1</sup>  
D. 5 ms<sup>-1</sup>

Q.12 A particle is moving eastwards with a velocity  $5 \text{ ms}^{-1}$ . In 10 s, the velocity changes to  $5 \text{ ms}^{-1}$  northward. The average acceleration in this time is

- A.  $\frac{1}{\sqrt{2}} \text{ ms}^{-2} \text{ NE}$
- B.  $\frac{1}{2} \text{ ms}^{-2} \text{ N}$
- C. Zero
- D.  $\frac{1}{\sqrt{2}} \text{ ms}^{-2} \text{ NW}$

Q.13 A force of 5 Newton acts on a body of weight 9.8 Newton. What is the acceleration produced in  $\text{ms}^{-2}$ ?

- A. 0.51
- B. 1.46
- C. 49.00
- D. 5.00


Q.14 Time of flight of projectile is

- A.  $\frac{v \sin \theta}{g}$
- B.  $\frac{2v \sin \theta}{g}$
- C.  $\frac{v \sin 2\theta}{g}$
- D.  $\frac{v^2}{g}$

Q.15 A cyclist riding at a speed of 5m/sec braked with uniform deceleration and stopped in 3m. The acceleration is

- A.  $2.16 \text{ m/sec}^2$
- B.  $-2.16 \text{ m/sec}^2$
- C.  $4.16 \text{ m/sec}^2$
- D.  $-4.16 \text{ m/sec}^2$

Q.16 A body moves from A to B in a straight line then in semi-circle of radius as shown in fig. What is the displacement of the body?



- A. 7 cm
- B. 9 cm
- C. 5 cm
- D. 12 cm

Q.17 Angle between action and reaction forces is

- A.  $0^\circ$
- B.  $\frac{\pi}{2} \text{ rad}$
- C.  $\pi \text{ rad}$
- D.  $2\pi \text{ rad}$

Q.18 The P.E gained by projectile when it reaches at maximum height is expressed by equation

- A.  $K.E_i \cos^2 \theta$
- B.  $K.E_i \tan \theta$
- C.  $K.E_i \sin^2 \theta$
- D.  $K.E_i \cot \theta$

Q.19 A 1 kg ball moving in some direction at 8 m/sec hits a 4 kg ball moving at 6 m/sec in the same direction. Assume that the 1 kg ball stops after collision. What will be the velocity, after collision, of the 4 kg ball if the 1 kg ball stops?

- A. 10 m/sec
- B. 0 m/sec
- C. 8 m/sec
- D. 12 m/sec

Q.20 A runner runs 100m in 10s, then turns around and jogs 50m back toward the starting point in 30s. What is his average speed and average velocity in m/s.

- A. 3, 1.25
- B. 3.75, 1.25
- C. 4.25, 1.25
- D. 3, 1

Q.21 A car of mass 1000 kg moving with  $25 \text{ ms}^{-1}$  collide straight with stationary truck of mass 1500 kg, and start moving together. Find the velocity of truck.

- A.  $15 \text{ ms}^{-1}$
- B.  $25 \text{ ms}^{-1}$
- C.  $10 \text{ ms}^{-1}$
- D.  $6 \text{ ms}^{-1}$

Q.22 A body moves from point P (3, 4) to Q (5, 7). Its displacement vector would be

- A.  $8\hat{i} + 10\hat{j}$
- B.  $8\hat{i} + 2\hat{j}$
- C.  $2\hat{i} - 2\hat{j}$
- D.  $2\hat{i} + 3\hat{j}$

Q.23 A particle starting from rest undergoes a rectilinear motion with acceleration a. The variation of 'a' with time t is shown in graph. Maximum velocity attained by the particle during the motion is:



- A. 55 m/s
- B. 110 m/s
- C. 550 m/s
- D. 650 m/s

Q.24 A missile is fired for maximum range with an initial velocity of 20 m/s. If  $g=10 \text{ m/s}^2$ , the range of the missile is

- A. 30 m
- B. 20 m
- C. 40 m
- D. 60 m


Q.25 A ball is projected horizontally from the top of a cliff on the surface of the earth with a speed of  $40 \text{ ms}^{-1}$ . Assuming that there is no air resistance, what will its speed be 3 s later?

- A.  $30 \text{ ms}^{-1}$
- B.  $40 \text{ ms}^{-1}$
- C.  $50 \text{ ms}^{-1}$
- D.  $60 \text{ ms}^{-1}$

Q.26 The rate of change of momentum of a body falling freely under gravity is equal to its

- A. Impulse
- B. Kinetic energy
- C. Power
- D. Weight

Q.27 Displacement time graph is shown in figure below acceleration will be



- A.  $5 \text{ ms}^{-2}$
- B.  $10 \text{ ms}^{-2}$
- C.  $2.5 \text{ ms}^{-2}$
- D. 0

Q.28 At which point for a projectile its kinetic energy is completely converted into potential energy

- A. At point of projection
- B. At the highest point
- C. Point to hit the ground
- D. Not possible

Q.29 A 10 kg object collides with stationary 5 kg object and after collision they stick together and move forward with velocity  $4 \text{ ms}^{-1}$ . What is the velocity with which the 10 kg object hit the second one?

- A.  $6 \text{ ms}^{-1}$
- B.  $8 \text{ ms}^{-1}$
- C.  $12 \text{ ms}^{-1}$
- D.  $10 \text{ ms}^{-1}$

Q.30 A force of 10 N acts on a body of mass 20 kg for 10 seconds. The change in its momentum is

- A. 50 kg-m/s
  - B. 100 kg-m/s
  - C. 300 kg-m/s
  - D. 1000 kg-m/s
- Q.31 An aircraft is moving with a velocity of  $300\text{ms}^{-1}$ . If all the forces acting on it are balanced, then
- A. It still moves with the same velocity
  - B. It will explode
  - C. It will lose its velocity gradually
  - D. It will fall down instantaneously

- Q.32 If a car covers  $\frac{2}{5}$ th of the total distance with  $v_1$  speed and  $\frac{3}{5}$ th distance with  $v_2$  then average speed is
- A.  $\frac{v_1 + v_2}{2}$
  - B.  $\frac{2v_1v_2}{v_1 + v_2}$
  - C.  $\frac{5v_1v_2}{3v_1 + 2v_2}$
  - D.  $\frac{1}{2}\sqrt{v_1v_2}$

Q.33 If initial velocity of projectile is doubled then which is correct

	Time of flight	Max. height	Horizontal range
A.	Same	Doubled	4 times
B.	Same	4 times	4 times
C.	Doubled	4 times	Doubled
D.	Doubled	4 times	4 times

- Q.34 Which one of the following is true in the case of inelastic collision?
- |                  |                |               |
|------------------|----------------|---------------|
| Total energy     | Kinetic Energy | Momentum      |
| A. Conserved     | Conserved      | Conserved     |
| B. Conserved     | Not conserved  | Conserved     |
| C. Conserved     | Conserved      | Not conserved |
| D. Not conserved | Not conserved  | Conserved     |

- Q.35 A body is thrown with a velocity of  $9.8\text{ m/s}$  making an angle of  $30^\circ$  with the horizontal. It will hit the ground after a time
- A. 1 sec
  - B. 2 sec
  - C. 3 sec
  - D. 1.5 sec

- Q.36 If  $R = \frac{R_{\text{max}}}{2}$  then angle of projection is
- A.  $30^\circ$
  - B.  $60^\circ$
  - C.  $15^\circ$
  - D.  $45^\circ$

- Q.37 A body is thrown horizontally from the top of a tower of height 5 m. It touches the ground at a distance of 10 m from the foot of the tower. The initial velocity of the body is ( $g = 10\text{ ms}^{-2}$ )
- A.  $2.5\text{ms}^{-1}$
  - B.  $5\text{ms}^{-1}$
  - C.  $10\text{ms}^{-1}$
  - D.  $20\text{ms}^{-1}$

- Q.38 A cricket ball is hit so that it travels straight up in air and it acquires 3 seconds to reach the maximum height. Its initial velocity is
- A.  $10\text{ms}^{-1}$
  - B.  $15\text{ms}^{-1}$
  - C.  $29.4\text{ms}^{-1}$
  - D.  $12.2\text{ms}^{-1}$

- Q.39 The distance travelled by a body is proportional to the square of time. The body is moving with
- A. Uniform acceleration
  - B. Uniform velocity
  - C. Variable acceleration
  - D. All of the above

- Q.40 Two balls projected at  $30^\circ$  and  $60^\circ$  with same initial velocities. The ratio of their maximum heights is
- A. 1:2
  - B. 1:3
  - C. 1:4
  - D.  $1:\sqrt{2}$

- Q.41 A 7.0 kg bowling ball experiences a net force of  $5.0\text{ N}$  what will be its acceleration?
- A.  $7.1\text{ ms}^{-2}$
  - B.  $5.0\text{ ms}^{-2}$
  - C.  $0.71\text{ ms}^{-2}$
  - D.  $35.0\text{ ms}^{-2}$

- Q.42 Two projectiles are fired at different angles with the same magnitude of velocity such that they have the same range. At what angles they might have been projected?
- A.  $35^\circ$  and  $75^\circ$
  - B.  $10^\circ$  and  $50^\circ$
  - C.  $25^\circ$  and  $65^\circ$
  - D. None of the above


- Q.43 A train takes 1 hour to go from one station to the other. It travels at a speed of  $30\text{ kmh}^{-1}$  for first half hour and at a speed of  $50\text{ kmh}^{-1}$  for the next half hour. The average speed of the train is:
- A.  $45\text{ kmh}^{-1}$
  - B.  $40\text{ kmh}^{-1}$
  - C.  $35\text{ kmh}^{-1}$
  - D.  $30\text{ kmh}^{-1}$

- Q.44 If a projectile is thrown with  $19.6\text{m/s}$  velocity at  $30^\circ$  with x-axis, time taken to reach highest point?
- A. 1 sec
  - B. 3 sec
  - C. 2 sec
  - D. 4 sec

- Q.45 The distance travelled is given by
- A. Area under speed-time graph
  - B. Area under distance-time graph
  - C. Slope of velocity-time graph
  - D. Slope of distance-time graph

- Q.46 If a body starts from a point, and returns back to the same point, then its
- A. Average velocity is zero but not average speed
  - B. Average speed is zero but not average velocity
  - C. Both average velocity and average speed are not zero
  - D. Both average speed and average velocity are zero

- Q.47 The variation of force acting on a body with time is shown



- What is the change in momentum of body after 4s?
- A. 10Ns
  - B. 40Ns
  - C. 20Ns
  - D. 80Ns

- Q.48 Velocity and acceleration are in the same direction when
- A. Velocity of a car is increasing on a straight road
  - B. Car is turning round a corner
  - C. Velocity of a car is decreasing on a straight road
  - D. None of these


- Q.49 When the average velocity of a moving body is equal to its instantaneous velocity then it is moving with
- A. Uniform velocity
  - B. Uniform acceleration
  - C. Variable velocity
  - D. Variable acceleration

Unit-1

- Q.50** If the initial speed of a projectile is doubled.  
 A. Its range will double  
 B. Its range will quadruple  
 C. Its range will be decreased by a factor of two  
 D. Its range will decrease by a factor of four
- Q.51** In straight line motion the  
 A. Acceleration is parallel (or antiparallel) to the velocity  
 B. Acceleration is vertical, while the velocity can be in any direction  
 C. Acceleration is perpendicular to the velocity  
 D. Acceleration is vertical and the velocity is horizontal
- Q.52** In projectile motion the  
 A. Acceleration is parallel (or antiparallel) to the velocity  
 B. Acceleration is vertical, while the velocity can be in any direction  
 C. Acceleration is perpendicular to the velocity  
 D. Acceleration is vertical and the velocity is horizontal
- Q.53** A baseball is thrown vertically into the air. The acceleration of the ball at its highest point is;  
 A. Zero  
 B.  $g$ , down  
 C.  $g$ , up  
 D.  $2g$ , down
- Q.54** A racing car traveling with constant acceleration increases its speed from  $10 \text{ m/s}$  to  $50 \text{ m/s}$  over a distance of  $60 \text{ m}$ . How long does this take?  
 A.  $2.0 \text{ s}$   
 B.  $5.0 \text{ s}$   
 C.  $4.0 \text{ s}$   
 D.  $8.0 \text{ s}$
- Q.55** An aeroplane is flying horizontally at a velocity  $v$ . It drops a packet from a height  $h$ . the time taken by the packet to reach the ground will be  
 A.  $\sqrt{\frac{2h}{g}}$   
 B.  $\sqrt{\frac{2v}{g}}$   
 C.  $\sqrt{\frac{h}{2g}}$   
 D.  $\sqrt{\frac{v}{h}}$
- Q.56** A particle is moving with a constant speed along straight-line path. A force is not required to  
 A. Change its direction  
 B. Increase its speed  
 C. Decrease its momentum  
 D. Keep it moving with uniform velocity
- Q.57** Two particles are moving with velocities  $v_1$  and  $v_2$ . Their relative velocity is maximum when the angle between their velocities  
 A. zero  
 B.  $\pi/2$   
 C.  $\pi$   
 D.  $\pi/4$
- Q.58** Horizontal distance travelled by a ball if it's thrown with initial velocity of  $20 \text{ m s}^{-1}$  at an angle of  $30^\circ$  is  
 A.  $24 \text{ m}$   
 B.  $56 \text{ m}$   
 C.  $35.3 \text{ m}$   
 D.  $36.3 \text{ m}$

Unit-1

- Q.59** A block of mass  $1 \text{ kg}$  slides along a frictionless surface at  $10 \text{ ms}^{-1}$  and it collides with a stationary block of mass  $10 \text{ kg}$ . After the collision, the first block rebounds at  $5 \text{ ms}^{-1}$  and the other one moves off at  $1.5 \text{ ms}^{-1}$ .



Identify which row in the table that correctly describes the collision.

	Momentum	Kinetic Energy	Type of Collision
A	conserved	conserved	elastic
B	conserved	not conserved	inelastic
C	conserved	not conserved	elastic
D	not conserved	not conserved	inelastic


- Q.60** A body is thrown with the velocity  $(4i + 3j) \text{ m/s}$ . Its maximum height is ( $g=10 \text{ m/s}^2$ )  
 A.  $2.5 \text{ m}$   
 B.  $0.8 \text{ m}$   
 C.  $0.9 \text{ m}$   
 D.  $0.45 \text{ m}$

# ANSWER KEY


1	A	11	D	21	C	31	A	41	C	51	A
2	B	12	D	22	D	32	C	42	C	52	B
3	A	13	D	23	A	33	D	43	B	53	B
4	C	14	B	24	C	34	B	44	A	54	A
5	A	15	D	25	C	35	A	45	A	55	A
6	D	16	C	26	D	36	C	46	A	56	D
7	A	17	C	27	D	37	C	47	B	57	C
8	B	18	C	28	D	38	C	48	A	58	C
9	A	19	C	29	A	39	A	49	A	59	B
10	C	20	B	30	B	40	B	50	B	60	D




- Q.1 Proton, electron, neutron and particles have the same momentum. Which of them have highest K.E?  
 A. Proton  
 B. Electron  
 C. Neutron  
 D. alpha-particle
- Q.2 What is the Average Power Required to Lift a Mass of 100kg to a Height of 50m in 50 Seconds?  
 A. 100  
 B. 980  
 C. 50  
 D. 5000
- Q.3 Which of the following graphs best represents graphical relation between momentum  $P$  and kinetic energy  $K$  for a body in motion?




A.



B.




C.




D.

- Q.4 A motorbike engine can develop a power of 90000 W in order to keep a constant velocity of 30 m/s. What is the pushing force?  
 A. 3000 N  
 B. 30000 N  
 C. 300000 N  
 D. 300 N
- Q.5 If a stone of mass 500 g strikes the ground with velocity of  $20 \text{ ms}^{-1}$  then kinetic energy at that point is  
 A. 10 J  
 B. 40 J  
 C. 75 J  
 D. 100 J
- Q.6 Force  $F$  on a particle moving in a straight line varies with distance  $d$  as shown in the figure. The work done on the particle during its displacement from 3m to 12m




- A. 13 J  
 B. 26 J  
 C. 18 J  
 D. 21 J
- Q.7 Two bodies with kinetic energies in the ratio 4 : 1 are moving with equal linear momentum. The ratio of their masses is:  
 A. 1 : 2  
 B. 1 : 4  
 C. 1 : 1  
 D. 4 : 1
- Q.8 How much work must be done by a force on 50 kg body in order to accelerate it from rest to 20 m/s in 10 s?  
 A.  $2 \times 10^3 \text{ J}$   
 B.  $10^3 \text{ J}$   
 C.  $10^4 \text{ J}$   
 D.  $4 \times 10^4 \text{ J}$
- Q.9 A force applied by an engine of a train of mass  $2.05 \times 10^6 \text{ kg}$  changes its velocity from 5m/s to 25m/s in 5 minutes. The power of the engine is  
 A. 6 MW  
 B. 1.025 MW  
 C. 5 MW  
 D. 2.05 MW

- Q.10 When a 300 N force is applied to a box weighing 600 N, the box moves 3.0 m horizontally in 20 s. What is the average power?



- A. 45 W  
 B. 90 W  
 C. 900 W  
 D. 1800 W
- Q.11 An engine pumps out 50 kg of water. If the water comes out vertically upwards with a velocity of 20 m/s. The power of engine is (take  $g = 10 \text{ m s}^{-2}$ )  
 A. 10 KW  
 B. 20 kW  
 C. 1000 W  
 D. 10 MW
- Q.12 A body, constrained to move in the  $y$ -direction, is subjected to a force of  $10 \text{ m}$  along the  $y$ -axis is  
 $F = -2\hat{i} + 15\hat{j} + 6\hat{k}$  Newton's. The work done by this force in moving the body a distance  
 A. 190 J  
 B. 160 J  
 C. 150 J  
 D. 20 J
- Q.13 Two particles of masses  $m$  and  $4m$  have linear momenta in the ratio of 2 : 1. What is the ratio of their kinetic energies?  
 A.  $\sqrt{2}$   
 B. 2  
 C. 4  
 D. 16
- Q.14 The long pendulum shown is drawn aside until the ball has risen 0.50 m. It is then given an initial speed of 3.0 m/s. The speed of the ball at its lowest position is:



- A. Zero  
 B. 0.89 m/s  
 C. 3.1 m/s  
 D. 4.3 m/s
- Q.15 Two men with weights in the ratio 5 : 3 run up a staircase (of same height) in times in the ratio 11 : 9. The ratio of power of first to that of second is  
 A. 11/15  
 B. 11/9  
 C. 15/11  
 D. 9/11
- Q.16 A body of mass  $m$  is dropped from a height  $h$  above the ground. The velocity  $v$  of the body when it has lost half its initial potential energy is given by  
 A.  $v = \sqrt{gh}$   
 B.  $v = \sqrt{2gh}$   
 C.  $v = \sqrt{\frac{gh}{2}}$   
 D.  $v = 2\sqrt{gh}$
- Q.17 A moving body need not have  
 A. Velocity  
 B. Momentum  
 C. Potential energy  
 D. Kinetic energy.

- Q.18 According to work energy principle work done on body will equal to change its  
 A. K.E only  
 B. P.E only  
 C. K.E and P.E both  
 D. All may correct
- Q.19 Adjacent figure shows the force-displacement graph of a moving body, the work done in displacing body from  $x=0$  to  $x=35\text{m}$  is equal to



- A. 25 J  
 B. 200 J  
 C. 287.5 J  
 D. 50 J
- Q.20 A body of mass 2 kg is projected vertically upwards with a velocity of  $2\text{msec}^{-1}$ . The K.E. of the body just before striking the ground is  
 A. 2J  
 B. 4J  
 C. 1J  
 D. 8J
- Q.21 A particle of mass  $m$  at rest is acted upon by a force  $F$  for a time  $t$ . Its K.E. after an interval  $t$  is  
 A.  $\frac{F^2 t^2}{m}$   
 B.  $\frac{F^2 t^2}{3m}$   
 C.  $\frac{F^2 t^2}{2m}$   
 D.  $\frac{Ft}{2m}$
- Q.22 An electric motor exerts a force of 40 N on a cable and pulls it by a distance of 30 m in one minute. The power supplied by the motor (in Watts) is  
 A. 200  
 B. 20  
 C. 10  
 D. 2
- Q.23 A weight lifter lifts 300 kg from the ground to a height of 2 meter in 3 second. The average power generated by him is  
 A. 5880 watt  
 B. 4410 watt  
 C. 2205 watt  
 D. 1960 Watt
- Q.24 Power of a water pump is 2 kW. If  $g=10\text{m/sec}^2$ , the amount of water it can raise in one minute to a height of 10 m is  
 A. 2000 litre  
 B. 1000 litre  
 C. 1200 litre  
 D. 100 litre
- Q.25 A car of mass 1000 kg accelerates uniformly from rest to a velocity of 54 km/hour in 5s. The average power of the engine during this period in watts is  
 A. 2000 W  
 B. 22500 W  
 C. 5000 W  
 D. 2250 W
- Q.26 A boy holds a 400 N weight at arm's length for 10s. His arm is 1.5m above the ground. The work done by the force of the boy on the weight while he is holding it is:  
 A. 0  
 B. 900J  
 C. 600J  
 D. 400J

- Q.27 A man does a given amount of work in 19 sec. Another man does the same amount of work in 20 sec. The ratio of the output power of first man to the second man is  
 A. 2/1  
 B. 1/2  
 C. 1  
 D. 4/1
- Q.28 A man of weight 500 newtons walks up to the top of a building of height of 20 m above the street level. The increase in the potential energy of man is  
 A.  $5 \times 9.8 \text{ J}$   
 B.  $500 \times 20 \text{ J}$   
 C.  $500 \times 9.8 \text{ J}$   
 D.  $\frac{500 \times 20}{9.8} \text{ J}$
- Q.29 Kinetic energy of a body moving with speed of  $10 \text{ ms}^{-1}$  is 30 J. If its speed becomes  $30 \text{ ms}^{-1}$  then its K.E becomes  
 A. 10 J  
 B. 90 J  
 C. 270 J  
 D. 180 J
- Q.30 A car is driven along a level road. The total energy input from the petrol is 60kJ and the car wastes 45kJ of energy, what is the efficiency of car  
 A. 25%  
 B. 15%  
 C. 45%  
 D. 75%
- Q.31 A 50kg man with 20kg load on his head climbs up 20 steps of 0.25m height each. The work done in climbing is?  
 A. 100 J  
 B. 350 J  
 C. 3430 J  
 D. 5 J
- Q.32 If force and displacement of particle in direction of force are doubled. Work would be  
 A. Double  
 B. 1/4 times  
 C. Half  
 D. 4 times
- Q.33 A person walks 2m with an acceleration of  $5\text{ms}^{-2}$ , holding an object of mass 2kg. The work done on the object is  
 A. 20 J  
 B. 10 J  
 C. 5 J  
 D. 0 J
- Q.34 The work done on an object does not depend upon the  
 A. Displacement  
 B. Angle between force and displacement  
 C. Force applied  
 D. Initial velocity of the object
- Q.35 Which of the following is correct order?  
 A.  $1\text{kwh} > 1\text{erg} > 1\text{J}$   
 B.  $1\text{J} > 1\text{erg} > 1\text{kwh}$   
 C.  $1\text{kwh} > 1\text{J} > 1\text{erg}$   
 D.  $1\text{kwh} = 1\text{J} = 1\text{erg}$
- Q.36 You lift a suit case from the floor and keep it on a table. The work done by you on the suitcase does not depend upon  
 A. The path taken by the suitcase  
 B. The time taken by you in doing work  
 C. Weight of the suitcase  
 D. A and B
- Q.37 A particle moves with velocity  $6\hat{i} - 4\hat{j} + 3\hat{k} \text{ ms}^{-1}$  under the influence of constant force  $\vec{F} = 20\hat{i} + 15\hat{j} - 5\hat{k} \text{ N}$ . The instantaneous power applied to particle is  
 A.  $45 \text{ Js}^{-1}$   
 B.  $35 \text{ Js}^{-1}$   
 C.  $25 \text{ Js}^{-1}$   
 D.  $195 \text{ Js}^{-1}$
- Q.38 A ball is dropped from a height of 10 m.  
 A. Its potential energy increases and kinetic energy decreases during the falls  
 B. Its potential energy is equal to the kinetic energy during the fall.  
 C. The potential energy decreases and the kinetic energy increases during the fall.  
 D. The potential energy and kinetic energy is maximum while it is falling.

- Q.39 How much time will be required to perform 520 J of work at the rate of 20 W?  
 A. 24s  
 B. 16s  
 C. 20s  
 D. 26s
- Q.40 Output of a truck is 4500 J and its efficiency is 50%. Input energy provided to truck is  
 A. 5000 J  
 B. 9000 J  
 C. 900 J  
 D. 500 J
- Q.41 Due to application of 5 N force an object moves 10 meter along perpendicular direction of the force. What amount work is done?  
 A. 50 J  
 B. 5 J  
 C. 15 J  
 D. 0 J
- Q.42 Work is independent of:  
 A. Force  
 B. Time  
 C. Displacement  
 D. All of these
- Q.43 When the speed of object is halved and the mass is quadrupled then the kinetic energy is:  
 A. Quartered  
 B. One Third  
 C. Twice  
 D. Remain same
- Q.44 For a body moving in a circular path, the work done by the centripetal force is  
 A. Negative  
 B. Constant  
 C. Positive  
 D. Zero
- Q.45 If a pump can lift 200 kg of water through a height of 6 m in 10 seconds, then its power is  
 A. 1100 watts  
 B. 1300 watts  
 C. 1000 watts  
 D. 1200 watts
- Q.46 A light and a heavy body have equal momenta. Which one has greater K.E.  
 A. The light body  
 B. The K.E. are equal  
 C. The heavy body  
 D. Data is incomplete
- Q.47 Two bodies of masses 2m and 1m have their K.E. in the ratio 8 : 1, then their ratio of momenta is  
 A. 1 : 1  
 B. 4 : 1  
 C. 2 : 1  
 D. 8 : 1
- Q.48 The decrease in the potential energy of a ball of mass 20 kg which falls from a height of 50 cm is  
 A. 968 J  
 B. 1980 J  
 C. 98 J  
 D. None of these
- Q.49 A body moves a distance of 10 m along a straight line under the action of a force of 5 N. If the work done is 25 joules, the angle which the force makes with the direction of motion of the body  
 A. 0°  
 B. 60°  
 C. 30°  
 D. 90°
- Q.50 A force  $F = (5\hat{i} + 3\hat{j})$  newton is applied over a particle which displaces it from its origin to the point  $r = (2\hat{i} - 1\hat{j})$  metres. The work done on the particle is  
 A. -7 joules  
 B. +7 joules  
 C. +13 joules  
 D. +11 joules
- Q.51 What happens to the kinetic energy of a moving object if the net work done is positive?  
 A. The kinetic energy increases  
 B. The kinetic energy remains the same  
 C. The kinetic energy decreases  
 D. The kinetic energy is zero

- Q.52 If momentum is increased by two times K.E. increases by  
 A. Two times  
 B. 3 times  
 C. four times  
 D. Remains
- Q.53 A particle moves with  $\vec{v} = -3\hat{j} + 5\hat{i} + 6\hat{k}$  ms<sup>-1</sup> under  $\vec{F} = 10\hat{i} + 10\hat{j} + 20\hat{k}$  N. the power applied  
 A. 200 Js<sup>-1</sup>  
 B. 170 Js<sup>-1</sup>  
 C. 40 Js<sup>-1</sup>  
 D. 140 Js<sup>-1</sup>
- Q.54 Effort × distance through which effort acts =  
 A. Output of machine  
 B. Input of machine  
 C. Efficiency  
 D. Work
- Q.55 Load × distance covered by the load =  
 A. Output of machine  
 B. Input of machine  
 C. Efficiency  
 D. Work
- Q.56 An object is thrown straight up. Which of the following is true about the sign of work done by the gravitational force while the object moves up and then down?  
 A. Work is positive on the way up, work is positive on the way down  
 B. Work is negative on the way up, work is positive on the way down  
 C. Work is negative on the way up, work is negative on the way down  
 D. Work is positive on the way up, work is negative on the way down
- Q.57 The burning of coal, wood or natural gas and releasing of heat and light energy in the air is a  
 A. nuclear reaction  
 B. kinetic reaction  
 C. chemical reaction  
 D. potential reaction
- Q.58 If a stone of mass 500 g is dropped from height 20m then kinetic energy while hitting the ground  
 A. 10 J  
 B. 40 J  
 C. 75 J  
 D. 100 J
- Q.59 If the linear momentum of the object is increased by 0.1%, then the kinetic energy is increased by  
 A. 0.1 %  
 B. 0.2%  
 C. 0.4%  
 D. 0.01%
- Q.60 A machine which has an efficiency of 80% raises a load of 50 N through a vertical height of 10 m. The work input to the machine is:  
 A. 400 J  
 B. 500 J  
 C. 800 J  
 D. 625 J

## ANSWER KEY

1	B	11	A	21	C	31	C	41	D	51	A
2	B	12	C	22	B	32	D	42	B	52	C
3	D	13	D	23	D	33	A	43	D	53	D
4	A	14	D	24	C	34	D	44	D	54	B
5	D	15	C	25	D	35	C	45	D	55	A
6	A	16	A	26	A	36	D	46	A	56	C
7	B	17	C	27	A	37	A	47	B	57	C
8	C	18	D	28	B	38	C	48	C	58	D
9	D	19	C	29	C	39	D	49	B	59	B
10	A	20	B	30	A	40	B	50	B	60	D

# 3 UNIT

## ROTATIONAL AND CIRCULAR MOTION


### SELF ASSESSMENT TEST

- Q. 1 When a body is whirled in a horizontal circle by means of a string the centripetal force is supplied by  
 A. Mass of a body  
 B. Velocity of body  
 C. The tension in the string  
 D. Centripetal acceleration
- Q. 2 If a wheel of radius  $r$  turns through an angle of  $30^\circ$  then the distance through which any point on its rim moves is  
 A.  $\pi/3 \times r$   
 B.  $\pi/6 \times r$   
 C.  $\pi/30 \times r$   
 D.  $\pi/180 \times r$
- Q. 3 The ratio of the angular speed of the minute hand of a clock to that of its hour hand is  
 A. 3600 : 1  
 B. 60 : 1  
 C. 24 : 1  
 D. 12 : 1
- Q. 4 A body of mass 10 kg, is moving with a velocity of 5 m/s in a circle of radius 5 m, what is the centripetal acceleration of the body?  
 A.  $5\text{m/s}^2$   
 B.  $25\text{m/s}^2$   
 C.  $0.5\text{m/s}^2$   
 D.  $50\text{m/s}^2$
- Q. 5 A particle moves on a circular path with decreasing speed. Choose the correct statement.  
 A. Angular momentum remains constant  
 B. Acceleration is towards the center  
 C. Particle moves on a spiral path with decreasing radius  
 D. The direction of angular momentum remains constant
- Q. 6 A disc is rotating about an axis through its centre and perpendicular to its plane. A point P on the disc is twice as far from the axis as a point Q. At a given instant, what is the value of the  $\frac{\text{the linear velocity of P}}{\text{the linear velocity of Q}}$ ?  
 A. 4  
 B. 2  
 C.  $\frac{1}{2}$   
 D.  $\frac{1}{4}$
- Q. 7 An angular ring with inner and outer radii  $R_1$  and  $R_2$  is rolling without slipping with a uniform angular speed. The ratio of the forces experienced by the two particles situated on the inner and outer parts of the ring,  $F_1 / F_2$  is  
 A. 1  
 B.  $\frac{R_1}{R_2}$   
 C.  $\frac{R_2}{R_1}$   
 D.  $\left(\frac{R_2}{R_1}\right)^2$
- Q. 8 If a body of mass  $m$  is rotating in a circle of radius  $r$  with frequency of rotation " $f$ " then centripetal force acting on it is  
 A.  $2\pi mrf$   
 B.  $4\pi^2 mrf$   
 C.  $4\pi^2 mrf^2$   
 D.  $\pi^2 mrf^2$

### Unit-3

## Rotational and Circular Motion

- Q. 9 A body is rotating clockwise with decreasing angular velocity. Its angular acceleration is directed  
 A. Into the plane of paper  
 B. Along the radius  
 C. Out of the plane of paper  
 D. Along the tangent to the circle
- Q. 10  $r\omega^2$  has unit of  
 A. N  
 B.  $\text{ms}^{-1}$   
 C.  $\text{ms}^{-2}$   
 D.  $\text{s}^{-1}$
- Q. 11 The figure shows a cylinder of radius 0.7m rotating about its axis at 10rad/s. The speed of the point P is:



- A. 7.0m/s  
 C.  $7.0\pi$  rad/s  
 B.  $14\pi$  rad/s  
 D. 0.70m/s
- Q. 12 Centripetal acceleration can be expressed as  
 A.  $\frac{v^2}{r}$   
 B.  $r\omega^2$   
 C.  $v\omega$   
 D. All of these
- Q. 13 In equation  $\vec{v} = \vec{\omega} \times \vec{r}$   $90^\circ$  is angle between  
 A.  $\vec{v}$  and  $\vec{r}$   
 B.  $\vec{r}$  and  $\vec{\omega}$   
 C.  $\vec{v}$  and  $\vec{\omega}$   
 D. All are correct
- Q. 14 What is the speed of the tip of second's hand of a clock if its length is 10 cm  
 A.  $1.05\text{cms}^{-1}$   
 B.  $2.05\text{cms}^{-1}$   
 C.  $1.05\text{m/s}$   
 D.  $3.05\text{cms}^{-1}$
- Q. 15 A particle of rigid body is at a distance 0.1 m from axis of rotation to rotate with linear speed 3 m/s. What is angular speed of the rigid body  
 A. 0.3 rad/s  
 B. 30 rads $^{-1}$   
 C. 3 rad s $^{-1}$   
 D. 1.5 rad/s
- Q. 16 If E is the K.E of body moving in circle of radius  $r$  then the centripetal force may be written as  
 A.  $F_c = \frac{K.E}{2r}$   
 B.  $F_c = \frac{K.E}{2r^2}$   
 C.  $F_c = K.E \times 2r$   
 D.  $\frac{2K.E}{r}$
- Q. 17 The ratio of displacement along diameter and total distance along circle is:  
 A.  $\pi : 1$   
 B.  $\pi : 2$   
 C.  $2 : \pi$   
 D.  $1 : \pi$
- Q. 18 A ball of mass 0.25 kg attached to the end of a string of length 1.96 m is moving in a horizontal circle. The string will break if the tension is more than 25 N. What is the maximum speed with which the ball can be moved?  
 A. 14 m/s  
 B. 3 m/s  
 C. 3.92 m/s  
 D. 5 m/s

Unit-3

- Q. 19 What happens to the centripetal acceleration of a revolving body if you double the orbital speed  $v$  and half the angular velocity  $\omega$
- The centripetal acceleration remains unchanged
  - The centripetal acceleration is halved
  - The centripetal acceleration is doubled
  - The centripetal acceleration is quadrupled
- Q. 20 A wheel is at rest. Its angular velocity increases uniformly and becomes 60 rad/sec after 5 sec. The total angular displacement is
- 600 rad
  - 300 rad
  - 75 rad
  - 150 rad
- Q. 21 A car of mass 1000 kg is moving with speed 72 km/h in a circular track of radius 100 m. The centripetal force acting on it is
- 4 N
  - 400 N
  - 40 N
  - 4000 N
- Q. 22 A circle of radius 1m rolls through some distance making an angle  $180^\circ$  at the centre; find the distance:
- 3.14 m
  - 3.14 rad
  - 5m
  - 2.8 m
- Q. 23 A particle comes round a circle of radius 1 m once. The time taken by it is 10 sec. The average velocity of motion is
- $0.2 \pi \text{ m/s}$
  - $2 \pi \text{ m/s}$
  - $2 \text{ m/s}$
  - Zero
- Q. 24 The acceleration of a train travelling with speed of 400 m/s as it goes round a curve of radius 160 m, is
- $1 \text{ km/s}^2$
  - $100 \text{ m/s}^2$
  - $10 \text{ m/s}^2$
  - $1 \text{ m/s}^2$
- Q. 25 An aircraft executes a horizontal loop of radius 1 km with steady speed of 900 km/h. What is its centripetal acceleration?
- $250 \text{ km/s}^2$
  - $75 \text{ m/s}^2$
  - $62.5 \text{ m/s}^2$
  - $60 \text{ m/s}^2$
- Q. 26 If the body is moving in a circle of radius  $r$  with a constant speed  $v$ , its angular velocity is
- $v^2/r$
  - $vr$
  - $v/r$
  - $r/v$
- Q. 27 Centripetal force may be equal to
- $\frac{mv^2}{r}$
  - $\frac{pv}{r}$
  - $\frac{p^2}{mr}$
  - All of these
- Q. 28 What is outward force acting on a mass of 10 kg when rotating at one end on an inelastic string 10m long at speed of 1m/s?
- 1 N
  - 2 N
  - 10 N
  - 100 N
- Q. 29 The angle subtended by an arc equal to radius is
- 1 rad
  - One degree
  - 1 Revolution
  - All of these

Unit-3

- Q. 30 If a particle moves in a circle describing equal angles in equal times, its velocity vector
- Remains constant
  - Changes in direction
  - Changes in magnitude
  - Changes both in magnitude and direction
- Q. 31 A motor cyclist going round in a circular track at constant speed has
- Constant linear velocity
  - Constant acceleration
  - Constant angular velocity
  - Constant force
- Q. 32 Which of the following statements is false for a particle moving in a circle with a constant angular speed?
- The velocity vector is tangent to the circle
  - The acceleration vector is tangent to the circle
  - The acceleration vector points to the centre of the circle
  - The velocity and acceleration vectors are perpendicular to each other
- Q. 33 The number of revolutions in  $3\pi$  radians
- 2
  - 3
  - $3/2$
  - $1/2$
- Q. 34 A Wheel of radius 50 cm having an angular speed of 5 rad  $\text{s}^{-1}$  have linear speed
- $1.5 \text{ ms}^{-1}$
  - $3.5 \text{ ms}^{-1}$
  - $4.5 \text{ ms}^{-1}$
  - $2.5 \text{ ms}^{-1}$
- Q. 35 In circular motion, if the angular velocity and angular acceleration becomes parallel, then the motion becomes:
- Slower
  - Faster
  - Constant
  - Both 'A' and 'C'
- Q. 36 An object is moving in a circle of radius 100 m with a constant speed of 31.4 m/s. What is its average speed for one complete revolution?
- Zero
  - 3.14 m/s
  - 31.4 m/s
  - $\sqrt{2} \times 31.4 \text{ m/s}$
- Q. 37  $1 \frac{\text{rev}}{\text{min}}$  is equal to:
- $\frac{\pi}{6} \text{ rad s}^{-1}$
  - $\frac{\pi}{15} \text{ rad s}^{-1}$
  - $\frac{\pi}{20} \text{ rad s}^{-1}$
  - $\frac{\pi}{30} \text{ rad s}^{-1}$
- Q. 38 A wheel rotates about an axis passing through the center and perpendicular to the plane with slowly increasing angular speed. Then it has
- Radial velocity and radial acceleration
  - Tangential velocity and radial acceleration
  - Tangential velocity and tangential acceleration
  - Tangential velocity but acceleration having both components
- Q. 39 A body moves in a circle of radius 4 m with constant speed 8 m/s experiences centripetal force 128 N. What is the mass of body?
- 2 Kg
  - 8 Kg
  - 4 Kg
  - 16 Kg
- Q. 40 When a body is whirled in a horizontal circle by means of a string the centripetal force is supplied by
- Mass of body
  - Tension in the string
  - Velocity of body
  - Centripetal acceleration

- Q. 41 Two bodies of mass 10 kg and 5 kg moving in concentric orbits of radii  $R$  and  $r$  such that their periods are the same. Then the ratio between their centripetal acceleration is  
 A.  $R:r$   
 B.  $R^2:r^2$   
 C.  $r:R$   
 D.  $r^2:R^2$
- Q. 42 The force which can do no work on the body on which it acts:  
 A. Frictional force  
 B. Elastic force  
 C. Gravitational force  
 D. Centripetal force
- Q. 43 A string breaks if its tension exceeds 10 newtons. A stone of mass 250 gm tied to this string of length 10 cm is rotated in a horizontal circle. The maximum angular velocity of rotation can be  
 A. 20 rad/s  
 B. 40 rad/s  
 C. 100 rad/s  
 D. 200 rad/s
- Q. 44 An electric fan has blades of length 30 cm as measured from the axis of rotation. If the fan is rotating at 1200 r.p.m. The acceleration of a point on the tip of the blade is about  
 A. 1600 m/sec<sup>2</sup>  
 B. 4740 m/sec<sup>2</sup>  
 C. 2370 m/sec<sup>2</sup>  
 D. 5055 m/sec<sup>2</sup>
- Q. 45 The expression for centripetal force is given by:  
 A.  $m\omega r^2$   
 B.  $\frac{m^2 v^2}{r^2}$   
 C.  $\frac{m^2 v^2}{r}$   
 D.  $\frac{mv^2}{r^2}$
- Q. 46 The period of circular motion is  
 A.  $T = \frac{2\pi}{\omega}$   
 B.  $T = 2\pi\omega$   
 C.  $T = \frac{\omega}{2\pi}$   
 D.  $T = \frac{2\pi\omega}{2}$
- Q. 47 The curved flight of fighter planes at high speed requires a large  
 A. Gravitational force  
 B. Frictional force  
 C. Centripetal force  
 D. Centrifugal acceleration
- Q. 48 Two particles of equal masses are revolving in circular paths of radius  $r_1$  and  $r_2$  respectively with the same speed. The ratio of their centripetal force is  
 A.  $\left(\frac{r_2}{r_1}\right)$   
 B.  $\sqrt{\left(\frac{r_2}{r_1}\right)}$   
 C.  $\left(\frac{r_2}{r_1}\right)^2$   
 D.  $\left(\frac{r_1}{r_2}\right)^2$
- Q. 49 In uniform circular motion, the factor that remains constant is  
 A. Linear velocity  
 B. Centripetal force  
 C. Acceleration  
 D. Speed

- Q. 50 A particle moves in a circle of radius 25cm at two revolutions per second. The radial acceleration of the particle is  
 A.  $\pi^2$   
 B.  $8\pi^2$   
 C.  $4\pi^2$   
 D. Zero
- Q. 51 A fly wheel rotates at a constant speed of 3000rpm. The angle described by the shaft in radian in one second is:  
 A.  $3000\pi$   
 B.  $50\pi$   
 C.  $100\pi$   
 D.  $2\pi$
- Q. 52 A stone is whirled in a vertical plane. The stone has  
 A. Radial acceleration only  
 B. Tangential acceleration only  
 C. Both radial and tangential accelerations  
 D. Neither radial nor tangential acceleration
- Q. 53 A body is rotating in circle of radius  $r$ . Keeping period of rotation constant but radius is doubled ( $2r$ ) then centripetal force become  
 A. Half  
 B. Same  
 C. Double  
 D. Four times
- Q. 54 A particle is moving in a circle of radius ' $r$ '. In one period of revolution its displacement and distance covered are  
 A.  $2\pi r, 2\pi r$   
 B.  $\pi, 2\pi r$   
 C.  $2\pi, 2\pi r$   
 D. zero,  $2\pi r$
- Q. 55 A particle is acted upon by a force of constant magnitude which is always perpendicular to the velocity of the particle. The motion takes place in a plane. It follows that  
 A. Its velocity is constant  
 B. Its acceleration is constant  
 C. Its motion is linear  
 D. Its motion is circular
- Q. 56 A particle describes a circular path of diameter 20 m every 2s. the average angular speed of the particle during 4s is  
 A.  $20/\pi$  rad/s  
 B.  $10/\pi$  rad/s  
 C.  $5/\pi$  rad/s  
 D.  $\pi$  rad/s
- Q. 57 A point on the wheel of 400cm diameter has a velocity of 1600cm/s. The angular velocity of the wheel is  
 A. 8 rad/s  
 B. 4 rad/s  
 C. 12 rad/s  
 D. 2 rad/s
- Q. 58 Four particles have the following masses (in terms of  $m$ ), speeds (in terms of  $v$ ), and radii (in terms of  $r$ ). Which particle have the same centripetal force as particle-1?

Particle	Mass	Speed	Radius
1	$m$	$v$	$r$
2	$m/2$	$2v$	$2r$
3	$2m$	$v/2$	$r$
4	$1m$	$2v$	$3r$

- A. Particle 1  
 C. Particle 3

- B. Particle 2  
 D. Particle 4

Unit-3

- Q. 59 Which of the following changes when a body performs uniform circular motion?  
 A. Speed  
 C. Direction  
 B. Mass  
 D. Kinetic energy
- Q. 60 A body is moving in a circular path with acceleration 'a'. If the speed of the body is increased to four times the initial value, the acceleration will become:  
 A. 4a  
 C. 16a  
 B. 8a  
 D. a

## ANSWER KEY


1	C	11	A	21	D	31	C	41	A	51	C
2	B	12	D	22	A	32	B	42	D	52	C
3	D	13	D	23	D	33	C	43	A	53	C
4	A	14	A	24	A	34	D	44	B	54	D
5	D	15	B	25	C	35	B	45	A	55	D
6	B	16	D	26	C	36	C	46	A	56	D
7	B	17	D	27	D	37	D	47	C	57	A
8	C	18	A	28	A	38	D	48	A	58	B
9	C	19	A	29	A	39	B	49	D	59	C
10	C	20	D	30	C	40	B	50	C	60	C

# 4 UNIT

## WAVES


### SELF ASSESSMENT

- Q. 1 The velocity of sound is greatest in  
 A. Water  
 C. Copper  
 B. Air  
 D. Ammonia
- Q. 2 Silence zone takes place due to  
 A. Constructive interference  
 C. Beats  
 B. Destructive interference  
 D. Resonance
- Q. 3 If the length of a simple pendulum is increased by 2%, then the time period  
 A. Increases by 1%  
 C. Increases by 2%  
 B. Decreases by 1%  
 D. Decreases by 2%
- Q. 4 The phase difference between the acceleration of a particle executing simple harmonic motion and the instantaneous velocity is?  
 A.  $\pi$   
 C. Zero  
 B.  $0.707\pi$   
 D.  $0.5\pi$
- Q. 5 Velocity of sound in air is  
 A. Faster in dry air than in moist air  
 C. Directly proportional to pressure  
 B. Directly proportional to temperature  
 D. Slower in dry air than in moist air
- Q. 6 The length of a string is 1m, tension in it is 40N and mass of the string is 0.1 kg. Then the velocity of transverse waves produced in the string will be:  
 A.  $400 \text{ ms}^{-1}$   
 C.  $180 \text{ ms}^{-1}$   
 B.  $80 \text{ ms}^{-1}$   
 D.  $20 \text{ ms}^{-1}$
- Q. 7 In strings, the position of antinodes is obtained at  
 A.  $\lambda, 2\lambda, 3\lambda$   
 C.  $0, \frac{\lambda}{2}, \lambda$   
 B.  $2\lambda, 4, 6\lambda$   
 D.  $\frac{\lambda}{4}, \frac{3\lambda}{4}, \frac{5\lambda}{4}$
- Q. 8 A sound source rotates anti-clock wise with an angular velocity  $\omega$ . Radius of the circle is R. A person is at P. The maximum frequency is heard when position of the source is at.




- A. Y  
 C. X  
 B. Z  
 D. W
- Q. 9 The wavelength of the sound produced by a source is 0.8m. If the source moves towards the stationary listener at  $32 \text{ ms}^{-1}$ , what will be apparent wavelength of the sound? The velocity of sound is  $320 \text{ ms}^{-1}$ .  
 A. 0.80 m  
 C. 0.72 m  
 B. 0.40 m  
 D. 0.32 m
- Q. 10 A source of sound is moving towards a stationary observer with  $\frac{1}{10}$  of the speed of sound. The ratio of apparent to actual frequency of sound is:  
 A.  $\frac{10}{9}$   
 B.  $\frac{11}{9}$   
 C.  $\frac{11}{10}$   
 D.  $\frac{9}{11}$

- Q. 11 When a wave moves through 10m, a point changes from crest to trough and time taken is 1s then wavelength of wave and its frequency are  
 A. 20 m, 0.5 Hz  
 C. 10 m, 1 Hz
- Q. 12 Displacement time graph of particle executing SHM is shown. The corresponding force-time graph of particle is




- Q. 13 At which temperature will the velocity of sound at 27°C become double?  
 A. 54 °C  
 C. 927 °C
- Q. 14 An observer moves towards a stationary source of sound, with a velocity one fifth of the velocity of sound. What is the percentage increase in the apparent frequency?  
 A. zero  
 C. 5%
- Q. 15 The graph shown below the variation of displacement with time for a particle moving with simple harmonic motion. Which one of the following correctly gives the value of the frequency and the amplitude of the motion?




	Frequency / kHz	Amplitude / $\mu\text{m}$
A.	5	2
B.	25	2
C.	25	4
D.	50	2

- Q. 16 The diagram shows the profile of a wave. Which of the following pairs of points are in phase?



- A. A, B  
 C. B, D  
 B. B, C  
 D. B, E

- Q. 17 A sound source moving with  $8 \text{ ms}^{-1}$  cross a stationary observer. The ratio of apparent frequency before and after crossing the observer (speed of sound =  $332 \text{ ms}^{-1}$ )  
 A.  $\frac{85}{81}$   
 C.  $\frac{81}{85}$
- Q. 18 In a sinusoidal wave, the time required for a particular point to move from maximum displacement to zero displacement is 0.5 sec. The frequency of the wave is  
 A. 0.2 Hz  
 C. 5 Hz
- Q. 19 A stretched string resonates with fundamental frequency of 50 Hz. The wavelength for its 3<sup>rd</sup> overtone is if speed of transverse wave in the string is  $100 \text{ ms}^{-1}$   
 A. 66 cm  
 C. 50 cm
- Q. 20 The string is attached to one end of vibrator having frequency 800 Hz. What will be the speed of the waves produced in the string as shown in figure.



- A.  $200 \text{ ms}^{-1}$   
 C.  $800 \text{ ms}^{-1}$
- Q. 21 A standing wave is established in a stretched string which is 100 cm long with fundamental frequency  $f$ . If tension in the string is increased to double its fundamental frequency would become  
 A.  $2f$   
 C.  $\sqrt{2}f$
- Q. 22 The sonometer wire is vibrating in its first overtone. There are  
 A. Two nodes and two antinodes  
 C. Three nodes and two antinodes
- Q. 23 In case of a moving source of sound which is moving away from an observer  
 A. The wavelength of sound appears to be less  
 B. The wavelength of sound appears same  
 C. The wavelength of sound appears to be more  
 D. Frequency and wavelength of sound appears to be same
- Q. 24 Distance and displacement traveled by a vibrating body in a time equal to  $\frac{3}{4}T$

where  $T$  is the period of the vibration

- A.  $3x_0, 3x_0$   
 C.  $3x_0, x_0$   
 B.  $3x_0, 0$   
 D.  $2x_0, 0$



- It-4
- Q. 25 The ratio of fundamental wavelengths of one end close pipe to the both end open pipe having same lengths  
 A. 1 : 2  
 C. 2 : 1  
 B. 1 : 4  
 D. 4 : 1
- Q. 26 An organ pipe  $P_1$  closed at one end vibrating in its first overtone and another pipe  $P_2$  open at both ends vibrating in its third overtone and another pipe tuning fork. The ratio of lengths of  $P_1$  and  $P_2$  is  
 A. 8 : 3  
 C. 4 : 3  
 B. 3 : 8  
 D. 3 : 4
- Q. 27 Two stretched strings have lengths  $l$  and  $2l$  while tensions are  $T$  and  $4T$  respectively. If they are made of same material the ratio of their frequencies is  
 A. 1 : 1  
 C. 2 : 1  
 B. 1 : 2  
 D. 1 : 4
- Q. 28 If the source of sound is moving away from the stationary observer with a speed half the speed of sound then the apparent wavelength for the observer would be if speed of sound is  $v$  and its original frequency is  $f$ .  
 A.  $\frac{v}{2f}$   
 C.  $\frac{3v}{2f}$   
 B.  $\frac{5v}{2f}$   
 D.  $\frac{v}{3f}$
- Q. 29 With what speed an observer should move towards a stationary source such that apparent frequency is double the actual frequency ( $v$  is speed of sound waves)  
 A.  $v$   
 C.  $2v$   
 B.  $\frac{v}{2}$   
 D.  $4v$
- Q. 30 For a closed organ pipe resonance is occurred when air columns of lengths are equal to  
 A.  $\frac{\lambda}{1}, \frac{\lambda}{2}, \lambda$   
 C.  $\frac{\lambda}{2}, \lambda, 3\lambda$   
 B.  $\frac{\lambda}{2}, \frac{3\lambda}{2}, \frac{5\lambda}{2}$   
 D.  $\frac{\lambda}{4}, \frac{3\lambda}{4}, \frac{5\lambda}{4}$
- Q. 31 The waves propagating on water surface are  
 A. Ultrasonic  
 C. Longitudinal  
 B. Unaudible  
 D. Transverse
- Q. 32 The wavelength of sound in air is 10 cm. Its frequency is (Take velocity of sound =  $330 \text{ ms}^{-1}$ )  
 A. 3.3 kHz  
 C. 330 mHz  
 B. 330 Hz  
 D.  $3 \times 10^9 \text{ Hz}$
- Q. 33 A wave generator produces 500 pulses in 10 seconds. Find period of pulses it produces  
 A. 50s  
 C.  $\frac{1}{5} \text{ s}$   
 B.  $\frac{1}{50} \text{ s}$   
 D.  $\frac{10}{50} \text{ s}$

- Unit-4
- Q. 34 The speed of a wave on a particular string is  $24 \text{ ms}^{-1}$ . If string is 6m long. Find its fundamental frequency  
 A. 2 Hz  
 C. 4 Hz  
 B. 6 Hz  
 D. 8 Hz
- Q. 35 The restoring force of SHM is maximum when particle:  
 A. Displacement is maximum  
 C. Crossing mean position  
 B. Half way between them  
 D. At rest
- Q. 36 The essential properties of a medium for the propagation of mechanical waves are  
 A. Inertia and mass  
 C. Elasticity only  
 B. Inertia and elasticity  
 D. Inertia only
- Q. 37 The nature of sound waves in gases is  
 A. Transverse  
 C. Stationary  
 B. Longitudinal  
 D. Electromagnetic
- Q. 38 The frequency of a rod is 200 Hz. If the velocity of sound in air is  $340 \text{ ms}^{-1}$ , the wavelength of the sound produced is  
 A. 1.7 cm  
 C. 1.7 m  
 B. 6.8 cm  
 D. 6.8 m
- Q. 39 Standing waves are produced in 10m long stretched string. If string vibrates in 5 segments and wave velocity is 20m/s, what is the frequency?  
 A. 10Hz  
 C. 20Hz  
 B. 5Hz  
 D. 4Hz
- Q. 40 When two identical traveling waves are superimposed, velocity of resultant wave  
 A. Decreases  
 C. Remains same  
 B. Increases  
 D. Becomes zero
- Q. 41 The distance between 1<sup>st</sup> node and 4<sup>th</sup> antinode is:  
 A.  $\frac{7}{4}\lambda$   
 C.  $13\frac{\lambda}{4}$   
 B.  $5\frac{\lambda}{4}$   
 D.  $11\frac{\lambda}{4}$
- Q. 42 If  $V$  is the speed of sound at pressure  $P$  then speed of sound at  $2P$ , keeping temperature constant, will be  
 A. 1 : 2  
 C. 2 : 1  
 B. 1 : 1  
 D.  $\sqrt{2} : 1$
- Q. 43 Which of the following has maximum value of  $\gamma = \frac{C_p}{C_v}$   
 A. Monoatomic gas  
 C. Diatomic gas  
 B. Polyatomic gas  
 D. All have same value
- Q. 44 Velocity of sound in air  
 A. Decreases with increase in temperature  
 C. Decreases with decrease of temperature  
 B. Increase with decrease in temperature  
 D. Does not depend on temperature
- Q. 45 At what temperature, the velocity of sound will be double its value at 273 K?  
 A.  $2 \times 273 \text{ K}$   
 C.  $4 \times 273 \text{ K}$   
 B.  $8 \times 273 \text{ K}$   
 D.  $16 \times 273 \text{ K}$

Q. 46 For all gases

A.  $v_t = v_0 \sqrt{1 - \frac{t}{273}}$

C.  $v_t = v_0 \sqrt{1 + 273t}$

B.  $v_t = v_0 \sqrt{1 + \frac{t}{273}}$

D.  $v_t = v_0 \sqrt{1 + \frac{273}{t}}$

Q. 47 Sound travels faster in moist air at STP because

- A. Moist air is heavier than dry air
- B. The pressure of moist air is greater than that of dry air
- C. The value of  $\gamma$  of moist air is greater than that for dry air
- D. The density of moist air is less than that of dry air

Q. 48 Newton assumed that sound propagation in a gas takes place under

- A. Isothermal conditions
- B. Isobaric condition
- C. Adiabatic conditions
- D. Isochoric condition

Q. 49 If  $v_a$ ,  $v_h$  and  $v_m$  are the speeds of sound in air, hydrogen and a metal at the same temperature, then

- A.  $v_h > v_a > v_m$
- B.  $v_m > v_h > v_a$
- C.  $v_h > v_m > v_a$
- D.  $v_a > v_h > v_m$

Q. 50 A rope of length 5 m is stretched to a tension of 80 N. If its mass is 1 kg, at what speed would a 10 Hz transverse wave travel down the string?

- A. 2 m/s
- B. 5 m/s
- C. 20 m/s
- D. 50 m/s

Q. 51 At which temperature the speed of sound in hydrogen will be same as that of speed of sound in oxygen at 100°C

- A. -148°C
- B. -317.5°C
- C. -212.5°C
- D. -249.7°C

Q. 52 A stationary wave is set up in the air column of a closed pipe. At the closed end of the pipe:

- A. Always a node is formed
- B. Neither node nor antinode is formed
- C. Always an antinode is formed
- D. Sometimes a node and sometimes an antinode is formed

Q. 53 A string, clamped at its ends, vibrates in three segments. The string is 100cm long. The wavelength is:

- A. 33.3cm
- B. 66.7cm
- C. 150cm
- D. 300cm

Q. 54 In stationary wave

- A. Strain is maximum at nodes
- B. Strain is minimum at nodes
- C. Strain is maximum at antinodes
- D. Strain is minimum at antinodes

Q. 55 A property of the progressive wave that does not depend upon other characteristics mentioned below is

- A. Wavelength
- B. Frequency
- C. Amplitude
- D. Wave velocity

Q. 56 If the frequency of oscillation of a particle doing simple harmonic motion is  $n$ , the frequency of kinetic energy is?

- A.  $2n$
- B.  $n$
- C.  $n/2$
- D.  $2/n$

Q. 57 A spring (spring constant =  $k$ ) is cut into 4 equal parts and two parts are connected in parallel. What is the effective spring constant?

- A.  $4k$
- B.  $16k$
- C.  $8k$
- D.  $6k$

Q. 58 If a simple harmonic oscillator has got a displacement of 0.02m and acceleration equal to  $2\text{m/s}^2$  at any time, the angular frequency of the oscillator is equal to

- A. 10 rad/s
- B. 0.1 rad/s
- C. 100 rad/s
- D. 1 rad/s

Q. 59 Water waves refract at a boundary between deep water and shallow water. What is the effect on the frequency, wavelength and speed of the waves at the boundary?

	Frequency	Wavelength	Speed
A.	Changes	Changes	Stays the same
B.	Changes	Stays the same	Stays the same
C.	Stays the same	Changes	Changes
D.	Stays the same	Stays the same	Changes

Q. 60 When a particle performing uniform circular motion of radius 10 cm undergoes the SHM, what will be its amplitude?

- A. 10 cm
- B. 5 cm
- C. 2.5 cm
- D. 20 cm

## ANSWER KEY

1	C	11	A	21	C	31	D	41	A	51	D
2	B	12	B	22	C	32	A	42	B	52	A
3	A	13	C	23	C	33	B	43	A	53	B
4	D	14	D	24	C	34	A	44	C	54	A
5	D	15	D	25	C	35	A	45	C	55	C
6	D	16	D	26	B	36	B	46	B	56	A
7	D	17	A	27	A	37	B	47	D	57	C
8	D	18	B	28	C	38	C	48	A	58	A
9	C	19	C	29	A	39	B	49	B	59	C
10	A	20	C	30	D	40	C	50	C	60	A

# 5 UNIT

## THERMODYNAMICS

### SELF ASSESSMENT TEST

Q. 1 A system which can neither exchange matter nor energy with the surroundings is called

- A. Open system  
B. Isolated system  
C. Closed system  
D. Ideal system


Q. 2 A system absorb 10 kJ of heat at constant volume and its temperature rises from 270 C to 370 C. The value of  $\Delta U$  is

- A. 100 kJ  
B. 10 kJ  
C. 0 kJ  
D. 1 kJ

Q. 3 Metabolism is the name of a process in which energy transformation takes place within

- A. Heat engine  
B. Human body  
C. Atmosphere  
D. Laboratory

Q. 4 It is known that curves A, B, C are Isobaric, Isothermal, Adiabatic process then when one is correct



- A. A - Adiabatic, B - Isothermal, C - Isobaric  
B. A - Isothermal, B - Adiabatic, C - Isobaric  
C. A - Isobaric, B - Isothermal C - Adiabatic  
D. None of these

Q. 5 If for a gas  $\left(\frac{R}{C_v}\right) = 0.67$  this gas is made up of molecules which are

- A. Diatomic  
B. Mixture of diatomic and polyatomic  
C. Mono atomic  
D. Polyatomic

Q. 6 Specific heat capacity at constant volume of gases in an adiabatic process is

- A.  $\infty$   
B. Zero  
C. Finite but not zero  
D.  $0 < C_v < \infty$

Q. 7 The molar specific heat at constant volume  $C_v$  for a diatomic gas is

- A.  $3/2R$   
B.  $7/2R$   
C.  $5/2R$   
D.  $9/2R$

Q. 8 Which of following process provide us maximum work done by system?

- A. Isothermal  
B. Adiabatic  
C. Isochoric  
D. Isobaric

Q. 9 For an adiabatic compression  $W = 100J$ , what will be  $\Delta U$ ?

- A. - 100 J  
B. 0 J  
C. + 100 J  
D. None

Q. 10 For an adiabatic expansion:

- A. T decreases  
B. Mechanical energy goes out of the system  
C. Q = constant  
D. All of these

## Unit-5

## Thermodynamics

Q. 11 For adiabatic contraction:

- A. T increases  
B. Mechanical energy comes into system  
C.  $\Delta Q = \text{zero}$   
D. All of these

Q. 12 Unit of molar specific heat is same as that of

- A. Entropy  
B. General gas constant  
C. Internal energy  
D. Pressure

Q. 13 For mono-atomic gas  $C_v = \frac{3R}{2}$ , therefore  $\gamma$  for this gas is

- A.  $\frac{3}{2}$   
B.  $\frac{5}{3}$   
C.  $\frac{3}{5}$   
D.  $\frac{3}{4}$

Q. 14 If 1 mole of an idea gas is heated at constant volume, then

- A.  $\Delta U = C_v \Delta Q$   
B.  $\Delta U = C_v \Delta T$   
C.  $\Delta T = C_v \Delta U$   
D.  $\Delta U = C_p \Delta T$

Q. 15 Compressed air coming out of punctured football becomes cooler because of

- A. Isothermal expansion  
B. Energy dissipation  
C. Adiabatic expansion  
D. Adiabatic compression

Q. 16 The amount of heat required to raise the temperature of one mole of substance through 1 Kelvin at constant pressure is called

- A. Specific heat  
B. Molar specific heat capacity at constant pressure  
C. Molar heat capacity at constant pressure  
D. Heat capacity at constant pressure

Q. 17 Mathematically molar specific heat at constant pressure can be expressed as

- A.  $C_p = \frac{Q_p}{\Delta T}$   
B.  $C_p = \frac{T}{\Delta Q_p}$   
C.  $C_p = \frac{\Delta T}{Q_p}$   
D.  $C_p = \Delta Q_p \times T$

Q. 18 The motion possess by mono-atomic gas molecules is

- A. Translatory  
B. Rotatory  
C. Vibratory  
D. None of these


Q. 19 Sound passes through air under the following process

- A. Isobaric  
B. Isothermal  
C. Adiabatic  
D. Both B and C


Q. 20 The specific heat of a gas in an isothermal process is

- A. Infinity  
B. Negative  
C. Zero  
D. Remain constant


Q. 21 Which of the substances A, B or C has the highest specific heat? The temperature vs heat graph is



- A. A  
C. C
- Q. 22 The curve represents isothermal process is called  
A. Isotherm  
C. Adiabatic
- Q. 23 Which one is not adiabatic process  
A. Escape of air from burst tyre  
C. Slow expansion
- Q. 24 The work done on ideal gas during the cycle is



- B. B  
D. All have equal specific heats
- B. Both "A" and "C"  
D. Either "A" or "C"
- B. Cloud formation  
D. None
- B.  $2PV$   
D. 0
- Q. 25 Heat energy added to a system under isothermal conditions appears as  
A. Work done by the system  
C. Increase in internal energy
- Q. 26 The change in internal energy can be defined as  
A.  $\Delta U - W$   
C.  $P\Delta V$
- Q. 27 A gas expands  $0.25 \text{ m}^3$  at constant pressure  $10^3 \text{ N/m}^2$ . The work done is  
A. 2.5 ergs  
C. 300J
- Q. 28 The work done by an ideal monoatomic gas along path LMNO is



- B. Work done on the system  
D. Increase in temperature
- B.  $Q - W$   
D.  $Q + W$
- B. 250J  
D. 150J
- B.  $3PV$   
D.  $4PV$

- A. PV  
C.  $2PV$

Q. 29 During adiabatic expansion the change in internal energy of 2 moles of a gas is 100J, the work done during this expansion is  
A. Zero  
C. 200J

B. -100J  
D. 100J

Q. 30 Which of the following is necessarily zero when the system is an ideal gas that undergoes a change at constant temperature?  
A. Work  
C. Change in Internal Energy

B. Pressure  
D. None of these

Q. 31 In which process, the change in internal energy of the system is zero  
A. Isochoric process  
C. Adiabatic process

B. Isobaric process  
D. Isothermal process

Q. 32 In an equation  $PV^\gamma = \text{constant}$

A.  $\gamma = \frac{C_p}{C_v}$

B.  $\gamma = \frac{C_v}{C_p}$

C.  $\gamma = C_p - C_v$

D.  $\gamma = C_p C_v$


Q. 33 The temperature of the system decreases in the process of

- A. Free expansion  
C. Isothermal expansion
- B. Adiabatic expansion  
D. Isothermal compression

Q. 34 If both temperature and volume of an ideal gas are doubled, the pressure

- A. Remains constant  
C. Is also doubled
- B. Increases by a factor of 4  
D. Is diminished by a factor  $\frac{1}{4}$

Q. 35 In the diagrams (i) to (iv) of variation of volume with changing pressure is shown. A gas is taken along the path ABCD. The change in internal energy of the gas will be



- A. Positive in all cases (i) to (iv)  
B. Negative in cases (i), (ii) and (iii) but zero in (iv) case  
C. Positive in cases (i), (ii) and (iii) but zero in (iv) case  
D. Zero in all four cases

Q. 36 We can express the work in terms of

- A.  $P\Delta U$   
C.  $P\Delta V$
- B.  $P\Delta A$   
D. All are correct


Q. 37 Internal energy is a function of state because change in internal energy

- A. Does not depend on path  
C. Depends on path
- B. Corresponds to isothermal process  
D. Corresponds to adiabatic process

Q. 38 The internal energy of ideal gas is

- A. Totally K.E  
C. Totally P.E
- B. Partly K.E and partly P.E  
D. Neither K.E nor P.E

Q. 39 Which of the substances A, B or C has the highest specific heat? The temperature vs time graph is shown.



A. A  
 B. C  
 C. B  
 D. All have equal specific heat




Q. 40 The value of  $\gamma$  of triatomic gas (linear arrangement) molecules is

A.  $\frac{5}{3}$   
 B.  $\frac{8}{6}$   
 C.  $\frac{7}{5}$   
 D.  $\frac{9}{7}$

Q. 41 Two different samples have the same mass and temperature. Equal quantities of energy are absorbed as heat by each. Their final temperatures may be different because the samples have different:

A. Thermal conductivities  
 B. Coefficients of expansion  
 C. Densities  
 D. Heat capacities





Q. 42 Which of the following graphs between pressure and volume correctly shows isochoric change?



Q. 43 In a given process of an ideal gas,  $\Delta W = 0$  and  $\Delta Q = -ve$ . Then for the gas

A. The temperature will decrease  
 B. The pressure will remain constant  
 C. The volume will increase  
 D. The temperature will increase

Q. 44 For an isothermal process heat added to an ideal system versus work performed by that system is represented graphically



Q. 45 Let 100J of work is done to compress a gas adiabatically, then the change in internal energy is

A. 100J  
 B. +50J  
 C. -100J  
 D. -50J

Q. 46 A monoatomic gas ( $\gamma=5/3$ ) is suddenly compressed to 1/8 of its original volume adiabatically, then the pressure of the gas will change to

A. 24/5  
 B. 40/3  
 C. 8  
 D. 32 times its initial pressure

Q. 47 Heat neither enters nor leave a system then its temperature:

A. Must remain same  
 B. May fall  
 C. May rise  
 D. Both B & C

Q. 48 The amount of heat required to raise the temperature of one mole of substance through 1 Kelvin is called

A. Specific heat  
 B. Specific heat at constant volume  
 C. Molar specific heat  
 D. Heat capacity

Q. 49 If 1 mole of an ideal gas is heated at constant pressure, then:

A.  $Q_p = C_p \Delta T$   
 B.  $Q_v = C_v \Delta T$   
 C.  $Q_p = C_p \Delta T$   
 D.  $Q_v = C_p \Delta T$

Q. 50 The amount of heat energy required to raise the temperature of a body of mass 1 kg through 1 k is called:

A. Specific heat  
 B. Molar specific heat  
 C. Heat capacity  
 D. Heat of vaporization

Q. 51 The molar specific heat constant pressure of an ideal gas is  $7R/2$ . The ratio of specific heat at constant pressure to that at constant volume is?

A. 9/7  
 B. 7/5  
 C. 8/7  
 D. 5/7

Q. 52  $C_p - C_v$  and  $\frac{C_v}{C_p}$  are respectively equal to

A.  $\left(\frac{1}{R}, \gamma^{-1}\right)$   
 B.  $\left(\frac{1}{R}, \gamma\right)$   
 C.  $(R, g)$   
 D.  $(R, \gamma^1)$

Q. 53 Which one is correct relation?

A.  $C_p + C_v = \gamma$   
 B.  $\gamma = \frac{C_p}{C_v}$   
 C.  $C_p = 1 + \frac{R}{C_v}$   
 D.  $C_p = 1 - \frac{R}{C_v}$


Q. 54 Four students found set of  $C_p$  and  $C_v$  (in cal/deg mole) as given below. Which of the following set is correct?

A.  $C_v = 4, C_p = 2$   
 B.  $C_v = 3, C_p = 3$   
 C.  $C_v = 2, C_p = 1$   
 D.  $C_p = 5, C_v = 3$

Q. 55 Molar specific heat of gas at constant volume is  $\frac{5}{2}R$ . Find ratio of specific heat at constant volume to specific heat at constant pressure

A.  $\frac{2}{7}$   
 B.  $\frac{1}{7}$   
 C.  $\frac{5}{7}$   
 D. 1

Q. 56 A thermodynamic system is explained with the help of the graph ABCD. What is the heat rejected by the system during the cycle?



- A.  $pV$   
 B.  $2pV$   
 C.  $4pV$   
 D.  $1/2pV$
- Q. 57 If  $C_p = 34.4 \text{ J/mol K}$  and value of  $R$  is  $8.314$  then  
 A.  $C_v = 26.1 \text{ J/mol K}$   
 B.  $C_v = 20.1 \text{ J/mol K}$   
 C.  $C_v = 10.1 \text{ J/mol K}$   
 D.  $C_v = 120.1 \text{ J/mol K}$
- Q. 58 Which of the statement is true?  
 A.  $R = (\gamma - 1) C_v$   
 B.  $R = (\gamma + 1) C_v$   
 C.  $C_v = R(\gamma + 1)$   
 D.  $C_p = \frac{R}{(\gamma - 1)}$

- Q. 59 100 g of water is heated from  $30^\circ\text{C}$  to  $50^\circ\text{C}$ . Ignoring the slight expansion of the water, the change in its internal energy is (specific heat of water is  $4184 \text{ J kg}^{-1} \text{ K}^{-1}$ )  
 A. 4.2 kJ  
 B. 8.4 kJ  
 C. 84 kJ  
 D. 2.1 kJ
- Q. 60 When an ideal diatomic gas is heated at constant pressure, the fraction of the heat energy supplied which increases the internal energy of the gas is  
 A.  $2/5$   
 B.  $3/5$   
 C.  $3/7$   
 D.  $5/7$

# ANSWER KEY


2	B	11	D	21	C	31	D	41	D	51	B
3	B	12	B	22	A	32	A	42	C	52	D
4	A	13	B	23	C	33	B	43	A	53	B
5	C	14	B	24	A	34	A	44	B	54	D
6	B	15	C	25	A	35	D	45	A	55	C
7	C	16	B	26	B	36	C	46	D	56	B
8	D	17	A	27	B	37	A	47	D	57	A
9	C	18	A	28	A	38	A	48	C	58	A
10	D	19	C	29	D	39	B	49	C	59	B
		20	A	30	C	40	D	50	A	60	D

### SELF ASSESSMENT TEST


- Q.1 A parallel plate air capacitor has a capacity of  $2 \text{ pF}$ . If the separation between its plates is doubled and a mica sheet is introduced between its plates, its capacity becomes  $6 \text{ pF}$ . What is the dielectric constant of mica?  
 A. 6  
 B. 5  
 C. 4  
 D. 3
- Q.2 An electric charge at rest produces  
 A. Only a magnetic field  
 B. Only an electric field  
 C. Neither electric field nor magnetic field  
 D. Both electric and magnetic fields
- Q.3 Two similar charges each of one coulomb placed in the air one meter apart repel each other with a force  
 A.  $9 \times 10^9 \text{ N}$   
 B.  $9.2 \times 10^4 \text{ N}$   
 C.  $9 \times 10^8 \text{ N}$   
 D.  $9 \times 10^7 \text{ N}$
- Q.4 3 capacitors with capacitances  $5 \mu\text{F}$ ,  $6 \mu\text{F}$  and  $7 \mu\text{F}$  are arranged in series combination with a battery of  $5\text{V}$ , then which capacitor has more potential difference across a plates  
 A.  $5 \mu\text{F}$  capacitor  
 B.  $6 \mu\text{F}$  capacitor  
 C.  $7 \mu\text{F}$  capacitor  
 D. Same potential difference across all
- Q.5 A test charge is moved from lower potential point to a higher potential point. The potential energy of test charge will  
 A. Remain the same  
 B. Increase  
 C. Decrease  
 D. Become zero
- Q.6 The force per unit charge is known as  
 A. Electric flux  
 B. Electric intensity  
 C. Electric potential  
 D. All of above are same
- Q.7 Uncharged capacitor is connected in a series with resistor, switch and a battery. When switch is closed the potential drop across resistor  
 A. Remain same  
 B. Initially maximum then decreases to zero  
 C. Become zero then increase maximum  
 D. Remain zero
- Q.8 Two charges placed in air repel each other by a force of  $10^{-4} \text{ N}$ . When oil is introduced between the charges, the force becomes  $2.5 \times 10^{-5} \text{ N}$ . The dielectric constant of oil is  
 A. 3  
 B. 4  
 C. 2  
 D. 0.25
- Q.9 Two charges each equal to  $2 \mu\text{C}$  are  $0.5\text{m}$  apart. If both of them exist inside vacuum, then the force between them is  
 A. 2.44 N  
 B. 0.144 N  
 C. 1.89 N  
 D. 3.144 N
- Q.10 The electric lines of force are  
 A. Imaginary  
 B. Physically existing every where  
 C. Physically existing near the charges  
 D. Depends upon case
- Q.11 A capacitor of capacitance  $C$  has charge  $Q$  and stored energy is  $W$ . If the charge is increase to  $2Q$ . The stored energy will be  
 A.  $W/4$   
 B.  $W/2$   
 C.  $2W$   
 D.  $4W$

- Q.12 A hollow metal sphere of radius 5 cm is charged so that the potential on its surface is 10 V. The potential at the center of the sphere is
- A. 10 V  
B. Same as at point 5 cm away from the surface  
C. 0 V  
D. Same as at point 25 cm away from the surface

- Q.13 The variation of electric potential due to a point charge with distance is represented by the graph (where V along y-axis and r along x-axis)




- Q.14 An electron of charge e is introduced between two metal plates a distance d apart. A potential difference V is applied to the plates as shown in the diagram.



- Which expression gives the electric force F on the electron?
- A.  $\frac{eV}{d}$   
B.  $eVd$   
C.  $\frac{V}{ed}$   
D.  $\frac{dV}{e}$

- Q.15 The energy stored per unit volume in an electric field of strength E volt/meter in a medium of dielectric constant K (in Joule/metre<sup>3</sup>) is:
- A.  $\frac{1}{2} \epsilon_0 E^2$   
B.  $\frac{1}{2} K \epsilon_0 E^2$   
C.  $\frac{1}{2} \frac{\epsilon_0 E^2}{K}$   
D.  $\frac{1}{2} K^2 \epsilon_0^2 E$

- Q.16 Which diagram represents the electric fields of a negative point charge, shown by.




- Q.17 A charge of 2C experiences a force 2000N in a uniform electric field. In this field the potential difference between two points separated by a distance 1cm is
- A. 2V  
B. 5V  
C. 10 V  
D. 20V


- Q.18 A 5 μF capacitor has a potential difference across its plates is 200 volts. The charge on the capacitor is
- A.  $2.5 \times 10^{-8}$  C  
B.  $10^{-3}$  C  
C.  $10^3$  C  
D.  $4 \times 10^3$  C

- Q.19 If the distance between the plates of parallel plates condenser is increased, its potential will
- A. Remain same  
B. Increase  
C. Decrease  
D. Decreases exponentially


- Q.20 The number of electrons in one coulomb charge is equal to
- A.  $6.2 \times 10^{18}$   
B.  $1.6 \times 10^{19}$   
C.  $6.2 \times 10^{21}$   
D.  $1.6 \times 10^{27}$
- Q.21 The amount of work done in joule in carrying a charge +q along the closed path PQRSP between the oppositely charged metal plates is (where E is electric field between the plates)



- A. Zero  
B. q  
C.  $qE(PQ + QR + SR + SP)$   
D.  $\frac{q}{\epsilon_0}$
- Q.22 The graph shows the growth of charge with potential difference between plates. The area under the graph shows




- A. Capacitance  
B. Separation of plates  
C. Energy stored  
D. Electric intensity
- Q.23 Value of  $\epsilon_r$  for various dielectrics is always
- A. Less than unity  
B. Equal to unity  
C. Larger than unity  
D. No hard and fast rule
- Q.24 In the electric field of a point charge q, a certain charge is carried from point A to B, C, D and E. Then the work done



- A. Is least along the path AB  
B. Is zero along all the paths AB, AC, AD and AE  
C. Is least along AE  
D. Is least along the path AD
- Q.25 A charge particle is free to move in an electric field. It will travel
- A. Along a line of force, if it has some initial velocity in the direction of an acute angle with the line of force  
B. Always along a line of force  
C. Along a line of force, if its initial velocity is zero  
D. None of these

Unit-6

Q.26 An uncharged sphere of metal is placed in between two charged plates as shown. The lines of force look like



- A. C  
B. A  
C. D  
D. B

Q.27 The intensity of electric field required to balance a proton of mass  $1.7 \times 10^{-27}$  kg and charge  $1.6 \times 10^{-19}$  C is nearly

- A.  $1 \times 10^{-7}$  V/m  
B.  $1 \times 10^{-5}$  V/m  
C.  $1 \times 10^7$  V/m  
D.  $1 \times 10^5$  V/m

Q.28 Two plates are 2 cm apart, a potential difference of 10 volt is applied between them, the electric field between the plates is

- A. 20 N/C  
B. 500 N/C  
C. 5 N/C  
D. 250 N/C

Q.29 A particle A has charge  $+q$  and a particle B has charge  $+4q$  with each of them having the same mass  $m$ . When allowed to fall from rest through the same electric potential difference, the ratio of their speed  $v_A/v_B$  will become

- A. 1:2  
B. 1:4  
C. 2:1  
D. 4:1

Q.30 What would happen to the electrostatic force between a pair of charged particles if both charges were doubled and the distance between them were also doubled?

- A. It would decrease by a factor of 4  
B. It would decrease by a factor of 2  
C. It would remain unchanged  
D. It would increase by a factor of 2.

Q.31 Two charges  $1 \mu\text{C}$  and  $5 \mu\text{C}$  separated by 20 cm, the ratio of electric forces acting on them will be:

- A. 1:2  
B. 1:1  
C. 1:5  
D. 5:1

Q.32 In bringing an electron towards the second electron the electrostatic potential of the system

- A. Increases  
B. Remains the same  
C. Decreases  
D. Becomes zero

Q.33 A capacitor is charged by using a battery which is then disconnected. A dielectric slab is then slipped between the plates, which results in

- A. Reduction of charge on the plates and increase of potential difference across the plates  
B. Increase in the potential difference across the plate, reduction in stored energy, but no change in the charge on the plates  
C. Decrease in the potential difference across the plates, reduction in the stored energy, but no change in the charge on the plates  
D. None of the above

Q.34 Unit of energy density of electric field is:

- A.  $\text{JC}^{-1}$   
B.  $\text{Jm}^{-3}$   
C.  $\text{JV}^{-1}$   
D.  $\text{JF}^{-3}$

Q.35 Coulomb's law applicable for

- A. Any two charges  
B. Point charges  
C. Both  
D. None

Q.36 Three charges  $2q, -q, -q$  are located at the vertices of an equilateral triangle. At the center of the triangle.

- A. The field is zero but potential is non-zero  
B. The field is non-zero but potential is zero  
C. Both field and potential are zero  
D. Both field and potential are non-zero

Q.37 An external agency carries  $-5 \text{ C}$  of charge from infinity to a point in an electrostatic field and performs 100 joule of work. The potential at the given point is

- A. 10 V  
B. 20 V  
C.  $-10 \text{ V}$   
D.  $-20 \text{ V}$

Q.38 A tin nucleus has charge  $+50e$ . If the proton is at a distance  $10^{-12}$  m from the nucleus, then the potential  $V$  at this position is [charge on the proton =  $1.6 \times 10^{-19}$  C]

- A.  $14.4 \times 10^4$  volt  
B.  $7.2 \times 10^8$  volt  
C.  $7.2 \times 10^4$  volt  
D.  $14.4 \times 10^8$  volt

Q.39 Four charges  $2\text{C}, -3\text{C}, -4\text{C}$  and  $5\text{C}$  respectively are placed at all the corners of a square. Which of the following statements is true for the point of intersection of the diagonals?

- A. Electric field is zero but electric potential is non-zero  
B. Electric field non-zero but electric potential is zero  
C. Both electric field and electric potential are zero  
D. Neither electric field nor electric potential is zero

Q.40 Two charges are placed at a certain distance. If the magnitude of each charge is doubled the force will become

- A.  $\frac{1}{4}$  th of its original value  
B.  $\frac{1}{8}$  th of its original value  
C. 4 times of original value  
D. 8 times of its original value

Q.41 Force between the plates of a charged parallel plate capacitor is attractive and its magnitude is

- A.  $\frac{Q^2}{A\epsilon_0}$   
B.  $\frac{Q^2}{2A\epsilon_0}$   
C.  $Q^2 A\epsilon_0$   
D.  $QA\epsilon_0$

Q.42 The electric intensity at infinite distance from the point charge is

- A. Zero  
B. Infinite  
C. 1-volt  $\text{m}^{-1}$   
D. Negative

Q.43 Value of  $\epsilon_r$  for air is;

- A. 1.6  
B. 1.96  
C. 1.986  
D. 1.0006

Q.44 What will be the electric potential energy of a 7 nC charge that is 2 cm from a 20 nC charge?


- A.  $6.3 \times 10^{-1}$  J  
B.  $6.3 \times 10^{-5}$  J  
C.  $6.3 \times 10^{-5}$  V  
D.  $1.3 \times 10^{-5}$  J

Q.45 The potential at a point, due to a positive charge of  $100 \mu\text{C}$  at a distance of 9 m, is


- A.  $10^4$  V  
B.  $10^5$  V  
C.  $10^6$  V  
D.  $10^7$  V



- Q.46 The quantity  $\frac{\Delta V}{\Delta r}$  is known as:  
 A. Potential Difference  
 C. Potential Gradient  
 B. Potential Energy  
 D. All of these
- Q.47 A unit positive charge +  $q_0$  placed anywhere in the vicinity of a positive point charge, experiences a repulsive force directed.  
 A. Radially inward  
 C. Radially zero  
 B. Radially out ward  
 D. None of these
- Q.48 If  $E$  be the electric intensity of an electrostatic field, then the electrostatic energy density is proportional to  
 A.  $E$   
 C.  $1/E^2$   
 B.  $E^2$   
 D.  $E^3$
- Q.49 The coulombs force between two point charges is  $F$ . If magnitude of each charge is doubled and distance between charges is halved, the forces between them becomes  
 A.  $F$   
 C.  $4F$   
 B.  $8F$   
 D.  $16F$
- Q.50 Relative permittivity  $\epsilon_r$  is defined by the following relation.  
 A.  $\frac{C_{med}}{C_{vac}}$   
 C.  $C_{med} \times C_{vac}$   
 B.  $\frac{C_{vac}}{C_{med}}$   
 D.  $C_{med} + C_{vac}$
- Q.51 Two point charges are separated by a distance of 4 m. The force between them is 4 N. What is the force between the charges, when the distance between them is 1 m?  
 A. 16 N  
 C. 1 N  
 B. 64 N  
 D. 32 N
- Q.52 An electric field exists in the space between two charged metal plates.



Which graph shows the variation of electric field strength  $E$  with distance  $d$  from X along the line XY?




- Q.53 Gaussian surface is  
 A. An imaginary surface  
 C. An open surface  
 B. A curved surface  
 D. A plane surface
- Q.54 12 J of work is to be done against an existing electric field to take a charge of 0.01 C from A to B. Find the potential difference between B and A.  
 A. 120 V  
 C. 1200 V  
 B. 1.2 V  
 D. 12 V
- Q.55 When one electron is taken towards the other electron, then the electric potential energy of the system  
 A. Decreases  
 C. Increases  
 B. Remains unchanged  
 D. Becomes zero

- Q.56 A parallel plate capacitor is charged and the charging battery is disconnected. Then a dielectric slab is introduced between the plates. The quantity that remains unchanged is \_\_\_\_\_  
 A. Potential  
 C. Energy  
 B. Capacity  
 D. Charge
- Q.57 Electric potential is \_\_\_\_\_  
 A. A scalar quantity  
 C. A dimensionless quantity  
 B. A vector quantity  
 D. Always a positive quantity
- Q.58 The work done in placing a charge of 8  $\mu\text{C}$  on a condenser of capacity 100 nF is \_\_\_\_\_  
 A.  $16 \times 10^{-5}$  J  
 C.  $3.2 \times 10^{-4}$  J  
 B.  $32 \times 10^{-6}$  J  
 D.  $16 \times 10^{-4}$  J
- Q.59 The metallic spheres A and B of radii 2m and 4m respectively carry the same charge  $4 \times 10^8$  C. If the spheres are connected by a copper wire  
 A. The charge will flow from A to B  
 C. No flow of charge will occur  
 B. The charge will flow from B to A  
 D. Both A and B are possible
- Q.60 Let  $Q$  denote charge,  $V$  denote potential difference, and  $U$  denote stored energy. Of these quantities, capacitors in parallel must have the same:  
 A.  $Q$  only  
 C.  $U$  only  
 B.  $V$  only  
 D.  $Q$  and  $U$  only

## ANSWER KEY

1	A	11	D	21	A	31	B	41	A	51	B
2	B	12	A	22	C	32	A	42	A	52	C
3	A	13	D	23	C	33	C	43	D	53	A
4	A	14	A	24	B	34	B	44	B	54	C
5	B	15	B	25	C	35	B	45	B	55	C
6	B	16	B	26	A	36	B	46	C	56	D
7	B	17	C	27	A	37	D	47	B	57	A
8	B	18	B	28	B	38	C	48	B	58	C
9	B	19	B	29	A	39	B	49	D	59	A
10	A	20	A	30	C	40	C	50	A	60	B

- Q.1 When a piece of aluminium wire of finite length is drawn through a series of dies to reduce its diameter to half its original value, its resistance will become  
 A. Two times  
 B. Four times  
 C. Eight times  
 D. Sixteen times
- Q.2 An electric wire is connected across a cell of e.m.f.  $E$ . The current  $I$  is measured by an ammeter of resistance  $R$ . According to ohm's law  
 A.  $E=I^2R$   
 B.  $E=IR$   
 C.  $E=R/I$   
 D.  $E=I/R$
- Q.3 Resistance of tungsten wire at  $150^\circ\text{C}$  is  $133\Omega$ . Its resistance temperature coefficient is  $0.0045/^\circ\text{C}$ . The resistance of this wire at  $500^\circ\text{C}$  will be  
 A.  $180\Omega$   
 B.  $225\Omega$   
 C.  $258\Omega$   
 D.  $317\Omega$
- Q.4 A bulb has a power of  $200\text{W}$ . What is the energy dissipated by it in 5 minutes?  
 A.  $60\text{J}$   
 B.  $1000\text{J}$   
 C.  $60\text{kJ}$   
 D.  $1\text{kJ}$
- Q.5 A wire of uniform area of cross section is cut into two parts of equal lengths. The resistivity of any part  
 A. Remain same  
 B. Is doubled  
 C. Is halved  
 D. One fourth
- Q.6 The resistance of a coil is  $4.2\text{ ohm}$  at  $100^\circ\text{C}$  and the temperature coefficient of resistance of its material is  $0.004/^\circ\text{C}$ . Its resistance at  $0^\circ\text{C}$   
 A.  $6.5\text{ ohm}$   
 B.  $5\text{ ohm}$   
 C.  $3\text{ ohm}$   
 D.  $4\text{ ohm}$
- Q.7 Two cells of e.m.f  $E_1$  and  $E_2$  and of negligible internal resistance are connected with two variable resistors as shown in the diagram.



When the galvanometer deflection is zero, the values of both resistances are  $P$  and

Q. what is the value of the ratio  $\frac{E_2}{E_1}$  ?

- A.  $\frac{P}{Q}$   
 B.  $\frac{Q}{(P+Q)}$   
 C.  $\frac{P}{(P+Q)}$   
 D.  $\frac{(P+Q)}{P}$

- Q.8 10,000 alpha particles per minute, are passing through a straight tube of radius  $r$ . The resulting electric current in approximately  
 A.  $0.5 \times 10^{-16}\text{ amp}$   
 B.  $0.5 \times 10^{12}\text{ amp}$   
 C.  $2 \times 10^{12}\text{ amp}$   
 D.  $2 \times 10^{-12}\text{ amp}$
- Q.9 An electric current source is actually source of  
 A. Current  
 B. Charge  
 C. Energy  
 D. Power
- Q.10 What can be used as the unit of energy  
 A. watt  $\times$  second  
 B. volt  $\times$  meter  
 C. volt per coulomb  
 D. newton per meter
- Q.11 A nichrome wire  $50\text{ cm}$  long and one square millimetre cross-section carries a current of  $4\text{A}$  when connected to a  $2\text{V}$  battery. The resistivity of nichrome wire in ohm metre is  
 A.  $1 \times 10^{-6}$   
 B.  $4 \times 10^{-7}$   
 C.  $3 \times 10^{-7}$   
 D.  $2 \times 10^{-7}$
- Q.12 Calculate the amount of charge flowing in 2 minutes in a wire of resistance  $10\text{ Ohm}$  when a potential difference of  $20\text{ V}$  is applied between its ends  
 A.  $4\text{ C}$   
 B.  $20\text{ C}$   
 C.  $240\text{ C}$   
 D.  $120\text{ C}$
- Q.13 An electric iron is marked  $20\text{ volts } 500\text{W}$ . The units consumed by it in using it for 24 hours will be  
 A. 12  
 B. 24  
 C. 5  
 D. 1100
- Q.14 The electric resistance of a certain wire of iron is  $R$ . If its length and radius are both doubled, then  
 A. The resistance will be halved and the specific resistance will remain unchanged  
 B. The resistance will be doubled and the specific resistance will be halved  
 C. The resistance will be halved and the specific resistance will be doubled  
 D. The resistance and the specific resistance, will both remain unchanged
- Q.15 The graphical representation of Ohm's law is \_\_\_\_\_  
 A. Hyperbola  
 B. Ellipse  
 C. Parabola  
 D. Straight line
- Q.16 SI unit of resistivity is \_\_\_\_\_  
 A.  $\Omega\text{-m}^2$   
 B.  $(\Omega\text{-m})^{-1}$   
 C.  $\Omega\text{-m}$   
 D.  $(\Omega\text{-m})^{-1}$
- Q.17 A cell is connected to a resistor. What is the e.m.f of the cell equal to?  
 A. The potential difference across the resistor for each unit of current  
 B. The power produced in the circuit for each unit of charge that passes  
 C. The work done in the circuit for each unit of charge that passes  
 D. The work done in the circuit for each unit of current
- Q.18 Electric current is generated by;  
 A. Diode  
 B. Transistor  
 C. Generator  
 D. Motor
- Q.19 A flow of  $10^7$  electron per second in a conductor constitutes a current of  
 A.  $1.6 \times 10^{-26}\text{ A}$   
 B.  $1.6 \times 10^{12}\text{ A}$   
 C.  $1.6 \times 10^{-12}\text{ A}$   
 D.  $10^7\text{ A}$

- Q.20 A 25 W, 220 V bulb and a 100 W, 220 V bulb are joined in series and connected to mains which bulb will glow brighter  
 A. 25 W  
 B. 100 watt bulb  
 C. First 20W and then 100 W bulb  
 D. Neither bulb will glow
- Q.21 A typical value of drift velocity is  
 A.  $1 \text{ mm s}^{-1}$   
 B.  $1 \text{ ms}^{-1}$   
 C.  $10 \text{ ms}^{-1}$   
 D.  $1000 \text{ km s}^{-1}$
- Q.22 The specific resistance of wire is  $50 \times 10^{-8} \Omega \text{ m}$ . The resistance of a cube of length 50 cm will be  
 A.  $10^{-6} \Omega$   
 B.  $2.5 \times 10^{-5} \Omega$   
 C.  $10^{-8} \Omega$   
 D.  $5 \times 10^{-4} \Omega$
- Q.23 5A of current is passed through a metallic conductor. The charge flowing in one minute in coulomb, will be  
 A. 5  
 B. 12  
 C.  $\frac{1}{12}$   
 D. 300
- Q.24 The length of the wire is doubled. Its conductance will be  
 A. Uncharged  
 B. Halved  
 C. Halved  
 D.  $\frac{1}{4}$  of the original value
- Q.25 1.6 mA current is flowing in conducting wire then the number of electrons flowing per second  
 A.  $10^{11}$   
 B.  $10^{16}$   
 C.  $10^{19}$   
 D.  $10^{15}$
- Q.26 A primary cell has an emf of 1.5V, when short circuited it gives a current of 3A. The internal resistance of the cell is:  
 A.  $4.5 \Omega$   
 B.  $2 \Omega$   
 C.  $0.5 \Omega$   
 D.  $\frac{1}{4.5} \Omega$
- Q.27 For which of the following the resistance decreases on increasing the temperature  
 A. Copper  
 B. Tungsten  
 C. Germanium  
 D. Aluminium
- Q.28 For a metallic wire, the ratio  $V/i$  ( $V$  = the applied potential difference,  $i$  = current flowing) is  
 A. Independent of temperature  
 B. Increases as the temperature rises  
 C. Decreases as the temperature rises  
 D. Increases or decreases as temperature rises, depending upon the metal
- Q.29 The resistances of a wire at temperatures  $t^\circ\text{C}$  and  $0^\circ\text{C}$  are related by  
 A.  $R_t = R_0(1 + \alpha t)$   
 B.  $R_t = R_0(1 - \alpha t)$   
 C.  $R_t = R_0^2(1 + \alpha t)$   
 D.  $R_t = R_0^2(1 - \alpha t)$
- Q.30 Total number of electrons present in 4 ampers current flowing for 1 sec is:  
 A.  $2.5 \times 10^{19}$   
 B.  $1.25 \times 10^{13}$   
 C.  $2.5 \times 10^{20}$   
 D.  $6 \times 10^8$

- Q.31 If a source of emf is traversed from positive to negative the potential change will be  
 A. Positive  
 B. Zero  
 C. Negative  
 D. Constant
- Q.32 Resistivity at a given temperature depends upon:  
 A. Area of cross-section  
 B. Length  
 C. Nature of material of conductor  
 D. Both length and area
- Q.33 A cell of emf  $E$  Volt and internal resistance  $r$  ohm is being charged with a current of  $i$  amp. Then the terminal potential difference is  
 A.  $E$   
 B.  $E + ir$   
 C.  $E - ir$   
 D.  $E - iR$
- Q.34 The temperature of a metal wire rises when an electric current passes through it because  
 A. Collision of metal atoms with each other releases heat energy  
 B. Collision of conduction electrons with each other releases heat energy  
 C. Collision of conduction electrons with the atoms of metal give them energy which appears as heat  
 D. all of these
- Q.35 If a bulb has a 20W power. If it is working at 60% efficiency, then its output power is  
 A. 12W  
 B. 15W  
 C. 20W  
 D. 18W
- Q.36 The substances which have a large number of free electrons and offer a low resistance are called  
 A. Insulators  
 B. semi-conductors  
 C. Inductors  
 D. Conductors
- Q.37 A light bulb draws 300 mA when the voltage across it is 240 V. The resistance of the light bulb is  
 A. 400 Q  
 B. 800 Q  
 C. 600 Q  
 D. 1000 Q
- Q.38 Two copper wires X and Y have the same volume. Wire Y is four times as long as wire X.




What is the ratio  $\frac{\text{resistance of wire Y}}{\text{resistance of wire X}}$ ?

- A. 4  
 B. 16  
 C. 8  
 D. 64
- Q.39 Which of the following materials has a negative temperature co-efficient of resistance?  
 A. Copper  
 B. Carbon  
 C. Aluminum  
 D. Brass
- Q.40 The terminal potential difference of a cell when short-circuited is ( $E$  = E.M.F. of the cell)  
 A.  $E$   
 B.  $E/2$   
 C. Zero  
 D.  $E/3$

- Q.41 By a cell a current of 0.9 A flows through 2 ohm resistor and 0.3 A through 7 ohm resistor. The internal resistance of the cell is  
 A. 0.5Ω B. 1.0Ω  
 C. 1.2Ω D. 2.0Ω


- Q.42 The emf of the cell in the following circuit is 9.0 V. The reading on the high resistance voltmeter 7.5 V?



- What is the current I?  
 A. 0.1 A B. 0.5 A  
 C. 0.6 A D. 2.0 A
- Q.43 Two bulbs having the ratings 40 W, 220 V and 20 W, 110 V. The ratio of their resistance is  
 A. 1 : 2 B. 1 : 1  
 C. 2 : 1 D. 1 : 4
- Q.44 A total charge of 100 C flows through a 12 W light bulb in a time of 50s. What is the potential difference across the bulb during this time?  
 A. 0.12 V B. 6.0 V  
 C. 2.0 V D. 24 V
- Q.45 Two bulbs of 500 W and 200 W rated at 250 V will have resistance ratio as  
 A. 4 : 25 B. 2 : 5  
 C. 25 : 4 D. 5 : 2
- Q.46 The resistance of a metallic wire becomes 8 times when;  
 A. Length is doubled B. Length is tripled  
 C. Length is doubled and radius is halved D. Length is halved and radius is doubled
- Q.47 The internal resistance of a cell is the resistance of  
 A. Electrodes of the cell B. Vessel of the cell  
 C. Electrode used in the cell D. Material used in the cell
- Q.48 An electron is circulating in a circular path with a frequency of 50 Hz. What is the associated current?  
 A.  $0.8 \times 10^{-17}$  A B.  $8 \times 10^{-17}$  A  
 C.  $0.4 \times 10^{-17}$  A D.  $80 \times 10^{-17}$  A
- Q.49 How many electrons per second constitute a current of one micro ampere?  
 A. One electron B.  $10^{-6}$  electrons  
 C.  $10^6$  electrons D.  $6.25 \times 10^{12}$  electrons
- Q.50 A steady current is flowing in a conductor of non-uniform cross-section. The charge passing through any cross-section per unit time is  
 A. Directly proportional to the area of cross-section  
 B. Proportional to square of the area of cross-section  
 C. Inversely proportional to the area of cross-section  
 D. Independent of the area of cross-section
- Q.51 In the case of gases, the charge carries are  
 A. Positive and negative ions B. Negative ions and electrons  
 C. Electrons and holes D. Positive ions and electron

- Q.52 What is meant by 5 A?  
 A. A charge of 5 C flows through a point in 1 second  
 B. 5V electricity is causing 1 C of charge to flow  
 C. 5 V electricity flows across 1Ω of resistance  
 D. A charge of 5 C flows through a point in 5 seconds.
- Q.53 Slope of the graph between "V" On X-axis and "I" on the Y-axis is  
 A. Resistance B. Conductance  
 C. Emf D. Capacitance
- Q.54 When we double the voltage in a simple electric circuits we double the  
 A. Current B. Power  
 C. Resistance D. Both 'A' and 'B'
- Q.55 The 40 watt, 100 watt and 150 watt bulbs are connected in series across 220-volt supply. Which bulb will be brightest?  
 A. 150 watt B. 100 watt  
 C. 40 watt D. All have same brightness
- Q.56 At what temperature will the resistance of a copper wire become three times its value at 0°C (Temperature coefficient of resistance for copper =  $4 \times 10^{-3}$  per °C)  
 A. 400°C B. 450°C  
 C. 500°C D. 550°C
- Q.57 Two electric bulbs, one of 200volt 40 watt and the other 200volt 100 watt are connected in a house wiring circuit  
 A. They have equal currents through them  
 B. The resistance of the filaments in both the bulbs is same  
 C. The resistance of the filament in 40watt bulb is more than the resistance in 100watt bulb  
 D. The resistance of the filament in 100watt bulb is more than the resistance in 40watt bulb
- Q.58 Two electric bulbs have 40W and 60W rating at 220V, then the ratio of their resistances will be  
 A. 3 : 2 B. 2 : 3  
 C. 3 : 4 D. 4 : 3
- Q.59 If there are two bulbs connected in series and one blows out, what happens to the other bulb?  
 A) The other bulb continues to glow with the same brightness  
 B) The other bulb stops glowing  
 C) The other bulb glows with increased brightness  
 D) The other bulb also burns out

Q.60 A student carries out an experiment and plots the V-I graph of three samples of nichrome wire with resistances  $R_1$ ,  $R_2$  and  $R_3$  respectively. Which of the following is true?



- A.  $R_1 = R_2 = R_3$   
 B.  $R_1 > R_2 > R_3$   
 C.  $R_2 > R_3 > R_1$   
 D.  $R_3 > R_2 > R_1$

## ANSWER KEY


1	D	11	A	21	A	31	C	41	A	51	D
2	B	12	C	22	A	32	C	42	B	52	A
3	C	13	A	23	D	33	B	43	C	53	B
4	C	14	A	24	B	34	C	44	B	54	A
5	A	15	D	25	B	35	A	45	B	55	C
6	C	16	C	26	C	36	D	46	C	56	C
7	C	17	C	27	C	37	B	47	D	57	C
8	A	18	C	28	B	38	B	48	A	58	A
9	C	19	C	29	A	39	B	49	D	59	B
10	A	20	A	30	A	40	C	50	A	60	C

# 8 UNIT


## ELECTROMAGNETISM AND ELECTROMAGNETIC INDUCTION

### SELF ASSESSMENT TEST

- Q.1 When the current in the current carrying conductor increases, what happens to the force in the conductor which is at right angles to the magnetic field?  
 A. Increases  
 B. Decreases  
 C. Remains the same  
 D. Becomes zero
- Q.2 The total flux in the core of an electrical machine is 20 mWb and its flux density is 1 T. The cross-sectional area of the core is:  
 A.  $0.05 \text{ m}^2$   
 B.  $0.02 \text{ m}^2$   
 C.  $20 \text{ m}^2$   
 D.  $50 \text{ m}^2$
- Q.3 A positive charge enters in a magnetic field and travels parallel to the field. Its speed experiences is  
 A. Increases  
 B. Decreases  
 C. Remain same  
 D. None of these
- Q.4 In the given figure, force on rod A and rod B are in direction respectively:






- A. Rightward direction, leftward direction  
 B. Rightward direction, Rightward direction  
 C. Leftward direction, Leftward direction  
 D. Leftward direction, Rightward direction
- Q.5 In velocity selector arrangement electric and magnetic field will be  
 A. Parallel each other  
 B. Anti-parallel each other  
 C. Perpendicular to each other  
 D. Inclined at  $45^\circ$  to each other
- Q.6 A proton is projected in a region containing both electric and magnetic field pointing in opposite direction to the motion of proton. The proton may  
 A. Move in same direction with increasing speed  
 B. Deflect downward with decreasing speed  
 C. Deflect upward with same speed  
 D. Move in same direction with decreasing speed



- Q.7 Magnetic flux would be zero when  
 A.  $\vec{B}$  is parallel to  $\vec{A}$   
 B.  $\vec{B}$  is along to  $\vec{A}$   
 C.  $\vec{B}$  is perpendicular to  $\vec{A}$   
 D. None of these
- Q.8 If an electron projected in a magnetic field with a velocity  $v$ , it will experience a force given by  
 A.  $\vec{F} = -e \vec{v} \times \vec{B}$   
 B.  $\vec{F} = +e \vec{v} \times \vec{B}$   
 C.  $\vec{F} = -e \vec{v} \cdot \vec{B}$   
 D.  $\vec{F} = +e \vec{v} \cdot \vec{B}$

- Q.9 A proton charge (+e coulomb) enters in a magnetic field of strength  $B$  (Tesla) making an angle  $30^\circ$  with the direction of magnetic field with speed  $v$ . The magnetic force on the proton is
- A.  $evB$   
 B. Zero  
 C.  $\infty$   
 D.  $\frac{evB}{2}$

- Q.10 The magnetic field intensity  $e$  at a distance ' $r$ ' from a long straight conductor carrying steady current  $I$  varies with ' $r$ ' as shown in the figure:




- Q.11 Magnetic field is produced by the flow of current in a straight wire. This phenomenon was discovered by
- A. Faraday  
 B. Coulomb  
 C. Maxwell  
 D. Oersted
- Q.12 Protons are shot perpendicular to a magnetic field
- A. The magnetic field will have no influence on the motion of the protons  
 B. The protons will continue to move in the opposite direction but will gain momentum  
 C. The protons will continue to move in the same direction but will gain momentum  
 D. They will bend in an arc of a circle
- Q.13 The magnetic induction at a point P which is distant 4 cm from a long current carrying wire is  $10^{-3}$  tesla. The field of induction at a distance 12 cm from the same current would be;
- A.  $3.33 \times 10^{-4}$  T  
 B.  $3 \times 10^{-3}$  T  
 C.  $1.11 \times 10^{-4}$  T  
 D.  $9 \times 10^{-3}$  T
- Q.14 A proton and an alpha-particle enter a uniform magnetic field with the same velocity. The period of rotation of the alpha-particle will be
- A. Four times that of the proton  
 B. Three times that of the proton  
 C. Two times that of the proton  
 D. Same as that of the protons
- Q.15 An electron moves at  $2 \times 10^2$  m/sec perpendicular to magnetic field of 2T. What is the magnitude of magnetic force?
- A.  $1 \times 10^{-6}$  N  
 B.  $6.4 \times 10^{-17}$  N  
 C.  $3.6 \times 10^{-24}$  N  
 D.  $4 \times 10^6$  N
- Q.16 When a charged particle moves through a magnetic field, the effect of the field changes the particles
- A. Speed  
 B. Mass  
 C. Energy  
 D. Direction

- Q.17 An electron is injected into a uniform magnetic field with components of velocity parallel to and normal to the field direction. The path of the electron is a
- A. Helix  
 B. Parabola  
 C. Circle  
 D. Straight line
- Q.18 If the current flowing through the conductor is made two times. Magnetic field strength due to it will increase;
- A. Two times  
 B. Remain same  
 C. Three times  
 D. Four times
- Q.19 Which of the following does not affect the motion of a moving electron?
- A. Electric field applied in the direction of motion  
 B. Electric field applied perpendicular to the direction of motion  
 C. Magnetic field applied in the direction of motion  
 D. Magnetic field applied perpendicular to the direction of motion
- Q.20 An electron is moving north in a region where the magnetic field is south. The magnetic force exerted on the electron is:
- A. Zero  
 B. Down  
 C. Up  
 D. East west
- Q.21 A charged particle travelling in a uniform field could have a circular trajectory if the field is
- A. Gravitational  
 B. Electrical  
 C. Magnetic  
 D. Gravitational or electrical
- Q.22 A uniform magnetic field will cause a charge to move in a circle if the charge is moving
- A. Perpendicular to the field lines and the field strength is constant  
 B. Perpendicular to the field and the field is increasing  
 C. Parallel to the field and the field is constant  
 D. Parallel to the field and the field is increasing
- Q.23 The magnetic flux through a wire loop in a magnetic field does not depend on
- A. The area of the loop  
 B. The shape of the loop  
 C. The magnitude of the field  
 D. The angle between the plane of the loop and the direction
- Q.24 A proton (or charged particle) moving with velocity  $v$  is acted upon by electric field  $E$  and magnetic field  $B$ . The proton will move undeflected if
- A.  $E$  is perpendicular to  $B$   
 B.  $E$ ,  $B$  and  $v$  are mutually perpendicular and  $v = E/B$   
 C.  $E$  is parallel to  $v$  and perpendicular to  $B$   
 D.  $E$  and  $B$  both are parallel to  $v$
- Q.25 The direction of magnetic lines of force produced by passing a direct current in a conductor is
- A. Perpendicular to the conductor and coming outward  
 B. Parallel to conductor  
 C. Perpendicular to the conductor and going inward  
 D. Surrounding the conductor and of circular nature
- Q.26 A charge  $q$  is moving with a velocity parallel to a magnetic field. Force on the charge due to magnetic field is
- A.  $qvB$   
 B. 0  
 C.  $qB/v$   
 D.  $Bv/q$


- Q.27 A straight wire of diameter 0.5 mm carrying current of 1A is replaced by another wire of 1 mm diameter carrying the same current. The strength of magnetic field far away is:
- A. Twice the earlier value  
B. One quarter of the earlier value  
C. One half of the earlier value  
D. No change
- Q.28 An electron accelerated through a potential difference  $V$ , passes through a uniform transverse magnetic field and experiences a force  $F$ . If the accelerating potential increased to  $2V$ , the electron in the same magnetic field will experience a force:
- A.  $F$   
B.  $\sqrt{2}F$   
C.  $F/2$   
D.  $2F$
- Q.29 The magnetic lines of force inside a bar magnet
- A. Are from north pole to south pole of the magnet  
B. Depend upon the area of cross-pole of the magnet  
C. Does not exist  
D. Are from south pole to north pole of the magnet
- Q.30 A charged particle moves in a uniform magnetic field. The path of the particle some instant makes an acute angle with the magnetic field. The path of the particle will be
- A. A straight line  
B. A helix with uniform speed  
C. A circle  
D. A helix with non-uniform speed
- Q.31 If  $F_1$  and  $F_2$  are forces acting on  $\alpha$ -particle and electron respectively, when moving perpendicular to the magnetic field then:
- A.  $F_1 = F_2$   
B.  $F_1 > F_2$   
C.  $F_1 < F_2$   
D.  $F_1 = 4F_2$
- Q.32 A uniform magnetic field of 3 G, exists in a  $+x$  direction. A proton shoots through the field in the  $+y$ -direction with a speed of  $5 \times 10^6$  m/s. The magnitude of the force on the proton is
- A.  $2.4 \times 10^{-16}$  N  
B.  $2.4 \times 10^{-6}$  N  
C.  $4 \times 10^{-16}$  N  
D. 4 N
- Q.33 20 Wb magnetic flux passes through the  $5\text{m}^2$  area of certain sheet, the magnetic flux density would be
- A.  $2 \text{ Wb m}^{-2}$   
B.  $4 \text{ Wb m}^{-2}$   
C.  $6 \text{ Wb m}^{-2}$   
D.  $8 \text{ Wb m}^{-2}$
- Q.34 The magnetic field in a certain region is  $5 \text{ Wb m}^{-2}$ . How much flux passes through an area of  $18\text{m}^2$ , if a loop is placed at right angle to the field
- A. 3.6Wb  
B. 90Wb  
C. 9.0 Wb  
D. 36Wb
- Q.35 When electrons are directed at right angle to a magnetic field directed into the plane of paper they will experience
- A. A variable force  
B. A constant force  
C. Force along the direction of velocity  
D. No force

- Q.36 G.P Thomson's early experiments on the diffraction of the electrons by crystals were criticized on the ground that the beams affecting the photographic plate might be X-rays. He proved that this was not so by placing bar magnets on each side of the beam as shown in the diagram.





How would the magnetic field due to magnetic affect the diffraction rings?


- A. The rings would be deflected in the direction A  
B. The rings would be deflected in the direction C  
C. The rings would be deflected in the direction B  
D. The rings would be deflected in the direction D
- Q.37 In a circuit with a coil of resistance 2 ohms, the magnetic flux changes from 2.0 Wb to 10.0 Wb in 0.2 second. The charge that flows in the coil during this time is
- A. 5.0 coulomb  
B. 4.0 coulomb  
C. 1.0 coulomb  
D. 0.8 coulomb
- Q.38 An aluminium ring B faces an electromagnet A. The current  $I$  through A can be altered



- A. Whether  $I$  increases or decreases, B will not experience any force  
B. If  $I$  decrease, A will repel B  
C. If  $I$  increases, A will attract B  
D. If  $I$  increases, A will repel B
- Q.39 A coil having an area  $2 \text{ m}^2$  is placed in a magnetic field which changes from  $1 \text{ Wb/m}^2$  to  $4 \text{ Wb/m}^2$  in an interval of 2 second. The e.m.f. induced in the coil will be
- A. 4 V  
B. 1.5 V  
C. 3 V  
D. 2 V
- Q.40 If the core of transformer is of substance whose hysteresis loop area is decreased, then the efficiency of transformer is
- A. Increased  
B. Decreased  
C. Same as original  
D. None of these is possible
- Q.41 A coil of wire is arranged with its plane perpendicular to a uniform magnetic field of flux density  $B$ , when the radius of the coil increases from  $r_1$  to  $r_2$  in time  $\Delta t$ , then what is the emf induced in the coil?
- A.  $\frac{\pi B (r_2^2 - r_1^2)}{\Delta t}$   
B.  $\frac{\pi B (r_2 - r_1)^2}{\Delta t}$   
C.  $\frac{B (r_2^2 - r_1^2)}{\Delta t}$   
D.  $\frac{\pi B (r_2^2 + r_1^2)}{\Delta t}$

- Q.42 The coils of a step down transformer have 500 and 5000 turns. In the primary coil an AC of 4 A at 2200 volts is sent. The value of the current and potential difference in the secondary will be.  
 A. 20 A, 22 V  
 B. 0.4 A, 22000 A  
 C. 40 A, 220 V  
 D. 40 A, 22000V
- Q.43 An alternating current or voltage \_\_\_\_\_  
 A. Fluctuates off and on  
 B. Varies in magnitude alone  
 C. Changes its direction again and again  
 D. Changes its magnitude continuously and reverses its direction of flow after regularly recurring intervals.
- Q.44 Lenz's law  
 A. Is the same as the right hand palm rule  
 B. Bears no relation to the law of conservation of energy  
 C. Determines the magnitude of an induced emf  
 D. Is useful in deciding the direction of an induced emf
- Q.45 A metal ring is held horizontally and a bar magnet is dropped through the ring, with its length along the axis of the ring. The acceleration of the falling magnet is:  
 A. Equal to g  
 B. More than g  
 C. Less than g  
 D. Depends on the diameter of the ring and length of magnet
- Q.46 A magnet is allowed to fall freely above a ring shaped metal which is having a cut in it as shown in the figure. When magnet fall towards the ring then  
 A. No emf will be induce in the ring  
 B. No current but some emf will be induced in ring  
 C. No current will be induce in the ring  
 D. No emf but some current will be induce in the ring
- Q.47 A.C voltage source changes its polarity \_\_\_\_\_ in one period  
 A. Once  
 B. Thrice  
 C. Twice  
 D. None
- Q.48 Refer to the figure maximum deflection in the galvanometer occurs when  
  
 A. The magnet is pushed toward the coil  
 B. The magnet is rotated in the coil  
 C. The magnet is stationary at the centre of coil  
 D. The number of turns in the coil is reduced
- Q.49 The emf induced in AC Generator is  $\epsilon$ . If the angular speed of the coil is triped, then the emf induced is  
 A.  $\epsilon$   
 B.  $3\epsilon$   
 C.  $2\epsilon$   
 D.  $4\epsilon$
- Q.50 A transformer has 200 turns in primary and 400 turns in secondary, if 50-watt power is given to input, what is output power?  
 A. 25 W  
 B. 50 W  
 C. 100 W  
 D. 200 W



- Q.51 A permanent magnet approaches a solenoid with a constant speed  $v$ . What is the magnetic pole induced at the portion Y of the solenoid and the direction of the induced current into the galvanometer?  
  
 A. 

Polarity at Y	Direction of the current
N-pole	From the left

 B. 

Polarity at Y	Direction of the current
N-pole	From the right

 C. 

Polarity at Y	Direction of the current
S-pole	From the left

 D. 

Polarity at Y	Direction of the current
S-pole	From the right
- Q.52 In a transformer 220 ac voltage is increased to 2200 volts. If the number of turns in the secondary are 2000, then the number of turns in the primary will be  
 A. 200  
 B. 50  
 C. 100  
 D. 20
- Q.53 A step-down transformer is connected to 2400 volts' line and 80 amperes of current is found to flow in output load. The ratio of the turns in primary and secondary coil is 20:1. If transformer efficiency is 100%, then the current flowing in primary coil will be  
 A. 1600 A  
 B. 4 A  
 C. 20 A  
 D. 1.5 A
- Q.54 The magnetic flux through a circuit of resistance R changes by an amount  $\Delta\phi$  in time  $\Delta t$ . Then the total quantity of electric charge q which passing during this time through any point of the circuit is given by  
 A.  $q = \frac{\Delta\phi}{\Delta t}$   
 B.  $q = -\frac{\Delta\phi}{\Delta t} + R$   
 C.  $q = \frac{\Delta\phi}{\Delta t} \times R$   
 D.  $q = \frac{\Delta\phi}{R}$
- Q.55 Primary secondary coils of a transformer have 50 and 200 turns respectively. When primary is connected to 9-volt battery secondary voltage is  
 A. 90  
 B. 18  
 C. 36  
 D. Zero
- Q.56 An electron is projected along the axis of a circular conductor carrying the same current. Electron will experience  
 A. A force along the axis  
 B. A force perpendicular to the axis  
 C. A force at an angle of  $4^\circ$  with axis  
 D. No force experienced
- Q.57 A coil having 500 sq. loops of side 10 cm is placed normal to magnetic flux which increases at a rate of 1 T/s. The induced emf is  
 A. 0.1 V  
 B. 0.5 V  
 C. 1 V  
 D. 5 V
- Q.58 North pole induces  
 A. Clockwise current  
 B. Anti-clockwise current  
 C. Zero current  
 D. Infinite current







- Q.59 Transformer ratings are given in  
 A. kVA  
 C. kVAR
- Q.60 Calculate the emf when a coil of 100 turns is subjected to a flux rate of 0.3 tesla/sec.  
 A. 3  
 C. -30
- B. HP  
 D. kW  
 B. 30  
 D. -300

## ANSWER KEY

1	A	11	A	21	C	31	B	41	A	51	C
2	B	12	D	22	A	32	A	42	C	52	A
3	C	13	A	23	B	33	B	43	D	53	B
4	A	14	C	24	B	34	B	44	D	54	D
5	C	15	B	25	D	35	B	45	C	55	D
6	D	16	D	26	B	36	A	46	B	56	D
7	C	17	A	27	D	37	B	47	A	57	D
8	A	18	A	28	B	38	D	48	A	58	B
9	D	19	C	29	D	39	C	49	B	59	A
10	B	20	A	30	B	40	A	50	B	60	B

- Q.1 In Compton scattering, the change in wavelength is maximum if  
 A. The angle of scattering is  $90^\circ$   
 C. The angle of scattering is  $180^\circ$
- Q.2 Calculate the energy of a photon of wavelength 6600 angstroms.  
 A.  $0.3 \times 10^{-19}$  J  
 C.  $30 \times 10^{-19}$  J
- Q.3 If the frequency of the incident radiation is equal to the threshold frequency, what will be the value of the stopping potential?  
 A. 0  
 C. 180 V
- Q.4 Which theory is confirmed by the Davisson - Germer experiment?  
 A. de - Broglie theory  
 C. Einstein's theory
- Q.5 The photoelectric effect can be understood on the basis of  
 A. Wave theory of light only  
 C. Quantum theory of light only
- Q.6 In the circuits given below, the correct full wave rectifier is \_\_\_\_\_



- Q.7 The momentum of a photon is  $p$ . The frequency associated with it is given by  
 A.  $pc/h$   
 C.  $ph/c$
- Q.8 There are  $n_1$  photons of frequency  $f_1$  in beam of light. In an equally energetic beam, there are  $n_2$  photons of frequency  $f_2$  then the correct relation is  
 A.  $\frac{n_1}{n_2} = 1$   
 C.  $\frac{n_1}{n_2} = \frac{f_1}{f_2}$
- B.  $\frac{h\nu}{p}$   
 D.  $\frac{h}{pc}$
- B.  $\frac{n_1}{n_2} = \frac{f_2}{f_1}$   
 D.  $\frac{n_1}{n_2} = \frac{f_1^2}{f_2^2}$

- Q.9 If we use two diodes and a centre tapped transformer, we will get  
 A. Half wave rectification B. A C current  
 C. Full wave rectification D. All of these
- Q.10 De-Broglie's hypothesis of wave nature of electrons was confirmed experimentally by:  
 A. Lummer and Pringsheim B. Davissan and Germer  
 C. Einstein and Max Planks D. Photoelectric equation
- Q.11 If an electron and a photon propagate in the form of waves having the same wavelength, it implies that they have the same  
 A. Velocity B. Energy  
 C. Angular momentum D. Momentum
- Q.12 The kinetic energy of electron and proton is  $10^{-32}$ J. Then the relation between their de-Broglie wavelengths is  
 A.  $\lambda_p = \lambda_e$  B.  $\lambda_p > \lambda_e$   
 C.  $\lambda_p < \lambda_e$  D.  $\lambda_p = 2\lambda_e$
- Q.13 If the energy of the photon is increased by a factor of 4, then its momentum  
 A. Does not change B. Increases by a factor of 4  
 C. Decreases by a factor of 4 D. Decreases by a factor of 2
- Q.14 Which light photon has the least momentum:  
 A. Red B. Blue  
 C. Yellow D. Green
- Q.15 De Broglie wave length associated with an electron at a speed of  $1 \times 10^6$  ms<sup>-1</sup>  
 A.  $7 \times 10^{-10}$ m B.  $6 \times 10^{-10}$ m  
 C.  $5 \times 10^{-10}$  m D.  $4 \times 10^{-10}$ m
- Q.16 Davission and Germer indicates \_\_\_\_\_ in their experiment.  
 A. Electron reflection B. Electron refraction  
 C. Electron polarization D. Electron diffraction
- Q.17 The electrons behave as waves because they can be:  
 A. Deflected by electric filed B. Deflected by magnetic field  
 C. Ionize as gas D. Diffracted by crystals
- Q.18 If the momentum of particle is doubled, then its de-Broglie wavelength:  
 A. Doubles B. Remain unchanged  
 C. Halves D. None of these
- Q.19 In Davison and Germer experiment, nickel crystal acts as a:  
 A. Perfect reflector B. Two dimensional grating  
 C. Perfect absorber D. Three dimensional grating
- Q.20 In Davison and Germer experiment, the angle which the incident beam makes with the normal to the nicked crystal is:  
 A.  $69^\circ$  B.  $90^\circ$   
 C.  $65^\circ$  D.  $180^\circ$
- Q.21 In a full wave rectifier input AC current has a frequency f, the output frequency of current is  
 A. 2f B. f/2  
 C. f D. 1.5 f
- Q.22 Of the following moving with same momentum, the one which has largest wavelength is:  
 A. An electron B. An  $\alpha$ -particle  
 C. A proton D. All have same de-Broglie wavelength


- Q.23 A particle of mass M at rest decays into two masses  $m_1$  and  $m_2$  with equal speed in opposite direction. The ratio of de-Broglie wave lengths of the particles  $\frac{\lambda_1}{\lambda_2}$  is  
 A.  $\frac{m_2}{m_1}$  B.  $\frac{\sqrt{m_1}}{\sqrt{m_2}}$   
 C.  $\frac{m_1}{m_2}$  D. 1 : 1
- Q.24 A transmitter radiates 30  $\mu$ W at 6.63 mm wavelength. The number of photons emitted per second are:  
 A.  $6.63 \times 10^{-34}$  B.  $6.63 \times 10^{34}$   
 C.  $10^{18}$  D.  $10^{11}$
- Q.25 Wave is associated with matter  
 A. When it is stationary  
 B. When it is in motion with the velocity of light only  
 C. When it is in motion with any velocity  
 D. None of these
- Q.26 The frequency of a photon, having energy 100 eV is ( $h=6.610^{-34}$ J-sec)  
 A.  $2.42 \times 10^{26}$ Hz B.  $2.42 \times 10^{12}$ Hz  
 C.  $2.42 \times 10^{16}$ Hz D.  $2.42 \times 10^9$ Hz
- Q.27 Which one is the correct expression of de Broglie equation for the wave length of atoms of mass m at temperature? T (k = Boltzmann's constant)  
 A.  $\lambda = \frac{h}{3mk}$  B.  $\frac{h}{\sqrt{3kTm}}$   
 C.  $\frac{h}{3kTm}$  D.  $\lambda = \frac{h}{\sqrt{3kT}}$
- Q.28 The ratio of momenta of an electron and an  $\alpha$ -particle which are accelerated from rest by a potential difference of 100 V is  
 A. 1 B.  $\sqrt{\frac{2m_e}{m_\alpha}}$   
 C.  $\sqrt{\frac{m_e}{m_\alpha}}$  D.  $\sqrt{\frac{m_e}{2m_\alpha}}$
- Q.29 In half wave rectifier if a resistance equal to load resistance is connected in parallel with the diode.  
 A. Output voltage would be halved B. Circuit will stop rectifying  
 C. Output voltage would be double D. Output voltage will remain unchanged
- Q.30 The PN junction diode is used as  
 A. A rectifier B. An oscillator  
 C. An amplifier D. A modulator
- Q.31 If n number of photon are striking on a metal surface, then total momentum exerted is \_\_\_\_\_  
 A.  $nh/\lambda$  B.  $2nh\lambda$   
 C. Zero D.  $n f \times t$

Unit- 9

- Q.32 The frequency of light beam A is twice that of light beam B. The ratio  $E_A/E_B$  of photon energies is  
 A. 1 B. 4  
 C. 1/2 D. 2
- Q.33 Which one of the following radiations has the strongest photon?  
 A. T.V waves B. Micro waves  
 C. X-rays D.  $\gamma$ -rays
- Q.34 In Davisson's and Germer's experiment if K.E of electron is increased by 4 times. Percentage change in  $\lambda$  will be:  
 A. 100% B. 400%  
 C. 200% D. 50%
- Q.35 In full wave rectification, the output D.C. voltage across the load is obtained for \_\_\_\_\_  
 A. The positive half cycle of input A.C. B. The negative half cycle of input A.C.  
 C. The complete cycle of input A.C. D. All of the above.
- Q.36 The output of a half wave rectifier is suitable only for  
 A. Running car radius B. Charging batteries  
 C. Running AC motors D. Running tape recorders
- Q.37 The primary function of a rectifier filter is to  
 A. Minimize a.c input variation  
 B. Stabilize d.c level of the output voltage  
 C. Suppress add harmonics in the rectifier output  
 D. Remove ripples, from the rectified output
- Q.38 The value of planck's constant can be determine by equation ( $E =$   
 A.  $h = \frac{E}{\lambda}$  B.  $h = \frac{Ec}{\lambda}$   
 C.  $h = \frac{E\lambda}{c}$  D.  $h = \frac{\lambda c}{E}$
- Q.39 An atomic particle of mass  $m$  moving at speed  $v$  is found to have wavelength  $\lambda$ . What is the wavelength of second particle with three times the speed and twice the mass?  
 A.  $\frac{3\lambda}{2}$  B.  $6\lambda$   
 C.  $\frac{2\lambda}{3}$  D.  $\frac{\lambda}{6}$
- Q.40 A proton, accelerated through a p.d  $V$  has a certain de Broglie wavelength. In order to have the same de Broglie wavelength, an  $\alpha$  - particle must be accelerated through a potential difference  
 A. 4V B.  $V/4$   
 C. 8V D.  $V/8$
- Q.41 The de-Broglie wavelength of the particle of mass  $m$  and energy  $E$  is  
 A.  $\lambda = \frac{h}{\sqrt{2mE}}$  B.  $\lambda = h\sqrt{2mE}$   
 C.  $\lambda = \frac{\sqrt{2mE}}{h}$  D.  $\lambda = \frac{1}{h\sqrt{2mE}}$

- Q.42 A proton and an  $\alpha$ -particle are accelerated through same voltage, the ratio of their de-Broglie wavelength will be  
 A. 1:2 B.  $2\sqrt{2}:1$   
 C.  $\sqrt{2}:1$  D. 2:1
- Q.43 If particles are moving with same velocity, then maximum de-Broglie wavelength will be for  
 A. Neutron B. Proton  
 C.  $\beta$ -particle D.  $\alpha$ -particle
- Q.44 An electron of mass  $m$  when accelerated through a potential difference  $V$  has de-Broglie wavelength  $\lambda$ . The de-Broglie wavelength associated with a proton of mass  $M$  accelerated through the same potential difference will be  
 A.  $\lambda \frac{m}{M}$  B.  $\lambda \frac{M}{m}$   
 C.  $\lambda \sqrt{\frac{m}{M}}$  D.  $\lambda \sqrt{\frac{M}{m}}$
- Q.45 Frequency of photon having energy 66 eV is  
 A.  $8 \times 10^{15} \text{ Hz}$  B.  $16 \times 10^{15} \text{ Hz}$   
 C.  $12 \times 10^{15} \text{ Hz}$  D. None of these
- Q.46 If the energy of the photon is increased by a factor of 4, then its momentum  
 A. Does not change B. Increases by a factor of 4  
 C. Decreases by a factor of 4 D. Decreases by a factor of 2
- Q.47 The energy of photon of radio waves is only  
 A.  $10^{-10} \text{ eV}$  B. 100 eV  
 C.  $10^{-5} \text{ eV}$  D. 1 MeV
- Q.48 If an electron is accelerated through a potential difference of 54 volts, its de-Broglie wavelength will be:  
 A.  $1.66 \times 10^{-8} \text{ m}$  B.  $1.66 \times 10^{-9} \text{ m}$   
 C.  $1.66 \times 10^{-10} \text{ m}$  D.  $1.66 \times 10^{-12} \text{ m}$
- Q.49 A particle which has zero rest mass and non-zero energy and momentum must travel with a speed  
 A. Equal to  $c$ , the speed of light in vacuum B. Less than  $c$   
 C. Greater than  $c$  D. Tending to infinity
- Q.50 Wavelength of a 1 keV photon is  $1.24 \times 10^{-9} \text{ m}$ . What is the frequency of 1 MeV photon?  
 A.  $1.24 \times 10^{15} \text{ Hz}$  B.  $1.24 \times 10^{18} \text{ Hz}$   
 C.  $2.4 \times 10^{20} \text{ Hz}$  D.  $2.4 \times 10^{23} \text{ Hz}$
- Q.51 The frequency of a photon, having energy 100 eV is ( $h=6.610^{-34} \text{ J-sec}$ )  
 A.  $2.42 \times 10^{20} \text{ Hz}$  B.  $2.42 \times 10^{12} \text{ Hz}$   
 C.  $2.42 \times 10^{16} \text{ Hz}$  D.  $2.42 \times 10^9 \text{ Hz}$
- Q.52 Energy of photon whose frequency is  $10^{12} \text{ MHz}$ , will be  
 A.  $4.14 \times 10^3 \text{ keV}$  B.  $4.14 \times 10^3 \text{ MeV}$   
 C.  $4.14 \times 10^2 \text{ eV}$  D.  $4.14 \times 10^3 \text{ eV}$
- Q.53 The momentum of photon is  
 A.  $\frac{h}{c}$  B.  $\frac{hc}{\lambda}$   
 C.  $\frac{h}{\lambda}$  D. Both B and C

- Q.54 If an electron and proton have the same de Broglie wavelength, which particle has greater speed?  
 A. Electron  
 B. Both have a same speed  
 C. Proton  
 D. None of these
- Q.55 Which photon have the maximum momentum  
 A. Green  
 B. Red  
 C. Blue  
 D. Have same momentum
- Q.56 Photons of energy 10.25 eV fall on the surface of the metal emitting photoelectrons of maximum kinetic energy 5.0 eV. What is the stopping voltage required for these electrons?  
 A. 10 V  
 B. 4 V  
 C. 8 V  
 D. 5 V
- Q.57 If the intensity of incident radiation in a photo-cell is increased, how does the stopping potential vary?  
 A. Increases  
 B. Remains the same  
 C. Decreases  
 D. Infinite
- Q.58 Which one of the following graphs represent correctly the variation of maximum kinetic energy  $E_{max}$  with the intensity of incident radiations having a constant frequency?



- Q.59 Half Wave diode rectifier circuit consists of  
 A. Single diode in Series with  $R_L$   
 B. Two diodes in Series with  $R_L$   
 C. Single diode in Parallel with  $R_L$   
 D. Two diodes in Parallel with  $R_L$
- Q.60 A simple diode rectifier has 'ripples' in the output wave which makes it unsuitable as a DC source. To overcome this one can use  
 A. A capacitor in series with the load resistance  
 B. A capacitor in parallel to the load resistance  
 C. Both of the mentioned situations will work  
 D. None of the mentioned situations will work

# ANSWER KEY

1	C	11	D	21	A	31	A	41	A	51	C
2	B	12	C	22	D	32	D	42	B	52	D
3	A	13	B	23	D	33	D	43	C	53	C
4	A	14	A	24	C	34	A	44	C	54	A
5	C	15	A	25	C	35	C	45	B	55	C
6	C	16	D	26	C	36	C	46	B	56	D
7	A	17	D	27	B	37	D	47	A	57	B
8	B	18	C	28	D	38	C	48	C	58	D
9	C	19	D	29	B	39	D	49	A	59	A
10	B	20	C	30	A	40	D	50	C	60	B

- Q.1 Gamma ( $\gamma$ ) radiation are fast moving  
 A. Electrons  
 B. Protons  
 C. Photons  
 D. Neutrons
- Q.2 If a radioactive element has a half-life of 40 minutes. The initial count rate was 1000 per minute, then how long will it take for the count rate to drop to 125 per minutes  
 A. 120 minutes  
 B. 90 minutes  
 C. 30 minutes  
 D. 60 minutes
- Q.3 The half-life of Carbon-14 is  
 A. 23 years  
 B. 1000 years  
 C. 1200 years  
 D. 5730 years
- Q.4 The activity of a sample of radioactive bismuth decreases to one-eighth of the original in 15 days. Its half-life is  
 A. 10 days  
 B. 15 days  
 C. 3 days  
 D. 5 days
- Q.5 The half-life of N is 6.5 s. A sample of this nuclide of hydrogen is observed for 32.5 s. The fraction of the original radioactive isotope remaining after this time is  
 A. 1/32  
 B. 1/16  
 C. 1/8  
 D. 1/4
- Q.6 If an electron jumps from 1st orbital to 3rd orbital, then it will  
 A. No gain of energy  
 B. Release energy  
 C. Absorb energy  
 D. None of these
- Q.7 An example of an absorption spectrum is the spectrum of:  
 A. Sodium vapour  
 B. Atomic hydrogen  
 C. Molten iron  
 D. Mercury vapour lamp
- Q.8 The spectral series that contains transitions terminating on the ground 1 hydrogen is called:  
 A. Paschen series  
 B. Balmer series  
 C. Pfund series  
 D. Lyman series
- Q.9 Paschen series is obtained when all the transitions of electron terminate on.  
 A. 2<sup>nd</sup> orbit  
 B. 4<sup>th</sup> orbit  
 C. 3<sup>rd</sup> orbit  
 D. 5<sup>th</sup> orbit
- Q.10 The relation for paschen series is given as  
 A.  $\frac{1}{\lambda} = R_H \left( \frac{1}{2^2} - \frac{1}{n^2} \right)$   
 B.  $\frac{1}{\lambda} = R_H \left( \frac{1}{4^2} - \frac{1}{n^2} \right)$   
 C.  $\frac{1}{\lambda} = R_H \left( \frac{1}{3^2} - \frac{1}{n^2} \right)$   
 D.  $\frac{1}{\lambda} = R_H \left( \frac{1}{5^2} - \frac{1}{n^2} \right)$
- Q.11 Isotopes are the atoms of the same element which contain equal number  
 A. Nucleons  
 B. Neutrons  
 C. Protons  
 D. Neutrons and protons

Q.12 What are the number of neutrons, protons and electrons in a neutral atom of  $^{235}_{92}\text{U}$ ?

	Number of neutrons	Number of protons	Number of electrons
A.	92	143	92
B.	143	92	235
C.	92	235	92
D.	235	92	


Q.13 Nucleus of an atom whose atomic mass is 24 consists of  
 A. 11 electrons, 11 protons and 13 neutrons  
 B. 11 protons and 13 neutrons  
 C. 11 electrons, 13 protons and 11 neutrons  
 D. 11 protons and 13 electrons

Q.14 As compared  $^{12}\text{C}$  atom,  $^{14}\text{C}$  atom has  
 A. Two extra protons and two extra electrons  
 B. Two extra neutrons and no extra electrons  
 C. Two extra protons but no extra electrons  
 D. Two extra neutrons and two extra electron

Q.15 Which one of the following pair is of isobars?  
 A.  $^{12}_6\text{C}$  and  $^{14}_6\text{C}$   
 B.  $^{14}_6\text{C}$  and  $^{14}_7\text{N}$   
 C.  $^4_2\text{He}$  and  $^3_1\text{H}$   
 D. Both A and C

Q.16 Diameter of nucleus is approximately \_\_\_\_\_  
 A.  $10^{-12}\text{m}$   
 B.  $10^{-10}\text{m}$   
 C.  $10^{-11}\text{m}$   
 D.  $10^{-14}\text{m}$

Q.17 A radioactive isotope 'W' decays to 'X' which decays to 'Y' and 'Y' decays to 'Z' as represented by the figure below




What is the change in the atomic number from 'W' to 'Z'?  
 A. Increase by 3  
 B. Increase by 5  
 C. Decrease by 3  
 D. Decrease by 5

Q.18 Consider a radioactive material of half-life 1 minute. If one of the nuclei decays now, the next one will decay  
 A. After one-minute  
 B. After any time  
 C. After two minutes  
 D. After  $\frac{1}{2}$  minute

Q.19 During a negative  $\beta$ -decay  
 A. An atomic electron is ejected  
 B. An electron which already present with in the nucleus is ejected  
 C. A neutron in the nucleus decays emitting an electron  
 D. A part of binding energy of nuclei is converted into electron

Q.20 Which of the arrangement about the particle is in accordance with the diagram?



- |    |          |         |          |          |
|----|----------|---------|----------|----------|
|    | P        | Q       | R        | Source   |
| A. | $\alpha$ | $\beta$ | $\gamma$ | S        |
| B. | $\alpha$ | $\beta$ | n        | $\gamma$ |
| C. | $\alpha$ | $\beta$ | n        | $\gamma$ |
| D. | $\gamma$ | n       | $\beta$  | $\alpha$ |

Q.21 When a radioactive nucleus emits a beta particle, the proton neutron ratio:  
 A. Decreases  
 B. Increases  
 C. Remains the same  
 D. None of the above

Q.22 Which statement concerning  $\alpha$ -particles, is correct?  
 A. An  $\alpha$ -particle has charge +4e  
 B. When  $\alpha$  particle travel through air, they cause ionization  
 C. An  $\alpha$ -particle is a helium atom  
 D. When  $\alpha$ -particle travel through a sheet of gold foil, they make the gold radioactive

Q.23 The decay of a nucleus of neptunium is accompanied by the emission of a  $\beta$ -particle and  $\gamma$ -radiation. What effect (if any) does this decay have on the proton number and the nucleon number of the nucleus?

	proton number	nucleon number
A.	increases	decreases
B.	unchanged	decreases
C.	decreases	increases
D.	increases	unchanged

Q.24 The transformation of a neutron into proton in the nucleus gives rise to emission of:  
 A. Beta particles  
 B. Gamma particles  
 C. Alpha particles  
 D. X-rays

Q.25 In the reaction  $^{234}_{92}\text{Th} \rightarrow ^{234}_{93}\text{Y} + ^0_{-1}\text{e}$ , the electron emits from the  
 A. 1<sup>st</sup> orbit  
 B. Nucleus  
 C. 2<sup>nd</sup> orbit  
 D. Valence Shell

Q.26 According to the equation  $^4_2\text{X} \rightarrow \text{Y} + 3 \alpha$ -particles, what are the atomic and mass numbers of 'Y'?  
 A. Z - 6, A - 12  
 B. Z + 1, A  
 C. Z - 2, A - 4  
 D. Z + 3, A


Q.27 The value of A in the following reaction is  
 $^9_4\text{Be} + ^4_2\text{He} = ^A_6\text{C} + ^1_0\text{n}$   
 A. 14  
 B. 10  
 C. 12  
 D. 16

Q.28 In a radioactive series,  $^{238}_{92}\text{U}$  changes to  $^{206}_{82}\text{Pb}$  through  $n_1$   $\alpha$ -decay processes and  $n_2$   $\beta$ -decay processes.  
 A.  $n_1 = 8, n_2 = 8$   
 B.  $n_1 = 8, n_2 = 6$   
 C.  $n_1 = 6, n_2 = 6$   
 D.  $n_1 = 6, n_2 = 8$

- Q.29 The most penetrating radiations out of the following is that of  
 A.  $\gamma$ -rays  
 B.  $\alpha$ -particles  
 C.  $\beta$ -rays  
 D. X-rays
- Q.30 Two radioactive elements X and Y have half-lives of 25 minutes and 75 minutes respectively. Sample of X and Y initially contain equal numbers of atoms. After 150 minutes what is the value of the following fraction?  

$$\frac{\text{No. of nuclei of X unchanged}}{\text{No. of nuclei of Y unchanged}}$$
  
 A. 1 : 16  
 B. 16 : 1  
 C. 1 : 8  
 D. 8 : 1
- Q.31 In a radioactive substance at  $t = 0$ , the number of atoms is  $8 \times 10^4$ . Its half-life period is 3 years. The number of atoms  $1 \times 10^4$  will remain after interval  
 A. 9 years  
 B. 8 years  
 C. 6 years  
 D. 24 years
- Q.32 If the decay or disintegration constant of a radioactive substance is  $\lambda$ , then its half-life and mean life are respectively  
 A.  $\frac{1}{\lambda}$  and  $\frac{\log_e 2}{\lambda}$   
 B.  $\frac{\log_e 2}{\lambda}$  and  $\frac{1}{\lambda}$   
 C.  $\lambda \log_e 2$  and  $\frac{1}{\lambda}$   
 D.  $\frac{\lambda}{\log_e 2}$  and  $\frac{1}{\lambda}$
- Q.33 In a sample of radioactive material, what fraction of the material will decay after half of its half-life.  
 A.  $\frac{1}{\sqrt{2}}$   
 B.  $\sqrt{2} - 1$   
 C.  $\frac{\sqrt{2} - 1}{\sqrt{2}}$   
 D.  $\sqrt{2}$
- Q.34 The decay constant  $\lambda$  of a radioactive sample  
 A. Decrease as the age of atoms increase  
 B. Increase as the age of atoms increase  
 C. Is independent of the age  
 D. Depends on the nature of activity
- Q.35 A radioactive decay rate of radioactive elements is found to be  $10^3$  disintegrations per sec at a certain time. If the half-life of the element is 1 second, the decay rate after one second is \_\_\_\_\_ and after 3 second is \_\_\_\_\_  
 A. 500, 125  
 B. 125, 500  
 C.  $10^3$ ,  $10^3$   
 D. 100, 10
- Q.36 If the radioactive decay constant of radium is  $1.07 \times 10^{-4}$  per year, then its half-life period is approximately equal to  
 A. 8,900 years  
 B. 6,476 years  
 C. 7,000 years  
 D. 2,520 years
- Q.37 Half-life of radioactive element depends upon  
 A. Amount of element present  
 B. Temperature  
 C. Pressure  
 D. None
- Q.38 A radioactive substance has a half-life of 60 minutes. After 3 hours, the fraction of atom that have decayed would be  
 A. 12.5%  
 B. 8.5%  
 C. 87.5%  
 D. 25.1%

- Q.39 Half-life of radium is 1600 years. Which of the following is the fraction of a sample of radium that would remain un-decayed after 6400 years?  
 A.  $\frac{1}{2}$   
 B.  $\frac{1}{4}$   
 C.  $\frac{1}{8}$   
 D.  $\frac{1}{16}$
- Q.40 A given radioactive sample is reduced from 20 g to 1.25 g in 40 days. Its half-life would be  
 A. 10 days  
 B. 8 days  
 C. 5 days  
 D. 6 days
- Q.41 A radioactive substance has a half-life of four months, three-fourths of the substance will decay in:  
 A. 3 months  
 B. 4 months  
 C. 8 months  
 D. 12 months
- Q.42 Calculate the half-life of bismuth-214 which has a decay constant of  $4.3 \times 10^3 \text{ s}^{-1}$ .  
 A.  $3.9 \times 10^3 \text{ s}$   
 B.  $2.9 \times 10^{-3} \text{ s}$   
 C.  $2.9 \times 10^3 \text{ s}$   
 D.  $1.6 \times 10^{-4} \text{ s}$
- Q.43 The radioactivity of a certain radioactive element drops to  $\frac{1}{64}$  of its initial value in 30 second. Its half-life is:  
 A. 2 second  
 B. 5 second  
 C. 4 second  
 D. 6 second
- Q.44 Cobalt-57 is radioactive, emitting  $\beta$  particles. The half-life for this is 270 days. If 100 mg of this is kept in an open container, the mass of Cobalt-57 after 540 days will be:  
 A. 50 mg  
 B.  $\frac{50}{\sqrt{2}}$  mg  
 C. 25 mg  
 D. Zero
- Q.45 When thorium ( ${}_{90}\text{Th}^{234}$ ) emits a  $\beta$ -particle its proton to neutron ratio would become  
 A. 90/144  
 B. 91/144  
 C. 91/143  
 D. 90/143
- Q.46 Three paths of radioactive radiations are observed as shown in the figure in the presence of electric field. Which type of radiation is shown in path 1?



- A. Alpha  
 B. Gamma  
 C. Beta  
 D. Cathode rays
- Q.47 In radiotherapy X-rays are used to:  
 A. Detect bone fractures  
 B. Detect heart diseases  
 C. Treat cancer by controlled exposure  
 D. Detect fault in radio receiving circuits
- Q.48 Radioactive iodine can be used to check person's \_\_\_\_\_ is working properly  
 A. Cancer  
 B. Lungs  
 C. Skin cancer  
 D. Thyroid gland

- Q.49 Phosphorus is used as tracer in  
 A. Industries  
 C. Agriculture  
 B. Electrical machinery  
 D. All of these
- Q.50 Cobalt-60 is used for treatment of  
 A. Cancer  
 C. Lungs  
 B. Kidneys  
 D. Thyroid
- Q.51 The  $\gamma$  - rays radiographs are used in:  
 A. Agriculture industry  
 C. Support Industry  
 B. Medical industry  
 D. All of above
- Q.52 In heavy elements ( $Z = \text{No. of protons}, N = \text{No. of neutrons}$ )  
 A.  $Z = N$   
 C.  $Z > N$   
 B.  $Z < N$   
 D.  $Z + N = A$
- Q.53 A nucleus emits a radiation neither its mass no. nor charge no. change the radiation is  
 A.  $\alpha$   
 C.  $\beta$   
 B.  $\gamma$   
 D. Neutron
- Q.54 An atom emits some radiation such that daughter nucleus is isotope of parent nucleus, the emitted radiation is  
 A.  $\alpha$   
 C.  $\beta$   
 B.  $\gamma$   
 D.  $1\alpha$  and  $2\beta$
- Q.55 In radioactivity  $\beta$  particles originate from  
 A. K-shell electron  
 C. Preexisting electron in nucleus  
 B. The decay of neutron  
 D. The decay of proton
- Q.56 CFC is used in  
 A. Refrigeration  
 C. Plastic foam industry  
 B. Aerosol spray  
 D. All of above
- Q.57 The age of fossil when C-14: C-12 in bone is one fourth of ratio in bone of living animal and half-life of C-14 is 5732 years is  
 A. 100 years  
 C. 1000 years  
 B. 11460 years  
 D. 1200 years
- Q.58 Technium-99 has a half-life of 6 hours. If there is 300 mg of it, how much will be left in 48 hours?  
 A. 1.58 mg  
 C. 1.17 mg  
 B. 1.25 mg  
 D. 2.56 mg
- Q.59 What is the ratio of minimum to maximum wavelength in the Balmer series?  
 A. 5:9  
 C. 1:4  
 B. 5:36  
 D. 3:4
- Q.60 When a hydrogen atom is in its first excited level, what is the relation of radius and Bohr radius?  
 A. Twice  
 C. Same  
 B. 4 times  
 D. Half

## ANSWER KEY

1	C	11	C	21	B	31	A	41	C	51	B
2	A	12	B	22	B	32	B	42	D	52	B
3	D	13	B	23	D	33	C	43	B	53	B
4	D	14	B	24	A	34	C	44	C	54	D
5	A	15	B	25	B	35	A	45	C	55	B
6	C	16	D	26	A	36	B	46	B	56	A
7	B	17	C	27	C	37	D	47	C	57	B
8	D	18	B	28	B	38	C	48	D	58	C
9	C	19	C	29	A	39	D	49	C	59	A
10	C	20	A	30	A	40	A	50	A	60	B


# POST-PREP ASSESSMENT

Note: Physics paper of NMDCAT contains 56 questions.

- Q.1 A battery is used to charge a parallel plate capacitor till the potential difference between the plates becomes equal to the electromotive force of the battery. The ratio of the energy stored in the capacitor and the work done by the battery will be
- A. 1  
B. 2  
C.  $\frac{1}{4}$   
D.  $\frac{1}{2}$

- Q.2 In gamma ray emission from a nucleus
- A. Both the neutron number and the proton number change  
B. There is no change in the proton number and the neutron number  
C. Only the neutron number changes  
D. Only the proton number changes

- Q.3 Charges are placed on the vertices of a square as shown. Let E be the electric field and V the potential at the centre. If the charges on A and B are interchanged with those on D and C respectively, then



- A.  $\vec{E}$  remain unchanged, V changes  
B. both  $\vec{E}$  and V change  
C.  $\vec{E}$  and V remains unchanged  
D.  $\vec{E}$  changes, V remains unchanged

- Q.4 A block of mass 'm' is connected to another block of mass 'M' by a spring (massless) of spring constant 'k'. The blocks are kept on a smooth horizontal plane. Initially the block are at rest and the spring is unstretched. Then a constant force 'F' starts acting on the block of mass 'M' to pull it. Find the force on the block of mass 'm'


- A.  $\frac{mF}{M}$   
B.  $\frac{(M+m)F}{M}$   
C.  $\frac{mF}{(m+M)}$   
D.  $\frac{MF}{(m+M)}$

- Q.5 A charged particle moves through a magnetic field perpendicular to its direction. Then
- A. The momentum changes but the kinetic energy is constant  
B. Both momentum and kinetic energy of the particle are not constant  
C. Both, momentum and kinetic energy of the particle are constant  
D. Kinetic energy changes but the momentum is constant

- Q.6 The resistance of a wire is 5 ohm at 50°C and 6 ohm at 100°C. The resistance of the wire at 0°C will be
- A. 2 ohm  
B. 1 ohm  
C. 4 ohm  
D. 3 ohm


- Q.7 A particle is projected at 60° to the horizontal with a kinetic energy K. The kinetic energy at the highest point is
- A. K  
B. Zero  
C. K/2  
D. K/4

- Q.8 Two springs, of force constants  $k_1$  and  $k_2$  are connected to a mass m as shown. The frequency of oscillation of the mass is f. If both  $k_1$  and  $k_2$  are made four times their original values, the frequency of oscillation becomes



- A.  $f/2$   
B.  $f/4$   
C. 4f  
D. 2f

- Q.9 The graph of range of projectile (on y-axis) as plotted against velocity, of projectile (x-axis) will be



- Q.10 A body is moving in a circle with period of 20π. What is angular frequency of body
- A. 0.1 Hz  
B. 100 Hz  
C. 2 Hz  
D. 10π Hz

- Q.11 A body is revolving with a constant speed along a circle. If its direction of motion reversed but the speed remains the same, then which of the following statement is true
- A. The centripetal force will not suffer any change in magnitude  
B. The centripetal force will have its direction reversed  
C. The centripetal force will not suffer any change in direction  
D. The centripetal force would be doubled

- Q.12 The angular velocity of the minute hand of a clock is

- A.  $\frac{2\pi}{60} \text{ rads}^{-1}$   
B.  $\frac{\pi}{24} \text{ rads}^{-1}$   
C.  $\frac{2\pi}{3600} \text{ rads}^{-1}$   
D.  $\frac{\pi}{3600} \text{ rads}^{-1}$

- Q.13 A simple harmonic oscillator has a time period of 10 seconds. Which equation relates acceleration a and displacement x?

- A.  $a = -10x$   
B.  $a = -(20\pi)^2x$   
C.  $a = -(20\pi)x$   
D.  $a = -(2\pi/10)^2x$

- Q.14 In stationary wave of wavelength 0.4 m the distance between node and antinodes is

- A. 0.1 m  
B. 0.4 m  
C. 0.2 m  
D. 0.5 m



Q.15 The minimum length of tube open at both ends that resonates with frequency of 350 Hz if  $v = 350 \text{ m s}^{-1}$ .

- A. 1 m
- C. 2 m

- B.  $\frac{1}{2}$  m
- D.  $\frac{1}{14}$  m

Q.16 The ratio of angular frequency and linear frequency is

- A.  $2\pi$
- C.  $\frac{1}{2\pi}$

- B.  $\pi$
- D.  $\frac{\pi}{2}$

Q.17 How many electrons per second constitute a current of one micro ampere?

- A. One electron
- C.  $10^{-6}$  electrons

- B.  $10^6$  electrons
- D.  $6.25 \times 10^{12}$  electrons

Q.18 A sound source is moving towards a stationary observer with  $\frac{1}{10}$  of the speed of sound.

- A.  $\frac{10}{9}$
- C.  $\left(\frac{11}{10}\right)^2$

- B.  $\frac{11}{10}$
- D.  $\left(\frac{9}{10}\right)^2$

Q.19 When we double the voltage in a simple electric circuits we double the

- A. Current
- C. Power

- B. Resistance
- D. Both A. and C.

Q.20 Which relation exactly described the isothermal process

- A.  $Q = W$
- C.  $Q = -\Delta U$

- B.  $W = -\Delta U$
- D.  $Q = \Delta U + W$

Q.21 The work done in moving a unit positive charge from one point to another against the electric field is a measure of

- A. Capacitance
- C. Intensity of electric field

- B. Potential difference between two points
- D. Resistance between two points

Q.22 Energy density in case of a capacitor is always proportional to

- A.  $E^2$
- C.  $V^2$


- B.  $\epsilon_0$
- D. C

Q.23 The capacitance of a parallel plate capacitor is given by

- A.  $C = \frac{A}{\epsilon_0 d}$
- C.  $C = \frac{A\epsilon_0}{d}$

- B.  $C = \frac{\epsilon_0 d}{A}$
- D.  $C = \frac{d}{\epsilon_0 A}$

Q.24 What's the total work performed on the gas as it's transformed from state a to state c, along the path indicated?



- A. 1,500 J
- C. 4,500 J

- B. 3,000 J
- D. 5,000 J

Q.25 The charge of an electron is  $1.6 \times 10^{-19} \text{ C}$ . How many electrons strike the screen of cathode ray tube each second when the beam current is 16 mA

- A.  $10^{19}$
- C.  $10^{15}$

- B.  $10^{17}$
- D.  $10^{17}$

Q.26 The following four wires are made of the same material and are at the same temperature. Which one of them has highest electrical resistance

- A. length = 100 cm, diameter = 1 mm
- C. length = 200 cm, diameter = 2 mm

- B. length = 50 cm, diameter = 0.5 mm
- D. length = 300 cm, diameter = 3 mm

Q.27 In a circuit with a coil of resistance 2 ohms, the magnetic flux changes from 2.0 Wb to 10.0 Wb in 0.2 second. The charge that flows in the coil during this time is

- A. 5.0 coulomb
- C. 4.0 coulomb

- B. 1.6 coulomb
- D. 0.8 coulomb

Q.28 The radiations emitted from hydrogen filled discharge tube show

- A. band spectrum
- C. continuous spectrum

- B. line spectrum
- D. none

Q.29 A radio station emits 10 kW power of 90.8 MHz. Find the number of photon emitted per second

- A.  $1.6 \times 10^{23}$
- C.  $1.6 \times 10^{27}$

- B.  $1.6 \times 10^{25}$
- D.  $1.6 \times 10^{22}$

Q.30 The number of turns in the primary and secondary coils of a transformer are 100 and 300 respectively. If the input power is 60 W. The output power will be

- A. 100 W
- C. 60 W

- B.  $3 \times 10^3 \text{ W}$
- D. 90 W

Q.31 In Davisson-Germer experiment, the diffracted electron beam from crystal shows

- A. Particle property
- C. Light property

- B. Wave property
- D. Quantum property

Q.32 The Lenz's law refers to induce

- A. emf
- C. Current

- B. Resistance
- D. Capacitance

Q.33 When a current carrying conductor is placed in a magnetic field. It moves from a region of

- A. Stronger to weak field
- C. Strong to weak if current is large

- B. Weak to strong field
- D. Weak to strong if current is large

Q.34 In electromagnetic induction, the induced e.m.f. in a coil is independent of

- A. Resistance of the circuit
- C. Change in the flux

- B. Time
- D. None

Q.35 In full wave rectifier with input frequency 50Hz the ripple in the output is mainly of frequency:

- A. 25 Hz
- C. 100 Hz

- B. 50 Hz
- D. zero

Q.36 What percentage of original radioactive atoms is left after five half lives

- A. 0.3%
- C. 31%

- B. 1%
- D. 3.125%

Q.37 A 2 MeV proton is moving perpendicular to a uniform magnetic field of 2.5 T. The force on the proton is:

- A.  $2.5 \times 10^{-10} \text{ N}$
- C.  $8 \times 10^{-11} \text{ N}$


- B.  $2.5 \times 10^{-11} \text{ N}$
- D.  $8 \times 10^{-12} \text{ N}$

Q.38 Unit of decay constant is

- A. s
- C. m

- B.  $\text{m}^{-1}$
- D.  $\text{s}^{-1}$

- Q.39 A certain radioactive mass decays from 64 gm to 2 gm in 20 days. What is its half-life?  
 A. 5 days B. 4 days  
 C. 3 days D. 5 days
- Q.40 An electron makes transition from  $n = 4$  to  $n = 1$  state in a hydrogen atom. The maximum possible number of photons emitted will be  
 A. 1 B. 3  
 C. 2 D. 6
- Q.41 A 60 kg man pushes a 40 kg man by a force of 60 N. The 40 kg man has pushed the other man with a force of:  
 A. 40 N B. 0  
 C. 60 N D. 20 N
- Q.42 The speed of a vehicle of mass 500kg increases from  $30\text{ms}^{-1}$  to  $70\text{ms}^{-1}$ . Calculate the work.  
 A.  $10^2$  B.  $10^3$   
 C.  $10^6$  D.  $10^9$
- Q.43 Work done by the gravitational force on a body of mass "m" moving on a smooth horizontal surface through a distance "s" is: (Given acceleration due to gravity = g)  
 A. mgs B. -mgs  
 C. 0 D. 2mgs
- Q.44 A plane is revolving around the earth with a speed of 100 km/hr at a constant height from the surface of earth. The change in the velocity as it travels half circle is  
 A. 200 km/hr B. 150 km/hr  
 C. 100 km/hr D. 0
- Q.45 Doppler effect is independent of \_\_\_\_\_  
 A. Velocity of source B. Velocity of listener  
 C. distance between source and observer D. None of the above
- Q.46 The following figure depicts a wave travelling in a medium. Which pairs of particles are in phase?



- A. B & F  
 C. A & D  
 B. C & E  
 D. B & G
- Q.47 First law of thermodynamics states that:  
 A. System can do work B. System has temperature  
 C. System has pressure D. Heat is a form of energy
- Q.48 A hollow metal sphere of radius 5 cm is charged so that the potential on its surface is 10 V. The potential at the center of the sphere is  
 A. 10 V B. Same as at point 5 cm away from the surface  
 C. 0 V D. Same as at point 25 cm away from the surface
- Q.49 A conducting resistance is connected to the battery and temperature of conductor decreases by the process of cooling then the value of current will be  
 A. Increased B. Decreased  
 C. Remain constant D. Zero

- Q.50 When a bar magnet is broken into two pieces  
 A. We will have a single pole on each piece B. Each piece will have two unlike poles  
 C. Each piece will have two like poles D. None of these
- Q.51 Whenever there is a change in the magnetic flux linked with a closed circuit, an emf and a current are induced in the circuit. This statement is referred to as;  
 A. Lenz's law  
 B. Faraday's second law of electromagnetic induction  
 C. Faraday's first law of electromagnetic induction  
 D. Laplace's law
- Q.52 A transformer:  
 A. Transforms energy B. Transforms frequency  
 C. Transforms voltage D. None
- Q.53 The wavelengths of a proton and a photon are same. Then:  
 A. Their velocities are same B. Their momenta are equal  
 C. Their energies are same D. Their speeds are same.
- Q.54 Bohr's model can explain:  
 A. The spectrum of hydrogen atom only  
 B. Spectrum of an atom or ion containing one electron only  
 C. The spectrum of hydrogen molecule  
 D. The solar spectrum
- Q.55 Radioactivity is affected by:  
 A. Temperature B. Pressure  
 C. Electric and magnetic field D. None of these
- Q.56 In alternating current  
 A. The direction of current is always positive  
 B. The direction of current is always negative  
 C. The direction of current changes constantly  
 D. The direction of current is either positive or negative

## ANSWER KEY

1	D	11	A	21	B	31	B	41	C	51	C
2	B	12	C	22	A	32	C	42	C	52	C
3	D	13	D	23	C	33	A	43	C	53	B
4	C	14	A	24	C	34	A	44	A	54	B
5	A	15	B	25	B	35	C	45	C	55	D
6	C	16	A	26	B	36	D	46	D	56	C
7	C	17	D	27	C	37	D	47	A		
8	D	18	A	28	B	38	D	48	A		
9	D	19	A	29	C	39	B	49	A		
10	A	20	A	30	C	40	D	50	B		

# EXPLANATORY NOTES

Q.1  $\frac{1}{2} qv = \frac{1}{2} qv$

Q.2 In gamma ray emission from a nucleus there is no change in the proton number and the neutron number

Q.3 As  $\vec{E}$  is a vector quantity.

Q.4  $Kx = ma = \frac{mF}{m+M}$

Q.5 A charged particle moves through a magnetic field perpendicular to its direction. Then the momentum changes but the kinetic energy is constant

Q.6

$$\frac{5}{6} = \frac{(1+50\alpha)}{(1+100\alpha)}$$

$$\alpha = \frac{1}{200}$$

$$5 = R_o(1+50\alpha)$$

$$R_o = 4$$

Q.7

$$K = K \cos^2 Q$$

$$= K \cos^2 60$$

$$= K \left(\frac{1}{2}\right)$$

Q.8

$$f = \frac{1}{2\pi} \sqrt{\frac{k_1 + K_2}{m}}$$

$$f' = \frac{1}{2\pi} \sqrt{\frac{k_1 + K_2}{m}} = 2f$$

Q.9  $R = \frac{v_i^2 \sin 2\theta}{g} \rightarrow R \propto v_i^2$  (Parabola along y-axis)

Q.10  $\omega = \frac{2\pi}{T} = \frac{2\pi}{20\pi} = 0.1$

Q.11  $\vec{f}_c = m\omega^2(-\vec{r})$  the magnitude of the centripetal force remains constant but its direction is continuously changing.

Q.12  $\theta = \omega t$   
 $\omega = \frac{\theta}{t} = \frac{2\pi}{3600} = \text{rads}^{-1}$

Q.13  $a = -x\omega^2$   
 $a = -x \left(\frac{2\pi}{T}\right)^2 = -x \left(\frac{2\pi}{10}\right)^2$

Q.14 Distance between node and antinode is  $= \frac{\lambda}{4} = \frac{0.4}{4} = 0.1$

Q.15  $v = f\lambda \rightarrow \lambda = \frac{v}{f} = \frac{350}{350} = 1$

For open pipe  $\lambda_1 = 2l \rightarrow l = \frac{1}{2} \text{m}$

Q.16  $\frac{\omega}{f} = \frac{2\pi f}{f} = 2\pi$

Q.17  $\frac{n}{t} = \frac{I}{e} \Rightarrow$  Number of free electrons per second  $= \frac{10^{-6}}{1.6 \times 10^{-19}} = 6.25 \times 10^{12}$  electrons / sec

Q.18  $f' = \left(\frac{v}{v-u_s}\right) f \Rightarrow f' = \left(\frac{v}{v-\frac{v}{10}}\right) f$

$$\frac{f'}{f} = \frac{v}{\frac{9}{10}v} = \frac{10}{9}$$

Q.19 According to Ohm's law  $I \propto V$

If voltage is doubled then current will also be doubled, so power will be four times as  $P \propto V^2$  also  $P \propto I^2$

Q.20 In isothermal process  $\Delta U = 0$

$$Q = \Delta U + W$$

$$Q = W$$

Q.21  $V = \frac{W}{q}$

Q.22  $U = \frac{1}{2} \epsilon_0 \epsilon_r E^2 \Rightarrow U \propto E^2$

Q.23  $C = \frac{\Delta E_0}{d}$

Q.24

$$W = P\Delta V = P(V_2 - V_1)$$

$$W = (3 \times 10^3)(25 - 10) \times 10^{-3} = 4500 \text{ J}$$

Q.25  $i = \frac{ne}{t} \Rightarrow 16 \times 10^{-3} = \frac{n \times 1.6 \times 10^{-19}}{1} \Rightarrow n = 10^{17}$

Q.26  $R \propto \frac{l}{r^2}$ . For highest resistance  $\frac{l}{r^2}$  should be maximum, which is correct for option

Q.27  $\epsilon = \frac{\Delta\phi}{\Delta t} \Rightarrow IR = \frac{\Delta\phi}{\Delta t} \Rightarrow \frac{Q}{\Delta t} = \frac{\Delta\phi}{R\Delta t}$

$$\Rightarrow Q = \frac{\Delta\phi}{R} = \frac{10 - 2}{2} = 4 \text{ C}$$

Q.28 Gas or vapor state of elements shows line spectrum

Q.29  $E = nhf \Rightarrow \frac{E}{t} = \frac{n}{t} hf \Rightarrow \frac{n}{t} = \frac{P}{hf} = 1.6 \times 10^{29}$

Q.30  $P_{in} = P_{out}$  [Ideal transformer]  $V_p I_p = V_s I_s$

Q.31 Diffracted electron beam from crystal shows wave nature.

Q.32 Lenz's law refers to induce current.

Q.33 Current carrying conductor will move from stronger to weaker magnetic field.

Q.34  $\epsilon = N \frac{\Delta\phi}{\Delta t}$

Q.35 In full wave frequency of the output is  $2f$

Q.36 Percentage fractional undecay  $= \frac{1}{2^n} \times 100 = \frac{1}{2^5} \times 100 = \frac{1}{32} \times 100 = 3.125\%$

Q.37  $K.E = \frac{1}{2}mv^2 \Rightarrow v = \sqrt{\frac{2K.E}{m}}$  ;  $F = qvB$

Q.38  $\lambda = \frac{\Delta N / N_0}{\Delta t}$   $\therefore$  unit  $= s^{-1}$

Q.39 Remaining atoms  $= \frac{\text{Original atoms}}{2^n}$

$$2^n = \frac{64}{2} = 32$$

$$n = 5$$

5 half-lives in 20 days

$$\text{So, half-life} = \frac{20}{5} = 4 \text{ days}$$

Q.40 Number of spectral lines  $= \frac{n(n-1)}{2} = \frac{4(4-1)}{2} = 6$

Q.41 To every action, there is an equal and opposite reaction. So when 60 kg man exerts a force of 60 N on 40 kg man, 40 kg man also exerts the same force of 60 N on the 60 kg man.

Q.42  $K.E_i = 1/2 \times 500 \times 30^2 = 250 \times 900 = 225000 \text{ J}$

$$K.E_f = 1/2 \times 500 \times 70^2 = 250 \times 4900 = 1225000 \text{ J}$$

$$\text{Work} = \text{Change in Kinetic energy} = K.E_f - K.E_i = 1225000 - 225000 = 1000000 = 10^6 \text{ J}$$

Q.43 As the gravitational force is perpendicular to the distance moved by body, the work done by gravity will be zero.

Q.44 As we know,

$$\Delta v = 2v \sin(\theta/2) = 2 \times v \times \sin 90^\circ$$

$$= 2 \times 100 = 200 \text{ km/hr}$$

Q.45 Doppler effect is independent of distance between source and observer

Q.46 Path difference between B and G is  $\lambda$ . so they are in the same phase.

Q.47  $\Delta Q = \Delta U + \Delta W$

Q.48 Since potential inside the hollow sphere is same as that on the surface.

Q.49 The resistance of the conducting wire will decrease according to equation  $R_t = R_0 (1 \pm \alpha \Delta t)$  due to which current will increase according to ohm's law

Q.50 When we break a magnet into two, both act as magnets, that is both have a north and south pole each. In both these pieces the corresponding north and south poles remain on the sides as in the original magnet. This means that the two broken sides must get oppositely polarized.

Q.51 Faraday's first law of electromagnetic induction state that whenever a conductor are placed in a varying magnetic field emf are induced which is called induced emf, if the conductor circuit are closed current are also induced which is called induced current.

Q.52 Transformers are capable of either increasing or decreasing the voltage and current levels of their supply, without modifying its frequency, or the amount of Electrical Power being transferred from one winding to another via the magnetic circuit.

Q.53 For Proton,  $\lambda = \frac{h}{mv} \rightarrow \lambda_1 = \frac{h}{p}$

for photon,  $E = \frac{hc}{\lambda}$ ,  $\lambda_2 = \frac{h}{E/c}$

$$\lambda_1 = \lambda_2 \Rightarrow P = \frac{E}{c}$$

Q.54 Bohr's model is valid to atom or ion having only one electron such as H, He<sup>+</sup>, Li<sup>2+</sup>, Be<sup>3+</sup>, etc

Q.55 Temperature, pressure, electric and magnetic field does not affect Radioactivity.

Q.56 Electric charge in alternating current changes direction periodically. The voltage in AC circuits also periodically reverses because the current changes direction.